

SKY and TELESCOPE

Spain 1960
Indiana University
1960
151960



Eclipse camp by moonlight

In This Issue:

★
Vol. XX, No. 5
NOVEMBER, 1960
50 cents
★

The Fine Structure
of Solar Prominences — I
Amateurs Photograph
the Lunar Eclipse
The Problem of Cygnus A

American Astronomers Report
Lunik II's Landing on the Moon
Convention at Haverford
Lick 120-inch Photographs — V

TEACHING WITH A FECKER 38" REFLECTOR



COMMENTS BY HARRY E. CRULL

*Director
J. I. Holcomb Observatory,
Butler University*

“The 38-inch reflector of the J. I. Holcomb Observatory at Butler University provides potentially the opportunity for utilization of a variety of accessories. The f/4 38-inch primary is large enough to have adequate light-gathering power and yet is of a size that can be conveniently mounted and easily manipulated. The Cassegrain optical system places the eyepiece near the center of motion, and the easily installed plate holder is a flexible accessory. The rugged and well balanced mounting is ideally suited to the addition of photometer, spectrograph, or other attachments. The 6-inch refracting guide telescope of long focus gives easy and accurate assistance in use of the instrument. Installation of a Newtonian diagonal and eyepiece or plate holder could be accomplished with a minimum of difficulty. The telescope has performed well for both public nights and class instruction over the past six years.”



This 38" Reflector Telescope was designed and built by J. W. Fecker for the J. I. Holcomb Observatory at Butler University. It is used for: student instruction, photography and visual use.

ACCESSORIES: 3" aperture wide-angle finder telescope, 6" aperture guide telescope, zenith eyepiece adapter.

FEATURES: fork mounting, sidereal drive, sidereal, hour and declination circles, solenoid clamping and motor driven slow motions in right ascension and declination.

For more information about this scope or other telescopes to meet your specific needs, write

j. w. fecker

Division of American Optical Co.
6592 Hamilton Ave., Pittsburgh 6, Pa.



COVER: The site of the September 5th lunar-eclipse expedition of the Tucson Astronomical and Astronautical Association to the Tucson Mountains, in Arizona. The picture was taken at 2:15 a.m. Mountain standard time, by Donald J. Strittmatter. The 13 members operated seven telescopes, ranging from two 8-inch reflectors to a 3-inch refractor. (See page 256.)

FRANK E. ROSS	251
THE FINE STRUCTURE OF SOLAR PROMINENCES - I — Donald H. Menzel and John G. Wolbach	252
AMATEURS PHOTOGRAPH THE LUNAR ECLIPSE	256
THE PROBLEM OF CYGNUS A — Otto Struve	259
AMERICAN ASTRONOMERS REPORT	263
LUNIK II'S LANDING ON THE MOON	265
CONVENTION AT HAVERFORD — William K. Hartmann	266
LICK 120-INCH PHOTOGRAPHS - V	269
AMATEUR ASTRONOMERS	272
A High School Observatory in California	
BOOKS AND THE SKY	294
Arabische Sternnamen in Europa	
The Search for Order	
Physics and Medicine of the Atmosphere and Space	
CELESTIAL CALENDAR	310
The Eclipsing Variable Star RZ Cassiopeiae	
GETTING ACQUAINTED WITH ASTRONOMY	277
The Planets — Mars - I	
GLEANINGS FOR ATM's	302
A Precision-Mounted Small Refractor	
An Altazimuth Reflector with a Drive	
LETTERS	262
NEWS NOTES	271
OBSERVER'S PAGE	282
The September Partial Eclipse of the Sun	
Lunar Rills in Eastern Mare Tranquillitatis	
Three Current Comets	
Deep-Sky Wonders	
OBSERVING THE SATELLITES	274
PLANETARIUM NOTES	280
QUESTIONS	278
STARS FOR NOVEMBER	313

FEATURE PICTURE: The nebulosity associated with the open star cluster M16 in Serpens, taken July 20, 1960, with the 120-inch reflector. Lick Observatory photograph. 268

SKY AND TELESCOPE is published monthly by Sky Publishing Corporation, Harvard College Observatory, Cambridge 38, Mass. Second-class postage paid at Boston, Mass.

Subscription: \$5.00 per year in the United States and possessions; \$9.00 for two years; \$13.00 for three years. To Canada, Mexico, and all countries of the Pan American Postal Union, \$6.00 for one year; \$11.00 for two years; \$16.00 for three years. To all other foreign countries, \$7.00 for one year; \$13.00 for two years; \$19.00 for three years. Canadian and foreign remittances should be made in United States currency. Single copy 50 cents. Back numbers, as available, 50 cents each.

All notices of change of address must be sent one month in advance and accompanied by old and new addresses. When sending your renewal order, or writing in regard to your subscription, your current mailing address must be given. Return our bill form with renewal payment. Circulation manager, Nancy R. Bolton. Office staff: Virginia K. Cox, J. Caroline Nason, Mary Silva, John Simmons, Helen B. Sniegiecki, Stacia Strong.

Editorial and advertising offices: 49-50-51 Bay State Rd., Cambridge 38, Mass. Advertising rates are listed in STANDARD RATE AND DATA SERVICE, or sent upon request. Advertising manager, Henry M. Corrado. Unsolicited articles and pictures are welcome, but we cannot guarantee prompt editorial attention, nor are we responsible for the return of unsolicited material. Editorial assistants: Gail B. Carney, William E. Shawcross, Raymond N. Watts, Jr.

The principal articles are indexed in THE READERS' GUIDE TO PERIODICAL LITERATURE.

Although the contents of this magazine, its text, pictures, diagrams, and charts, are copyrighted with all rights reserved, we are pleased to give permission, upon request, for the use of SKY AND TELESCOPE material for educational and scientific purposes, with the exception of the monthly star and planet charts, which may not be copied or reproduced in any manner for any purpose. Permission to use matter from signed articles, and photographs and diagrams for which credit is given, must also be obtained from the person, observatory, institution, or agency concerned.

Frank E. Ross

ONE of the most versatile of all American astronomers was Frank E. Ross, who died September 21st at the age of 86. Today he is perhaps first remembered for the large-field lenses he developed for precise astronomical photography, and for the correcting lenses he invented to be used with great reflectors to enlarge the field of sharp definition.

Optical design was, however, only one of Dr. Ross' careers. Trained in celestial mechanics by Armin Leuschner at the University of California, in 1902 he joined the Nautical Almanac Office in Washington, D. C. There he collaborated with Simon Newcomb in a monumental study of the moon's motion. The *American Ephemeris* still bases its predicted positions of the planet Mars and of Saturn's satellite Phoebe upon orbits determined by Dr. Ross.

From 1905 to 1915 he was in charge of the International Latitude Observatory at Gaithersburg, Maryland, and then spent a decade as a physicist with Eastman Kodak Co., finally joining the staff of Yerkes Observatory in 1924. There he prepared with Mary Calvert a magnificent photographic atlas of the Milky Way, and discovered many hundreds of variables and stars with large proper motions.

For the last 30 years of his life, Dr. Ross was also connected with Mount Wilson and Palomar Observatories, and served as a consultant in the design of the 200-inch Hale telescope. At Mount Wilson he secured a famous series of photographs, through color filters, of the planet Venus; these long remained the high-water mark in the observational attack on the nature of that planet.

Precise photographic photometry was also a favorite subject of his, and led to the invention of the Ross microphotometer. He collaborated at Mount Wilson with F. H. Seares and Mary Joyner in preparing a large catalogue of accurate magnitudes of stars north of declination +80°.

In the range of astronomical fields to which he made first-rate contributions, few other American astronomers can be compared to Frank E. Ross.

J. A.

Sky and Telescope Binders

A dark blue fabrikoid binder priced at \$3.50 each, postpaid in the United States; \$4.00 in Canada. Two sizes: Binder C is for volumes up to XVIII; Binder D is for volume XIX and after. When ordering, please specify the volume number for which the binder is to be used.

Your name can be gold-stamped for 75¢ extra, the volume number for 50¢, both for \$1.25; print desired lettering clearly. Payment must accompany order. (Sorry, but no foreign orders accepted.)

Sky Publishing Corporation

Harvard Observatory, Cambridge 38, Mass.




Fig. 1. These loops of glowing gas, photographed on the limb of the sun by the Sacramento Peak 15-inch prominence camera, are solar prominences of class ASI (see table on page 254). The diagram below identifies particular details for discussion. All pictures with this article are from Harvard and Sacramento Peak observatories, courtesy U.S. Air Force.

The Fine Structure of Solar Prominences—I

DONALD H. MENZEL and JOHN G. WOLBACH, *Harvard College Observatory*




Fig. 1 (key). The loop prominence shown above and sketched here may be seen in its entirety in Fig. 2. The bright luminous structure at A is an excellent example of a filament resolved into thin parallel threads. At B, neighboring fine features brighten at contiguous places along the filaments. Some of the elongated, linked clouds that the authors liken to sausages are seen at C.

UNTIL the year 1868, astronomers could observe the red clouds of glowing hydrogen at the sun's edge — the solar prominences — only during total eclipses. Then N. Lockyer and J. Janssen independently discovered a spectroscopic technique for visual observations of these formations at any time. The modern era in the study of prominences began with B. Lyot's invention of the coronagraph (1931) and the introduction shortly afterward of optical filters with very narrow passbands.

The 6-inch coronagraph and the 15-inch prominence camera of Sacramento Peak Observatory and instruments of other solar stations have provided detailed mo-




Fig. 2. In these remarkable ASI-type loops, hydrogen gas flows down each side of the nearly circular trajectory like water escaping from a piece of hose open at both ends. A number of bright knots or condensations are in the gas streams at D. The brighter filaments are partly overexposed in this 6-inch coronagraph picture taken on June 28, 1957.

tion pictures of prominence activity. These hydrogen-alpha photographs show clearly that prominences are not uniform clouds of gas, but frequently possess an intricate filamentary structure of various degrees of complexity.

The simplest form consists of one or more filaments of matter, usually moving downward from the corona. Few of these trajectories appear to be uniformly luminous; they generally display, as in Figs. 1 and 2, an irregular pattern of knots or condensations closely following one another along each filament. In size, the individual knots of gas are comparable to the earth or the moon.

As the photographs show, the knots are not necessarily round, but often elongated. These individual "dashes" sometimes lie parallel to the filament along the direction of motion and resemble a string of moving sausages. Less frequently the dashes lie perpendicular to the filament and the direction of motion





Fig. 3. This October 9, 1957, photograph obtained with the 15-inch prominence camera shows rain of class ASa. Several distinct flow patterns arise in various coronal sources, E. The sausagelike formations that prevail generally in this region contrast markedly with the isolated examples of nearly uniform streamers at F. Near G are a number of fine threads, both parallel and perpendicular to the general downward motion of the material. Examples of fishbone structure are at lower right.



(Figs. 3 and 4), and their pattern may be referred to as fishbone.

Under highest magnification, a filament occasionally is resolved into a series of fine, parallel threads with either a sausage or fishbone structure superposed. Fig. 1 shows threads of this type. Such detailed examination reveals the varying complexity of prominences, ranging from simple threads to multiple and interwoven or tangled filaments that form a vertical curtain (Fig. 5).


A single prominence may simultaneously exhibit sausagelike structure in one region and fishbone in another. We do not know why prominences show so wide a range of structural form, and we place no special emphasis on these details. Our desire is to emphasize how intricate the filamentary or fibrous structure can be. We also wish to show that some prominences possess diffuse or hazy filaments,

while others have sharply defined fibers, with diameters less than the effective resolving power of the instrument.

To illustrate the variations of internal appearance, we have selected a variety of typical prominences from the voluminous available observational material. For each, we give the classification on a system devised by D. H. Menzel and J. W. Evans, as modified by F. Q. Orrall.

To describe both behavior and appearance of prominences, the system first divides them into two classes: A, in which the luminous matter comes from above, and B, in which it comes from below. Each class is then separated into two groups consisting of prominences associated only with sunspots, subclass S, and prominences not so associated, subclass N. The table shows the full system.

An additional behavioral description is necessary, because in the course of its de-





PROMINENCE CLASSIFICATION

A (Descending)

S (Spot)	N (Non-spot)
a. rain	a. coronal rain
f. funnel	b. tree trunk
l. loop	c. tree
	d. hedgerow
	f. suspended cloud
	m. mound

B (Ascending)

S (Spot)	N (Non-spot)
s. surge	s. spicule
p. puff	

writers term such behavior "eruptive," as distinguished from "quiescent," a category to which most class A prominences were once assigned. Actually the behavior of these prominences does not justify the use of either older term.

Matter continually filters through an A prominence, becoming luminous at the top and descending slowly with approximately uniform velocity until it flows from the bottom. The speed of descent usually ranges between 10 and 20 kilometers per second, so that the substance






Fig. 4 (above and left). Fishbone patterns occur where matter breaks from the funnel-shaped cloud at H, and moves toward L, in the direction of the arrow. At the same time, the bright knots lengthen across this direction (I). Luminous material cascades in graceful arcs to the surface at J, while a transition from complex to simple patterns occurs at K. This ANf formation occurred July 14, 1957.

development a prominence of class A may suddenly ascend. At some stage of their development, many hedgerow prominences display heightened activity. The rate of downward flow increases markedly. Then the matter suddenly elevates to form an expanding arch, giving the illusion that such prominences belong to class B rather than A. Indeed, many

Fig. 5 (below and right). Resolution of this ANd hedgerow into filaments is evident. The dominant motion is downward, with unusual upward flow at M. Fishbone arrangement occurs at N. Three days after this prominence was photographed, it appeared as a tree, ANc, then vanished.






Fig. 6. A sketch to illustrate the Menzel model of a structureless prominence. The heavy horizontal lines represent the solar surface, while lighter ones indicate the magnetic field in the sun's atmosphere.

of a large hedgerow prominence some 150,000 kilometers high undergoes a complete change about once every three hours. Meanwhile, the appearance of the prominence generally remains substantially the same. As a result, we conclude that the physical conditions and the force fields responsible for the shape and distribution of the gases possess a singular degree of permanence.

With a spectroheliograph or a Lyot filter we can observe hedgerows on the face of the sun itself. In $H\alpha$ light they appear as dark filaments against the bright background of the photosphere, because they absorb more light than they emit along the line of sight. Frequently, they persist for several solar rotations of approximately 27 days each.

Some years ago, one of us (Menzel) proposed a physical model for a structureless prominence, in terms of a solution of the equations of magnetohydrostatics. To understand this model and apply it to real prominences, consider the following elementary situation.

Imagine a magnetic field embedded in the hot corona, with lines of force parallel to the solar surface, as in Fig. 6a. Now suppose that, in the region indicated as O between the dotted lines, some disturbance causes the corona to lose energy rapidly, perhaps through radiation in the far ultraviolet. The gas cools, contracts, and subsides, carrying the magnetic lines of force with it, as shown in 6b, wherein P represents a prominence supported by a balance of gas pressure and magnetic forces. The prominence is supposed to extend a relatively large distance above and below the plane of the paper.

It is commonly believed that ionized gas can move only along — not across — magnetic lines of force. This statement requires some modification. The natural motion does tend to be along the field. However, motion may occur across the field, and then the lines of force are stretched and distorted as in 6c. When the displacement perpendicular to the field is excessive, in the sense that it exceeds the width of the disturbance,

second-order forces come into play which may cause bubbles of ionized gas to break through the lines of force, each bubble containing some of the original magnetic field.

There will thus be a tendency for a column of bubbles to form, these in turn tending to elongate in the direction of the field, under the compression of both magnetic and hydrostatic forces (Fig. 6d). An alternative form is in 6e, where the lines of force are heavily stretched downward and the successive bubbles of gas are themselves compressed by the effect of the overlapping layers as the matter slowly descends along the vertical column.


The rupture of the field may occur at intervals, resulting in parallel columns of gas bubbles. Figs. 7 and 8 illustrate the rib structure formed as successive layers of matter condense and leak through the lower layers of the magnetic field. Thus, this simple model appears to account for the major features of downward-moving prominences.

(To be continued)



Fig. 7 (left). This type-ANf suspended cloud was photographed on October 9, 1959, with the 15-inch camera at Sacramento Peak. Material is flowing predominantly downward and filtering slowly from the bottom of the cloud. The arrow at Q shows the direction of motion of the material as it flows across the axis of the fishbone structure, parallel to the magnetic field.

Fig. 8 (right). The suspended cloud seen in Fig. 7 is shown about three hours later, when the seeing at Sunspot, New Mexico, had deteriorated markedly. The bubbles of luminous gas at R now flow downward, indicated by the arrow. The ribbed pattern due to multiple ruptures of the field may be noted in both pictures.





J. Russell Smith secured these pictures of the eclipse of the moon on September 5th with the 16-inch Newtonian reflector of his Skyview Observatory in Eagle Pass, Texas. The prints are contrasty, so in each case the penumbral darkening along the edge of the umbra is emphasized. From left to right, the exposure times on Tri-X film are: 1/100 second at 9:31 Universal time, 1/50 at 9:59, and 1/50 at 10:14. The first picture was taken while the moon was in the penumbral shadow, five minutes before predicted umbral contact. All times given with this article are UT.

Amateurs Photograph the Lunar Eclipse

ON September 5th, generally fair weather favored most amateur observers of the total eclipse of the moon, although in the United States large areas of the East and Midwest were clouded out. Reports and pictures have been received from places as widely separated as Rhode Island, Hawaii, British Columbia, Costa Rica, and Australia. On these pages are some samples of the many

excellent photographs that were taken.

Activities on eclipse night ranged from the backyard vigil of the lone amateur to large eclipse parties held by societies, and many groups made expeditions to country sites far from city lights (see front cover).

As the moon entered deep in the earth's shadow and the brightness of a full-moon night faded away, a spectacular auroral display became visible during totality


for watchers in many places. One observer in British Columbia described it as "often so bright that I thought dawn was coming." This display was seen as far south as Amarillo, Texas.

The December issue of *SKY AND TELESCOPE* will contain a synopsis of the visual observations of the eclipse, and a listing of the names of all persons who submitted reports or pictures of the event.




Eight-inch reflectors and eyepiece projection were used for both of these pictures. At the left, penumbral darkening is recorded at 9:28, eight minutes before umbral eclipse began. Alan L. Larson, of Burlington, Iowa, took this 1/300-second exposure on Tri-X film, through a low-power eyepiece. Below is seen the moon at 12:15, 10 minutes after totality ended. Harold J. Kendis, Jr., Los Angeles, California, used a special lens system to obtain an effective focal length of 19.5 feet, the moon's image on the negative being over two inches in diameter.






Above: Eight stages in the disappearance of the moon into the earth's shadow were recorded on a single negative by Richard Cromwell at Des Moines, Iowa. The first exposure, at upper left, was made at 9:36, at the moment of umbral contact, and shows strong penumbral darkening. Subsequent exposures were made at 10-minute intervals until two minutes before the moon was completely engulfed at totality.

Right: The series above was taken with a Mamiya C camera attached to the homemade 6-inch Springfield reflector that Mr. Cromwell is operating at the right. The telescope tube (mottled gray) is supported by a simple but sturdy pipe mounting that is firmly embedded in the ground. Tom Morehead, at the left, adjusts the eyepiece of the 10-inch portable reflector (white tube) that was used for visual observations. Mr. Cromwell took this self-photograph by pulling a black thread attached to the shutter mechanism of his camera.




At San Carlos, California, Victor Nikolashin took pictures of the moon while it was in the penumbra, or light outer shadow of the earth, just before the umbral phase (left) and while it was emerging from the umbra (right). The 1/25-second exposures were secured at the Newtonian focus of a 6-inch f/6 homemade reflector. The first picture, at 9:36 (umbral contact), was on a 103a-O spectroscopic plate developed for four minutes in D-19 to bring out the penumbral shading. The other, at 12:45, was on Polaroid film with an ASA rating of 200, developed in D-11 for about five minutes. As with all other photographs in this article, the moon is shown uninverted, with north at the top.





Above: Thirteen members of the observing section of the Tucson Astronomical and Astronautical Association watched the eclipse at Gates Pass picnic area, 10 miles west of the Arizona city (see the front-cover photograph). This series of moon shots by group leader Donald J. Strittmatter shows the moon in the umbral shadow at 10-minute intervals, beginning at 10:06 and ending two minutes before totality. The image formed by his 8-inch reflector was projected through a 32-mm. Erfle eyepiece and recorded by a Bronica single-lens reflex camera on Royal-X Pan film. The large dark sea at the right is Mare Crisium, with Mare Tranquillitatis to its left and Mare Foecunditatis below.

Left: For one year, Eddie Ries, a 14-year-old high school sophomore from Pico Rivera, California, has taken about 250 pictures of the moon, including a record of the entire lunar cycle, for his science fair project. The eclipse photograph here was made at 12:53 on Royal Pan film, with his 6-inch reflector at 60x. Tycho is prominent at the lower left; Plato is the dark crater at the upper right.



Part of the equipment set up by Fred Larsen of Los Angeles, California, for viewing the lunar eclipse, as well as the eclipse of the sun on September 20th (see page 282). His main instrument, at the left, is an 8-inch Cassegrainian used for close-ups of the last part of the lunar eclipse. At the right, an old De Vrey camera was attached to a 5-inch achromatic telescope for color movies, taking one picture per second. The moon's image almost filled the 16-mm. frame.

The Problem of Cygnus A

OTTO STRUVE

*National Radio Astronomy
Observatory**

IN 1944, Grote Reber's pioneer radio map gave a first rough view of the northern heavens in radiation of 1.9 meters wave length. This historic chart (reproduced on page 191 last month) showed a broad band of emission stretching across the sky, corresponding in position with the Milky Way. Superimposed on the band were three strong concentrations in the constellations Sagittarius, Cassiopeia, and Cygnus, that are now known as three of the most intense discrete radio sources in the sky.

The first of these, Sagittarius A, coincides with the center of our galaxy. F. D. Drake has recently resolved it into four symmetrically placed components. The inner two are clouds of hot hydrogen gas, while the outer pair probably represent a gaseous ring, whose plane coincides with the plane of symmetry of our galaxy, the ring radiating by the synchrotron process. The source Cassiopeia A, which I discussed last month, is probably the remnant of a supernova that exploded about 260 years ago.

Cygnus A remained unidentified with any optically observable feature until, nine years ago, F. G. Smith at Cambridge, England, accurately measured its coordinates as $19^{\text{h}} 57^{\text{m}} 45^{\text{s}}.3$, $+40^{\circ} 35'.0$

*Operated by the Associated Universities, Inc., under contract with the National Science Foundation.



Fig. 1. This greatly enlarged portion of a photograph taken with the 200-inch shows the peculiar object — possibly a pair of colliding galaxies — in the position of the cosmic radio source Cygnus A. The very faint halo is larger than this field. Mount Wilson and Palomar Observatories photograph.

(1950). In this same spot, W. Baade and R. Minkowski photographed a peculiar 18th-magnitude galaxy, evidently the source of the radio emission.

This faint object has an abnormal appearance, consisting of a pair of distinct condensations about two seconds of arc apart, and surrounded by a large, dim halo (Fig. 1). The spectrum of the galaxy is also remarkable. Superimposed on the faint continuous background from the hosts of member stars there are many diffuse emission lines, attributed to forbidden transitions (Fig. 2). According to W. A. Baum, more than half of the visible light of the object is produced by emission lines. In this respect, the Cygnus galaxy differs radically from ordinary

systems, such as M31 or the Milky Way, whose spectra exhibit weak emission lines and a relatively strong continuum.

Baade and Minkowski, in 1954, explained the pair of condensations as two separate galaxies in the act of colliding nearly face on. Their motion relative to each other would make an angle of about 45 degrees to the line of sight. Earlier, in 1951, Baade and L. Spitzer, Jr., had computed the chances of collisions between galaxies. The probability that a particular field galaxy will collide with some other one is extremely small. In dense clusters of galaxies, however, such encounters should be much more frequent.

Thus, when Baade and Minkowski discovered that the Cygnus A object is actually one of the brightest members of a remote, rich cluster of galaxies, it seemed not unreasonable to suggest that it is a pair of galaxies undergoing a collision. Since each of the two systems may be of the order of 100,000 light-years in diameter, and the relative velocity perhaps 3,000 kilometers per second, it might easily take something like a million years for one galaxy to pass through the other.

"As far as the stars of the colliding systems are concerned," the Palomar astronomers wrote in 1954, "such a collision is an absolutely harmless affair. The average distance between two stars is so large that the two galaxies penetrate each other without any stellar collisions. The situa-




Fig. 2. The spectrum of Cygnus A, recorded by W. Baade and R. Minkowski with the 200-inch telescope. Very bright lines of ionized oxygen, nitrogen, and neon make this spectrum quite abnormal for a galaxy. The Palomar astronomers used a grating spectrograph, giving overlapping orders. From the "Astrophysical Journal."

tion is very different for the gas and dust embedded in the two systems. Because of the much shorter free paths of the gas and dust particles, the collision of the two galaxies means a real collision of the imbedded gas and dust, which are heated up to very high temperatures, since the collisional velocities range from hundreds to thousands of kilometers per second."

The hypothesis of colliding galaxies as strong radio sources has since been the subject of many investigations, and until recently most astronomers adopted it. Some doubt arose when radio astronomers observed that the centimeter and meter emission of Cygnus A does not come from the pair of condensations, but from two large blobs, located about 80 seconds of arc (some 250,000 light-years) to either side of the optical object. No visible light is associated with these blobs. The optical feature itself has a longest diameter of only about 30 seconds of arc (90,000 light-years).

In November, 1959, Minkowski rediscovered the problem of Cygnus A when the




Fig. 3. Successive stages in the collision of two galaxies, as visualized by I. S. Shklovsky in 1954. As the galaxies interpenetrate, the gas and dust clouds recoil. From "Cosmic Radio Waves."

National Academy of Sciences met at Indiana University. There he stated: "This difference of optical and radio appearance does not contradict the identification. . . . [It has] now become strikingly clear from unpublished investigations by Bolton of the source Centaurus A, whose identification with NGC 5128 cannot be doubted." In both cases, there are optically unobservable sources of radio emission, symmetrically placed with respect to the optical object. For Centaurus A, the two radio blobs are each about four degrees from NGC 5128, or approximately $1\frac{1}{2}$ million light-years.

The Soviet radio astronomer I. S. Shklovsky had accepted the colliding-galaxy hypothesis in his book, *Cosmic Radio Waves*, whose English translation, containing its author's revisions, was published earlier this year. He gives an important analysis of the phenomena that would result from the encounter of two galaxies, each surrounded by a spherical halo of exceedingly tenuous but hot gas. When the outer edges of the halos



Fig. 4. The intense southern radio source Centaurus A is associated with this unusual elliptical galaxy, NGC 5128. Minkowski has called this two galaxies in collision, interpreting the dark band as a spiral seen edge on. Shklovsky, however, thinks it a single galaxy with an obscuring belt, and suggests that remote Cygnus A would appear similar, if it could be viewed from nearby. Hale telescope photograph from Mount Wilson and Palomar Observatories.

clashed, a hydrodynamic wave would be propagated rapidly through the gas, causing the appearance of many high-energy charged particles (Fig. 3). Their velocities, in the case of electrons, would approach the speed of light. Such so-called relativistic electrons produce strong radio waves.

In his book, Shklovsky calculated from the assumed density of the halos and their relative velocity that the kinetic energy of the Cygnus A colliding gas masses is 5×10^{50} ergs. The radiation rate is approximately 2×10^{41} ergs per second — most of it as radio emission. He supposed that the collision has been going on for about 10 million years, which would mean that some 5×10^{38} ergs have already been radiated away, approximately 10 per cent of the available kinetic energy.

All this seemed reasonably consistent with the state of our knowledge in early 1960. However, in an article about to appear in the *Russian Astronomical Journal*, Shklovsky has changed his position and now proposes a new interpretation of Cygnus A. I am grateful to Professor Shklovsky for an advance copy of his manuscript.

He and other astronomers had felt concern about the identification of several radio sources, notably Centaurus A, which Baade and Minkowski had linked with NGC 5128. That galaxy is near enough for detailed photographs (Fig. 4). A few years ago, it was widely regarded as a pair of colliding galaxies — a spherical one and an edgewise spiral. The former was identified with the symmetrical bright body, the latter with the broad band of obscuration. This interpretation was, however, never considered fully convincing, despite Minkowski's defense of it in November, 1959.

Additional doubt was created in Shklovsky's mind by the magnificent recent interferometer work on Centaurus A by J. Bolton and his associates. Their radio mapping indicates that Centaurus A consists of at least three sources — the galaxy NGC 5128 itself and the two distant blobs (*Publications, Astronomical Society of the Pacific*, February, 1960, page 29). And Shklovsky did not know that C. M. Wade, at the National Radio Astronomy Observatory, had recently discovered a new radio source, not far from Virgo A, that seems from preliminary data to be a close counterpart of Centaurus A. Optically, Wade's source is also an elliptical galaxy with a band of obscuration across it. If anything, this discovery tends to augment the doubts expressed by the Soviet astronomer.

During the past few years, several other strange galaxies have been found to be sources of strong radio emission (for example NGC 4486, associated with Virgo A), but the radio emission does not come from the brightest part of the optical feature. NGC 4486 shows no indication of collision; instead, from its nucleus a




Fig. 5. From the nucleus of NGC 4486 (M87) extends a beaded ray that is highly polarized. These Hale telescope views, taken in polarized light of four orientations (electric vector marked), differ in appearance.

luminous straight jet extends in position angle 290 degrees (Fig. 5). Photographs of this jet show several condensations whose light is strongly polarized, each condensation having a different angle of polarization.

In his new paper, Shklovsky has marshaled several arguments against the collision hypothesis for Cygnus A. Consider first the problem of the energy involved. If the radio emission of Cygnus A is produced by the synchrotron mechanism, the total energy contained in the relativistic electrons is about 5×10^{50} ergs, a result based on independent computations by Shklovsky and G. R. Burbidge.

Where can this enormous amount of energy come from? The optically observed galaxies may contain 200 billion solar-type stars, with a combined mass of 4×10^{44} grams. The corresponding nuclear energy is given by Einstein's famous formula as the product of the total mass by the square of the velocity of light — 4×10^{58} ergs. However, only about one per cent of this energy can be released by known nuclear processes, unless we assume that the complete annihilation of mass and its conversion into energy are occurring in galaxies. And of this one per cent, only a minute fraction would appear as kinetic energy, with very little of that becoming radio emission.

As a second argument, Shklovsky points out that the optical spectrum of the Cygnus A object does not show double emission lines, as might be expected in the case of colliding galaxies. Instead of the relative velocity of 3,000 kilometers per

second he assumed in his book, he now argues with considerable emphasis that the spectra by Baade and Minkowski suggest a relative velocity of only some 100 or 200 kilometers per second. Since the kinetic energy is proportional to the square of the relative velocity, it becomes perhaps 400 times smaller than the 5×10^{50} ergs he had given in the book. If so, the kinetic energy of a collision would be insufficient to maintain the radio emission of Cygnus A over the past 10 million years.

A third line of argument follows from Shklovsky's belief that each of the two galaxies of Cygnus A has an absolute magnitude of the order of -21 . They would thus be giant systems, at least as large and as massive as the Milky Way and M31. Shklovsky therefore agrees with V. A. Ambarzumian that in this case the probability of collision as computed by Baade and Spitzer does not apply. The two American astronomers had considered average galaxies, but even in as dense a cluster as the Cygnus one, the probability of observing at the present time a collision between two extremely large galaxies is vanishingly small, because such objects are relatively rare.

These and other discrepancies have led Shklovsky to a new interpretation of Cygnus A. He now suggests that the apparently double optical object is not two colliding galaxies, but is a peculiar single galaxy resembling NGC 5128, with a broad band of obscuration across it.

If this view is correct, then Cygnus A and Centaurus A are very much alike,

but differing greatly in distance, since the first is 700 million light-years away, the other only 20 million. The two systems also differ in evolutionary age, according to Shklovsky, Cygnus A being intrinsically a much more powerful emitter of radio waves. From the synchrotron mechanism he estimates that Cygnus A may be only a few tens of millions of years old, while the age of Centaurus A may be between one and three billion years.

The source of the radio emission of both galaxies is attributed to tenuous gas clouds ejected from supernovae that exploded during the early stages of the two systems. The supernovae would have occurred in their central regions, where the stars are very strongly concentrated. The ejected clouds of gas, moving along magnetic lines of force from the nucleus of each galaxy toward its poles, would produce large numbers of the relativistic electrons that give rise to synchrotron radiation.

Shklovsky suggests that all galaxies undergo this stage of evolution relatively early in their histories. If so, some five or 10 billion years ago our Milky Way galaxy and M31 would have been intense radio sources similar to Cygnus A or Centaurus A, for only in the youngest galaxies would this be likely to occur.

The Moscow radio astronomer does not attempt to explain in his article the details of what happens when two galaxies really collide. Baade and Spitzer's calculations of the probability of collisions are not challenged; in fact they have been confirmed by G. A. Harrower in the *Astrophysical Journal* for July, 1960. Astronomers throughout the world will be awaiting new observational and theoretical discussions of the exciting ideas raised in Shklovsky's article.

ASTRONOMY IN THAILAND

Chulalongkorn University in Bangkok, Thailand, has recently enlarged its curriculum to include astronomy. Seeking to develop the program, Lt. Gen. Phya Salwidhannidhes, professor of engineering, conferred with astronomers at Harvard Observatory during his visit to the United States in September and October as a participant in the foreign leaders program of the Department of State. He is currently president of the Science Society of Thailand.

PEKING PLANETARIUM

Since its opening in September, 1957, the Zeiss planetarium at Peking, China, has had an average daily attendance of over 2,000 persons. There are five to seven scheduled demonstrations daily, and eight or nine on holidays. In the same building are an astronomical museum and a lecture room seating 500, while a neighboring dome contains a 5-inch refractor. The director is Lu Shu-yuan.

LETTERS

Sir:

Observational astronomy has long been harassed by the results of population growth and increased industrialization. Recently a new and potentially more serious threat has arisen.

Last July 6th, I was reassured by the National Aeronautics and Space Administration that the Echo I satellite would have a "relatively short" life. Now it appears that Echo I may be in orbit for a number of years, and the launching of an Echo II is planned for late this year. The orbiting of as many as 50 of these balloons has been proposed.

Admittedly Echo I not only has scientific but prestige value. However, if more of these bright objects are placed in similar orbits, there may be significant interference with some astronomical research. For example, long-exposure photography, especially with wide-field instruments, could be hampered.

A large number of slow-moving bright objects would markedly change the appearance of the sky and could only cause its disfigurement. Whether any group of people has the right to change the appearance of the night sky is questionable. Much fainter satellites have been successfully tracked, and I can see no useful purpose in having Echo satellites so bright. Blackening them might help.

Other potentially harmful proposals include orbiting or placing high in the atmosphere various particles, gases, and dusts. Unfortunately, once such objects are in orbit nothing can be done about them. Letters sent to members of Congress and to NASA right now should bring more careful thought and action to the problem. Only by immediate attention can we preserve what is so important to us.

ALAN MCCLURE

649 S. Olive St.
Los Angeles 14, Calif.

Sir:

During several passes of the Echo I balloon satellite in August and early September, I found that it twinkled very rapidly. This phenomenon was seen in 7 x 50 binoculars, shaken to cause the satellite's image to trace out a curve. The traces were broken into short segments, showing that the light was varying on the order of 100 times per second. The frequency could be judged by comparison with a distant fluorescent light viewed in the same manner. Such a lamp on 60-cycle current flashes 120 times a second.

On one occasion, I viewed the balloon through a 6-inch telescope, and found that it did not appear to twinkle, even when the telescope was shaken.

These observations can be explained in terms of atmospheric turbulence and the satellite's rapid motion. Scintillation of stars is known to be caused by very slight variations of air density, on a scale of

only a few inches, in the lowest mile or so of the atmosphere. Pockets of denser or lighter air act as positive or negative lenses of very long focus, alternately increasing and decreasing the intensity of the starlight reaching the observer.

The 100-foot aluminized balloon serves as a convex mirror, giving essentially a point image of the sun. Hence the density fluctuations must have been enough smaller than the 6-inch telescope mirror for their effects mostly to cancel out. On the other hand, the fluctuation scale was evidently larger than the small effective diameter of the binocular objectives.

When Echo I, traveling about 4.5 miles per second, is moving perpendicularly to the line of sight, a ray of light from it to the observer's eye cuts through the atmosphere with a velocity of 24 feet per second at a height of one mile. If the scale of the density fluctuations is three inches, the resulting twinkling would have a frequency of nearly 100 per second, in plausible agreement with my estimate of the observed rate in binoculars.

WILLIAM D. ROSS

120 Birch Ave.
Wilmington 5, Del.

Sir:

Although astronomical equipment is rather scarce in our country, there are 700 members in the Yugoslav Astronomical Society. We publish a quarterly magazine *Vasiona (Universe)*, written largely by staff members of the Belgrade Astronomical Observatory.

We believe there may be some amateurs in the United States who would like to read astronomical articles in Serbian. Since our funds are limited, we would appreciate a subscription to *SKY AND TELESCOPE* in exchange for *Vasiona*. We can also send another publication, *Zemlja i Svemir (The Earth and the Universe)*, a quarterly printed by the Zagreb Astronomical Observatory.

DRAGOSLAV EKSINGER

Koce Kapetana 44/III
Belgrade, Yugoslavia

ALICE FARNSWORTH DIES

A well-known teacher of astronomy at Mt. Holyoke College from 1920 to 1957, and director of its Williston Observatory since 1936, Alice H. Farnsworth died on October 1st at Newton, Massachusetts. Born in Williamstown, she was 66. Her doctorate in astronomy was from the University of Chicago, and she had worked at Lick Observatory.

Dr. Farnsworth served as president of the American Association of Variable Star Observers from 1929 to 1931. She was a regular observer of occultations and sunspots, in collaboration with her students. In addition, she published several papers on the distribution of stars in space, and on the absorption of starlight by interstellar dust.

AMERICAN ASTRONOMERS REPORT

Here are highlights of some papers presented at the 106th meeting of the American Astronomical Society at Mexico City, August 22-25, 1960. Complete abstracts will appear in the Astronomical Journal.

Rotation of Stars in Orion

An important difference has been found by D. H. McNamara, Brigham Young University, between the *B*-type stars belonging to the Orion association and similar stars that are scattered over the sky.

Dr. McNamara measured the rotational velocities of 28 stars from the broadening of their spectral lines. All 28 had spectral types between *B0.5* and *B3*, and all but four were main-sequence objects. For any individual star, he could not tell whether low rotational velocity is an intrinsic property, or whether that star's axis of rotation happens to be tilted toward the earth. But he could establish, for the stars as a group, that they are spinning with an average velocity of 126 kilometers per second. This result assumes the stars' axes are oriented at random.

Earlier, a similar study of 67 *B1* to *B3* field stars had been made at Perkins Observatory by A. Slettebak and R. Howard, who found an average rotational rate of 200 kilometers per second. It is not known why the *B* stars belonging to the Orion association rotate so much more slowly.

High-Altitude Balloons

For astronomers interested in placing their instruments well above most of the earth's atmosphere, James R. Smith of Raven Industries, Inc., suggested an increased use of high-altitude plastic balloons, such as employed in Project Stratoscope to observe granulation on the sun. He described these carriers and explained what they can do.

Currently, maximum payloads of 2,500 to 3,500 pounds are being lifted successfully. By using the largest balloons available, these loads can be carried above 100,000 feet, but Mr. Smith pointed out that smaller sizes (perhaps two million cubic feet instead of 10 million) can serve

reliably to carry loads to 80,000 feet and be well within the proven limits of balloon operation.

One serious limitation has been that surface winds are often too strong to permit launchings at will. But at Holloman Air Force Base in southern New Mexico (near Sacramento Peak Observatory), a long-term study indicates that satisfactory balloon launching conditions occur on more than four days in five.

A new result from the Project Stratoscope flight of August 17, 1959, was reported by J. D. R. Bahng and M. Schwarzschild, Princeton University Observatory. They studied an eight-minute sequence of high-definition photographs of the sun's surface, and found the average lifetime (as technically defined) of solar granules to be 8.6 minutes, with a probable error of ± 0.2 minute.

Chemistry of Jupiter's Atmosphere

Recent laboratory experiments to provide new information about the chemistry of planetary atmospheres have been undertaken by Carl Sagan, Yerkes Observatory, and Stanley L. Miller, University of California.

In one type of experiment, they passed electric corona discharges through mixtures of hydrogen, methane, and ammonia in the proportions of 30:3:1, this being roughly the composition of Jupiter's atmosphere above the cloud layer. The gas mixture was allowed to stand until equilibrium was attained, and then samples were extracted for analysis with an infrared spectrograph and a mass spectrograph.

The molecules produced in this way included ethane, ethylene, acetylene, hydrogen cyanide, acetonitrile, and formaldehyde, the last being formed only

when water vapor was present. The experimenters point out that these results disprove the statement frequently encountered in astronomical literature that more complex molecules cannot be produced from methane and ammonia in an excess of hydrogen.

Inclusion of hydrogen cyanide in the list is interesting, since G. P. Kuiper has observed a feature in the infrared spectrum of Jupiter, at a wave length of 1.53 microns, which has been tentatively identified as due to HCN.

The possible presence of acetylene on Jupiter is suggestive, since this gas is known to polymerize explosively under conditions similar to those in the planet's atmosphere, forming a variety of compounds of high molecular weight, some of them strongly colored. This may have bearing on the colorations observed in Jupiter's belts, and it is also possible that such chemical reactions may contribute to Jupiter's radio noise.


Stellar Aluminum Oxide

Among the many absorption bands in the spectra of the cool red stars of type *M* are some due to molecules of aluminum oxide. Under the conditions of temperature and pressure in stellar atmospheres, however, this compound has the composition AlO , rather than the Al_2O_3 familiar on earth. Peculiarities in the behavior of stellar aluminum-oxide bands were described in a paper by Philip C. Keenan, Perkins Observatory.

The bands usually appear as absorption features in long-period variable stars, and as weaker absorptions in ordinary *M*-type giants and small-amplitude red variables. They are stronger in the coolest *M* stars. Using the Perkins 69-inch reflector, Dr. Keenan and his colleagues have made a



In the picture at the left, Rupert Wildt of Yale Observatory talks astronomy with Rudolph Minkowski, Mount Wilson and Palomar Observatories, in the science-auditorium foyer of the Mexican National University. At the right, Arthur N. Cox of Los Alamos Scientific Laboratory converses with Lick Observatory astronomer George H. Herbig, who is holding the abstracts for papers given at the meeting. Photographs by David Tellez.





Among the many small groups that gathered between sessions at Mexico City were the two shown here. At the left, Smithsonian Astrophysical Observatory astronomers (from left to right) Jack W. Slowey, Don A. Lautman, and Leon Campbell, Jr., talk with former associate Karl G. Henize, now at Dearborn Observatory. At the right are Henrietta H. Swope and A. H. Joy, Mount Wilson and Palomar; Helen Pillans, Mills College; and Mrs. Joy. Photos by D. Tellez.

survey of the spectra of 60 long-period variables, finding that aluminum-oxide features differ from star to star. R Hydrae, RT Librae, and SU Virginis have unusually strong AIO absorption bands, but they are weak in RT Cygni.

A hitherto unique anomaly occurred in the spectrum of Mira for several weeks in 1924, when the blue bands of aluminum oxide appeared in emission. Now Dr. Keenan reports that the same event occurred in the spectrum of R Serpentis during March, 1960. This variable passed through a maximum (magnitude 7.9) much fainter than average on March 20th, according to the American Association of Variable Star Observers. The 1924 maximum of Mira was also below normal, but both stars have had other faint maxima at which no AIO emission was observed.

One remarkable exception to the rule that aluminum-oxide bands are prominent only in late *M* stars was provided by the supergiant variable Rho Cassiopeiae, normally an *F* star, when it was unusually faint in 1946.

Zeta Aurigae Masses

One of the most important eclipsing binaries is Zeta Aurigae, which consists of a blue star, spectral type *B7*, and a cool *K4* star of very large dimensions. They revolve around each other in a period of about 32 months, and the *B* star is totally eclipsed by the other for 37 days. Although much is known about this system from its light curve and spectrum, the distance from us is uncertain and the masses are imperfectly established.

Now, at the University of California in Los Angeles, Daniel M. Popper has made a new mass determination by means of spectrograms obtained with the 100-inch Mount Wilson telescope expressly for his study. In all there are 35 spectrum photographs, strongly exposed in the near-


ultraviolet region at 3700 to 4000 angstroms, with a dispersion of 10 angstroms per millimeter. At this large scale, microdensitometer tracings like the one shown here give a very detailed portrayal of the spectrum.

The broad dips in the tracing are the Balmer hydrogen lines of the *B* star, but superimposed on them and badly distorting them are the very numerous sharp lines of metals in the atmosphere of the *K* star. Dr. Popper sought a method of allowing for this distortion, so the true wave lengths of the *B* star's hydrogen lines could be measured and Doppler shifts determined for different positions in its orbit. In this way the motion of the *B* star could be obtained, while the velocity variations of the *K* component were already well known. Most of the spectrograms were obtained near times of extremes of velocity variation, that is, when one star was approaching us and

the other receding, producing the maximum Doppler shift in their lines toward opposite ends of the spectrum.

In order to calibrate the method of reduction, 26 composite photographs of stars of known velocity were used. Each of these spectrograms consists of the superposed spectra of two stars of types similar to those of Zeta Aurigae's components. For each particular hydrogen line measured (*H*₈, *H*₉, *H*₁₀, *H*₁₂, *H*₁₃, and *H*₁₄), a correction to the observed position was obtained, thus making allowance for the distortion by the *K* star's metallic lines.

The provisional results for the masses are 7.9 for the *K* star and 5.5 for the *B* star, in terms of the sun's mass. The *B* component thus seems more nearly normal than was indicated by the value of 10.2 (22.0 for the *K* star) obtained by P. Wellmann in 1951. The estimated mean error of the new values is 15 per cent.



Daniel M. Popper's tracing of the spectrum of Zeta Aurigae was made from a 100-inch-telescope spectrogram. Courtesy University of California.

Lunik II's Landing on the Moon

ON the afternoon of September 13, 1959, as the Soviet rocket Lunik II was nearing the moon, the news agency Tass distributed a prediction that the object would strike the lunar surface at 21:01 Universal time, in the region between Mare Tranquillitatis, Mare Serenitatis, and Mare Vaporum. The forecast led many European observers to watch this area for visible signs of impact, such as a short-lived dust cloud. (At the time in question, the moon was still below the horizon for the United States.)

In the next few days a number of press notices reported optical observations of impact, but the evidence appeared unsatisfactory. There was disagreement on the location of the point of fall, and in most cases the details were lacking. Only the sudden cessation of the space probe's transmissions at the predicted time, and the analyses of radio tracking observations in the United States, Great Britain, Japan, and the U. S. S. R., demonstrated that Lunik had in fact reached the moon.

This situation has now been placed in an entirely new light by L. Detre, an internationally known astronomer who is director of the Budapest-Szabadsaghegy Observatory in Hungary. In No. 45 of the *Contributions* of that institution, he gives a full account of certain decisive observations in Hungary and Sweden.

At the Budapest Observatory, staff member M. Lovas was viewing the moon at the crucial time with the 7-inch refractor. The seeing was good, and he could use a magnification of 500. With him in the

dome were two other staff astronomers, Julia Balazs and B. Balazs. Very shortly after 21:02:30 UT, Lovas said, "There is a dark point I did not see before." A few seconds later, he called out that the spot was expanding. Both his colleagues took a turn at the eyepiece, and were able to verify the expansion. As the spot grew, it faded, and was about 40 kilometers in diameter when it was last definitely seen, at 21:07.

Lovas estimated the position of the marking as close to a small crater plotted in the International Astronomical Union's lunar atlas at selenographic co-ordinates $\xi = +.024$, $\eta = +.434$. This is in Mare Imbrium, south of the crater Autolycus and not far from the Apennines, but somewhat outside of the predicted impact area of Lunik II.

The same dark marking was independently seen in a 10-inch reflector by M. Ill at the public observatory and satellite-tracking station of Baja, in southern Hungary. He devoted particular attention to fixing the location of the spot, whose co-ordinates he estimated as $\xi = +.03$, $\eta = +.45$, a position 25 kilometers northwest of Lovas'. Mr. Ill could keep the marking in view until 21:08. Because his observations were reported in Hungarian provincial newspapers the next day, before he could know of the details of the Budapest sighting, this appears to be an independent confirmation.

Also on the night of September 13th, a good series of lunar photographs was taken at Uppsala, Sweden, by E. Tengström, with the Markowitz moon camera of the university's geodetic institute. Professor Detre wrote to Uppsala to ask if these photographs showed any unusual feature at the co-ordinates $+024, +434$.

Tengström replied that he found nothing definite in that position, but 25 kilometers northwest there seemed to be a very small, sharp, dark spot on his photograph of 21:03.6 UT, and a trace of it on the next exposure, taken 1.9 minutes later.

Professor Detre summarizes his conclusions in the following words:

"1. Shortly after the radio signals from Lunik II ended, an expanding spot on the moon could be observed in the position $+03, +45$, which was at first black and pointlike, a few minutes later gray

In M. Ill's sketch map, right, his dark spot appeared at the place marked X. Most of the same area is shown on the Mount Wilson 100-inch-telescope photograph at the left. Both drawings on this page are from the "Contributions" of the Budapest-Szabadsaghegy Observatory in Hungary.



The impact cloud of Lunik II, drawn by Budapest observers at one-minute intervals. One inch on this picture corresponds to about 50 miles.

and diffuse, with a diameter of about 40 kilometers. The reality of the phenomenon is beyond doubt. It was most probably a cloud of dust.


"2. There is a faint photographic trace, still awaiting confirmation, of this phenomenon on the plates taken by Tengström in Uppsala, which should furnish a precise positional determination.

"3. The phenomenon observed by the Hungarian astronomers is the only one compatible with the impact point subsequently deduced by Soviet radio tracking of the instrument section of Lunik II."

EAST-WEST EXCHANGE

An American solar astronomer is spending six months at the Crimean Astrophysical Observatory while a Soviet specialist in stellar astrophysics is working at the University of California, in an exchange program sponsored by the U. S. Department of State and supported by the Guggenheim Foundation.

The American is Harold Zirin, of the High Altitude Observatory in Colorado, who will study solar magnetic fields. His Soviet counterpart is Crimean astronomer A. A. Boyarchuk.





The ALPO exhibit at the Astronomical League meeting. Photo by the author.

Convention at Haverford

WILLIAM K. HARTMANN,

Allegheny Valley Amateur Astronomers Association

THERE were 278 registrants from 25 states, the District of Columbia, Canada, Puerto Rico, and Cuba at the 1960 convention of the Astronomical League, held over Labor Day weekend at Haverford College in Pennsylvania. Philadelphia's Rittenhouse Astronomical Society was the host, with Edwin F. Bailey as general chairman.

The meeting opened with a business session on Saturday, September 3rd. Mrs. Wilma Cherup, executive secretary, stated that the league had gained 16 new member societies, nine of them in the Northeast Region, during the past year. A description was given of league services now available to members. These include a manual for holding regional conventions, a 10-per-cent discount on astronomical books, and an information bureau on instruments. In the planning stage are filmstrips with lecture tapes, as well as manuals on preparing papers and organizing club projects.

Five talks were given at the first general session that afternoon, including an account by Miss E. T. Vadala of Philadelphia of her observing experiences in high-flying balloons, and a lively presentation by Dr. Earle G. Linsley of Honolulu, Hawaii, on Polynesian astronomical lore. This was followed by the motion picture *Universe*, a production of the National Film Board of Canada and winner of the animation prize at the

Cannes, France, film festival. It is an introduction to astronomy, and shows with beautiful perspective numerous planetary and stellar animated sequences.

After the movie, a junior session was held. Among the speakers who were later awarded prizes for their presentations were Robert Kenison, Madison, Wisconsin; Charles Downton, Jr., and Robert Dragon, both of Springfield, Massachusetts. A proposed classification system for deep-

sky objects by George Doschek and James Mullaney of Pittsburgh aroused much comment. Tim Wyngaard, Madison, concluded the program with an illustrated account of his trip to the Canary Islands with the United States solar eclipse expedition of October, 1959.

That evening three observatories were visited. On the Haverford campus, excellent views of Jupiter and Saturn were had through the instruments of the Strawbridge Memorial Observatory; passage of the Echo I satellite enlivened the star party. Nearby Villanova Observatory was open to delegates. John J. Ruiz, of Danemora, New York, was one of those who was invited to the Flower and Cook Observatory by its director F. B. Wood. Mr. Ruiz's description follows.

"The observatory is used exclusively for research work and is not ordinarily open to the public. This was a special demonstration of part of the electronic equipment for those amateurs who either had themselves built photometers and made photoelectric observations or who were seriously considering doing so.

"Mr. E. G. Reuning showed us the 28½-inch reflector, which has an infrared photoelectric photometer that utilizes lead-sulfide cells and works at wave lengths greater than 9000 angstroms. Appropriate filters isolate a region lying around 1.6 microns for brighter objects. The instrument is currently employed for measures of certain intrinsic and eclipsing variables, especially Algol.

"On the 15-inch horizontal refractor, Dr. William Blitzstein demonstrated a pulse-counting photometer that is being used by graduate students for multicolor observations of eclipsing systems. We were also shown a conventional direct-current photometer that is employed by L. Binnendijk and his students on the reflector, and the instruments for scintillation studies by W. M. Protheroe. We are in-



This convention group picture by Ray Satori is continued on the next page.



Seated at the head table for the honors dinner on Monday evening were, from left to right, Dr. Louis C. Green, Mrs. Chandler Holton, Mr. Holton, Edwin F. Bailey, Rev. Edward Jenkins, Mrs. Wilma Cherup, Manfred Kamin-tius, and Walter H. Haas. Photograph by Frank Delaney.


debted to Dr. Wood and Mr. Reuning for devoting so much of their time and patience in answering our many questions."

On Sunday, an all-day bus trip took the delegates from Pennsylvania to New Jersey, and into Delaware. The first stop was at the Franklin Institute for a showing of the Zeiss projector of the Fels Planetarium and inspection of the rooftop observatory. On the way to Edmund Scientific Company in Barrington, New Jersey, where a steak dinner was served, the buses passed such historical sites as Independence Hall and Betsy Ross house. Then the tour went to Spitz Laboratories in Yorklyn, where the new Spitz model A-3-P planetarium projector was unveiled. After a buffet supper given by the company, the buses left for Swarthmore, Pennsylvania, where the party inspected Sproul Observatory's 24-inch refractor, used for trigonometric parallax determinations.

The Association of Lunar and Planetary Observers held its meeting on Mon-

day morning, with Walter H. Haas, Edinburg, Texas, as chairman. The Jupiter recorder, Philip R. Glaser, of Menomonee Falls, Wisconsin, reported that a disturbance in that planet's south equatorial band seems to be imminent. According to Phillip W. Budine, Binghamton, New York, activity on Saturn was unusually great this year, a number of prominent long-lasting white spots having been observed in fairly high latitudes. At the league's instrument session that afternoon, seven papers were presented, ranging from a null test for paraboloids by Charles Spoelhof, Rochester, New York, to the Lehigh Valley Amateur Astronomical Society's planetarium project (*SKY AND TELESCOPE*, June, 1959, page 430).

At the final business meeting, the following officers were elected for the coming year: Norman C. Dalke, Seattle, Washington, president; Ralph K. Dakin, Pittsford, New York, vice-president; Dr. Herman C. Schested, Ft. Worth, Texas, secretary; and



G. E. Audy, Wilmington, Delaware, built this 8-inch reflector for an amateur in the Canal Zone. Photograph by the author.

Leonard G. Pardue, Miami Springs, Florida, treasurer. The Middle East Region named the following officers: David D. Meisel, Fairmont, West Virginia, chairman; Anthony Doschek, Pittsburgh, vice-chairman; Emil Volchek, Wilmington, Delaware, secretary; and Edward Naylor, Harrisburg, treasurer. Detroit, Michigan, will be the site for the 1961 convention, which will be held over the July 4th weekend.

The closing event was the honors dinner on Monday night. Daniel Kleinman, Louisville, Kentucky, was named the outstanding junior of the year, and an ALPO award was presented to Mr. Meisel. The banquet speaker was Dr. Louis C. Green, of Haverford College, who talked on extending observational astronomy beyond the earth's atmosphere.



The 14th general convention of the Astronomical League, at Haverford College, Pennsylvania, on September 3-5, 1960.





Lick 120-inch Photographs—V

FACING PAGE: This great gaseous nebula in Serpens was photographed in a 40-minute red-light exposure with the 120-inch Lick Observatory reflector by N. U. Mayall on July 20, 1960. The emulsion was Eastman 103a-F and the filter OG1. North is at the left, west above, and the scale is 4.1 seconds of arc per millimeter. The open star cluster to the left and above center is Messier 16, whose distance from us is nearly 4,000 light-years. Noteworthy are the many, very small, black globules scattered over the nebula, some fine examples being situated on the southeastern outskirts of M16. These compact dust clouds may be condensing to form stars; their reality can be verified by a comparison with the photograph taken by the 200-inch telescope and reproduced on pages 246-7 of the August, 1951, issue. There Otto Struve discusses the problem of "Clouds with Silver Linings," since this nebulosity is also remarkable for the bright-rimmed lanes of obscuring material that extend into it from many sides.

ABOVE: Messier 27, the famous Dumbbell nebula in the constellation of Vulpecula, is one of the largest and brightest planetary nebulae in the sky. An enormous expanding cloud of highly rarified gas, it is excited to shine by a faint but very hot central star. Its distance is estimated at 1,000 light-years. Dr. Mayall obtained this photograph with the 120-inch telescope on June 23, 1960, a 15-minute exposure on a 103a-O plate. North is toward the bottom in this reproduction, which has a scale of 3.1 seconds of arc per millimeter. The nebula is a difficult object for pictorial photography because of the great range in luminosity between the dim outlying parts and the bright central regions, whose flocculent structure is well shown here. Compare this view with the drawing ("Sky and Telescope," October, 1959, page 665) made in 1860 by William Lassell from visual observations with a 48-inch reflector. These photographs and those on the next page are courtesy Lick Observatory.



The large-scale picture above is of h Persei (NGC 869), the western member of the Double Cluster in Perseus, as shown in a five-minute exposure with the 120-inch reflector. It was taken December 19, 1959, by N. U. Mayall, using Eastman 103a-E emulsion and an RG1 filter for red light. The Double Cluster is at a distance from us of 7,300 light-years. In an f/5 mirror, the field of sharp definition is small, and all stars not close to the optical axis show fan-shaped images, distorted by coma. Hence, the 120-inch telescope has a correcting lens, placed a short distance in front of the photographic plate, and this negative was exposed through it. The effectiveness of the corrector is indicated by the neatness and roundness of the star images in the boxed area (enlarged fourfold at the left), which is about 15 minutes of arc southeast of the cluster center. The fine quality of the big Lick instrument is further shown by its light grasp, for in a 30-minute exposure on fast blue-sensitive plates, stars close to photographic magnitude 23 are recorded. Pictures made under the most favorable conditions allow images of galaxies to be distinguished from stars to as faint as about magnitude 22½. This impressive performance is surpassed only by the 200-inch telescope on Palomar Mountain.

NEWS NOTES

PLANETARY ORIGIN AND LITHIUM ABUNDANCES

The cosmic distribution of the third lightest element, lithium, is far from uniform. In the sun it is relatively less abundant than on earth. Recently, W. K. Bonsack and J. L. Greenstein have found that the T Tauri stars are abnormally rich in lithium, the surfaces of these stars having at least 10 times as much of this element per gram of material as do the nebulae surrounding them (see August issue, page 81). These facts have now led Thomas Gold, Cornell University, to discuss their bearing on the problem of the origin of the solar system. In the *Astrophysical Journal* for July, he writes:

"It has often been suggested that the planetary material may once have been part of the sun in the form of an outer envelope that was later thrown off and that then, perhaps through magnetic interaction, it obtained almost all the angular momentum initially resident in the sun. Such a thin, far-flung disk of gas may then have condensed into the planets. The lithium evidence would now strongly suggest that this indeed has been the course of events. The sun in its early stage would, as a consequence of shedding its magnetic energy, have formed an atmosphere rich in lithium just like the T Tauri stars presently observed and then, in forming a planetary disk out of this material, would have assured the high lithium abundance in the eventual planets."

Dr. Gold points out that if stars generally go through a T Tauri stage and then shed their atmospheres rich in lithium (and possibly deuterium), we might identify by spectroscopic means a small nebula around some such star. This case would then be similar to the solar system before the formation of planets.

COPERNICUS AND CALVIN

After the death of Copernicus in 1543, his doctrine of planetary motions around the sun gained only gradual acceptance, and for a century it was the source of religious as well as astronomical controversy.

It is well known that Martin Luther was a severe critic of Copernicus, and the same is often said of John Calvin (1509-64), the French Protestant reformer at Geneva, Switzerland.

The question of Calvin's actual views on astronomy has recently been carefully surveyed by Edward Rosen, City College of New York, in the *Journal of the History of Ideas* (July-September, 1960). Dr. Rosen shows that the astronomical allusions in Calvin's writings are pre-Copernican. Far from being an opponent of Copernicus, Calvin probably

was unaware of the Polish astronomer's work.

The legend that the Genevan reformer was a critic of Copernicus originated, according to Dr. Rosen, in a mistake by the English ecclesiastical writer F. W. Farrar (1831-1903).

INVITATIONS TO BERKELEY IAU MEETING IN 1961

Attendance at the August, 1961, meeting of the International Astronomical Union will be restricted to its members and invited guests. Invitations to non-members are to be limited to the following categories:

1. Astronomers who are being proposed for IAU membership.
2. New astronomers who are potential members.
3. Outstanding scientists in other fields who have wide astronomical interests.

The U. S. national committee of the IAU has asked astronomy departments to nominate the top 10 per cent of their graduate students for invitations under category 2, also recent recipients of the doctoral degree if they are deemed potential union members. Persons in the third category should be recommended by two members.

Correspondence concerning invitations should be sent promptly to Dr. Frank K. Edmondson, Goethe Link Observatory, Indiana University, Bloomington, Ind.

CHRISTMAS MEETINGS

On December 28-31, the American Astronomical Society will hold its 107th meeting, together with Section D of the American Association for the Advancement of Science. The sessions for papers are open to all persons registering for the 127th AAAS meeting, which takes place in New York City December 26-31. The astronomical headquarters are to be at the Hotel Roosevelt.

The program features a joint symposium to be held with Section B (physics), "Plasma: the Fourth State of Matter," on Wednesday morning, December 28th. The next day, when AAS paper sessions begin, Dr. Ira S. Bowen, director of Mount Wilson and Palomar Observatories, is scheduled to give the address of the retiring chairman of Section D. That evening, the Helen B. Warner lecturer will be Halton C. Arp, also of Mount Wilson and Palomar.

NEW YORK PLANETARIUM'S 25TH ANNIVERSARY

On October 2nd, the American Museum-Hayden Planetarium marked completing its first 25 years of operation. Members of the original 1935 staff were invited to a banquet and showing of the current demonstration, "20,000 Leagues Above the Seas," the Sunday evening public audience

IN THE CURRENT JOURNALS

INTERFERENCE FILTERS, by D. M. Hunten, *Journal of the Royal Astronomical Society of Canada*, August, 1960. "The first interference filters were made about 20 years ago by Geffcken of the Zeiss firm, but they did not see much application until some time after the war. Their advantages over conventional absorption filters are that they can isolate a very narrow region of the spectrum and that this region can be at any desired wave-length."

SURVIVAL POSITION LOCATION USING STAR SIGHTING, by E. H. Sharkey, *Navigation*, Vol. 6, No. 8, 1960. "A method using star sighting which can be used by persons having no knowledge of celestial navigation has been developed and tested. Neither sextant nor almanac is required. The necessary equipment fits into a shirt pocket and provides an accuracy of 10 to 20 miles under realistic field conditions."

ASTRONOMERS IN TURMOIL, by Otto Struve, *Physics Today*, September, 1960. "The explosion which we are witnessing today is mainly due to the sudden recognition by our people of the importance of what the popular writers call the conquest of space. . . . But I believe we are in fact living in a period of vigorous, though much more gradual, increase of ideas, which has little or no relation to the sputniks and which began in the early years of this century. . . ."

PHENOMENA OF THE SOLAR ATMOSPHERE, by R. Grant Athay, *Science*, September 16, 1960. "Among the many unanswered questions regarding phenomena of the solar atmosphere, those that stand out are: (i) What is the mechanism of energy supply? (ii) What is the role of magnetic fields? (iii) What is a flare? (iv) What are the instabilities that give rise to spicules and solar activity? (v) What are the nature and variability of solar corpuscular and short-wavelength radiation incident upon the earth as a result of solar activity?"

partaking of coffee and dessert after the performance.

Attendance at the planetarium in its first year was more than 600,000, a figure not exceeded until the opening of the satellite era three years ago. Early this year extensive improvements in the physical plant were made, including the installation of theater-type seats and a new Zeiss projector (*SKY AND TELESCOPE*, March, 1960, page 271). Further plans call for addition of a new wing to the building with exhibit space, classrooms, an enlarged library, and a research laboratory.

Amateur Astronomers

A HIGH SCHOOL OBSERVATORY IN CALIFORNIA

A DREAM of many years has come true for the astronomy club of the Mount Whitney High School here in Visalia, California. In the past, as sponsor of the group, I made my 6-inch home-built reflector available for celestial observations, to augment constellation studies and discussions of theory. Now we have an observatory that not only fills our needs but affords an opportunity to introduce astronomy to elementary and junior high school classes, and to the adults of our community as well.

The observatory is a result of special efforts by our senior class last year, some faculty members, and several townspeople. Early in the 1959 school year, under the National Defense Education Act, our federal government agreed to pay half the cost of a 10-inch reflecting telescope. Our school district offered the rest, if we could finance and construct the building. The senior class accepted the challenge.

There were about 20 boys and five teachers on hand when ground was broken the day after Thanksgiving. The crew varied in size from one to 15 students, who worked evenings and weekends to finish the project just before school closed last June. Nearly all technical work and supervision were done by six teachers from the mathematics, chemistry, physics, auto shop, metal shop, and mechanical drawing departments.

A lecture room 20 by 20 feet is attached to the observatory, and both are made from concrete blocks. The floors are of asphalt tile, with acoustical tile on the ceiling. The dome skeleton is of channel iron, and it is covered with plaster lath. Fiberglass was blown onto the outside, and acoustical plaster on the inside. The dome is carried by wheels with four-inch bearings and rides on a circular channel-iron ring.

To earn funds for the building materials, nearly \$2,000, the students conducted a fashion show, a Christmas tree

sale, and cake sales. But the biggest fund-raising effort was a basketball game between the school coaching staff and members of the San Francisco 49'ers professional football team.

The observatory is located within 50 feet of the science building. Although we must contend with city lights and several trees, we do have a considerable area of dark sky in which to observe.

ARTHUR SHAHZADE
Mount Whitney High School
Visalia, Calif.

JONES BEACH, NEW YORK

"Understanding the Solar System" is the general theme of this season's meetings of the Star Gazers of Jones Beach, New York. Each of these public sessions starts with a brief lecture at Fieldhouse No. 1, and if the skies are clear, is followed by outdoor viewing through 10 manned telescopes. The meetings are on November 6th and 20th, December 4th and 18th, at 7 p.m.

Percy M. Proctor, of Babylon High School, is the lecturer. The get-togethers are cosponsored by the Long Island Park Commission and Abraham and Straus department store.

CLEVELAND OPEN NIGHTS

Warner and Swasey Observatory is continuing its series of public nights during the coming year. On November 3rd and 4th, the topic is "The Sun"; December 1st and 2nd, "Comets, Meteors, and Dust"; February 2nd and 3rd, "Stars and Star Clusters"; March 2nd and 3rd, "Variable Stars"; and April 6th and 7th, "From Atoms to Galaxies."

Lectures begin at 8 p.m., with Thursday nights reserved primarily for adults. Reservations may be made with the observatory, Taylor and Brunswick Rds., E. Cleveland 12, Ohio; phone, Glenville 1-5625.

*** AMATEUR BRIEFS ***

More than 400,000 persons have attended the annual star parties cosponsored by the Cleveland Press and the Cleveland Astronomical Society since the first one was held in 1929. Dr. David Dietz, science editor for the Scripps-Howard newspapers, promotes these events. Telescopes are provided by society members.


From the bulletin of the Astronomical Society of South Australia: "When is it safe for an ATM to finish his mirror spherical? . . . As a back-garden amateur writing for back-garden amateurs, we would say — stick to spherical mirrors, and make the tube as long as may be convenient. This means that observations made at the zenith should not call for the assistance of a box or a stepladder in order to reach the eyepiece. These wooden aids are uncomfortable to use, and they are a menace in other ways. Keep both feet on the ground with a 6-inch f/10. For short men, f/9. For tall men, f/11. If visitors are to be considered, f/8.2 as per Texereau. If a richest field telescope is contemplated — but that, as Kipling would say, is another story."

During last summer, J. Russell Smith conducted astronomy classes for 70 students of East Texas State College in Commerce, Texas. His Skyview Observatory in Eagle Pass is always open to amateurs passing through the area.

Each member of the Feather River Astronomy Club in Oroville, California, is required to handle the program at one meeting during the year. The society is currently polishing the mirror for a 10½-inch reflector.

Jim Heinemann, of the Denver, Colorado, Astronomical Society, solved the problem of transporting his 10-inch reflector to a club star party by removing the entire back seat of his car, thereby allowing the telescope tube to extend into the trunk.

Clubs desiring to enliven their meetings may be interested in two sound motion pictures now available for rental



The exterior (left) and interior (above) of the Mount Whitney High School observatory in California. The twin staircase aids group viewing.

from Contemporary Films, 267 W. 25th St., New York 1, N. Y. One, entitled Universe, is a description of a night-long observing session at the David Dunlap Observatory in Canada; the other, The Inquisitive Giant, is a study of the huge radio telescope at Jodrell Bank in England. The rental fees are \$7.00 and \$7.50, respectively; each film runs for 28 minutes.

Amateurs make newspaper headlines! Arnold Norman, Jr., of Little Rock, Arkansas, had one of his moon photographs on the front page of the *Arkansas Gazette*, and the next day there was an account of his equipment in the *Arkansas Democrat*. Sherman W. Schultz's home-made observatory was the cover story for the Sunday pictorial magazine section of the St. Paul, Minnesota, *Pioneer Press*. Two large pictures of a Kenosha Amateur Astronomers' star party were first-page items of the Kenosha, Wisconsin, *Evening News*.

Travelers passing through Amarillo, Texas, are invited to take in the Friday night observing sessions held by the Panhandle Astronomical Society at Old Soncy School, on U. S. highway 66 west of the city limits.

January 1st is the date by which the Buffalo Astronomical Association hopes to have its observatory in operation. The equipment will include a 7-inch refractor, a 12-inch Newtonian reflector, a remote-controlled all-sky camera, and an adjoining building for offices. The observatory is at the Newstead site of the Cornell Aeronautical Laboratories' ionospheric research center, about a mile east of Clarence, New York.

H. M. C.

NEW ENGLAND AMATEURS MEET IN RHODE ISLAND

About 80 amateurs from all parts of New England were present for the annual astro-assembly sponsored by Skyscrapers, Inc., at their Seagrave Observatory in North Scituate, Rhode Island, on October 1st. Refreshments were served by the hosts in the afternoon, while several astronomical competitions were being judged.

The Amateur Telescope Makers of Boston not only won the prize for largest attendance, but swept the telescope contest with entries by William Sheehan, Ann Sudarich, and Walter Redding. The best astronomical photograph was submitted by the Aldrich Astronomical Society of Worcester, Massachusetts, and an award was given to Arthur Crowe of the host club for his test rig for Maksutov optics.

A steak dinner in the evening preceded an informal series of astronomical ghost stories by Raymond Watts, Jr. Clear skies prevailed for the star party that followed, guests using the Skyscrapers' 8-inch Alvan Clark refractor and other telescopes.



Senior and junior members of the Racine Astronomical Society constructed this colorful float to publicize their club during the July 4th parade held in the Wisconsin city last summer. Photograph by George A. Dechant.

RACINE, WISCONSIN

IN order to promote civic interest in its activities, the Racine Astronomical Society entered the float pictured here in the city's annual Fourth of July parade. Both senior and junior members volunteered time and money to build the 7-by-14-foot display. Total cost of the project was under 50 dollars.

The float was planned on a red, white, and blue motif. We fashioned the observatory dome frame from iron wire hoops with struts running to the top, and then covered it with chicken wire. Artificial flowers were made from 7,400 paper napkins, and each was individually stuffed into the frame! Four senior and two junior members rode on the platform, along with three telescopes. The rear wheel hubs of the farm tractor donated to pull the exhibit were covered by a cardboard representation of the earth and an orbiting satellite ball. Our planet did not rotate on its proper axis, but this was intentional, as we wished to have it readily recognized.

The float carried a sign publicizing a star party we held later in the month. Nearly 400 persons attended, despite imperfect observing conditions, and three signed up as new members.

GEORGE A. DECHANT
2225 Shoop St.
Racine, Wis.

CHICAGO ASTRONOMY CLASSES

A series of courses in astronomy, telescope making, and navigation is currently being given at the Adler Planetarium. For information, write the planetarium at 900 E. Acheson Bond Dr., Chicago 5, Ill., or phone Wabash 2-1428.

THIS MONTH'S PROGRAMS

Baltimore, Md.: Baltimore Astronomical Society, 8 p.m., Enoch Pratt Free Library main building. November 21, William P. Gonce, Sr., Martin Co., "Living in Space."

Cleveland, Ohio: Cleveland Astronomical Society, 8 p.m., Warner and Swasey Observatory. November 11, Dr. S. W. McCuskey, Warner and Swasey Observatory, "Satellites and Their Orbits."

Dallas, Tex.: Texas Astronomical Society, 8 p.m., Health and Science Museum. November 28, Ted F. Gangl, "Living with Radiation."

New Orleans, La.: Pontchartrain Astronomy Society, 8 p.m., Tulane University Observatory. November 4, Dr. J. Frazer Thomson, Tulane University Observatory, "The Stars."

New York, N. Y.: Amateur Astronomers Association, 8 p.m., American Museum of Natural History. November 2, Dr. Henry Miller, S. J., Fordham University, "Journey to the Center of the Earth — 1960."

New York, N. Y.: Junior Astronomy Club, 8 p.m., Waverly building, New York University. November 18, Dr. Otto Struve, National Radio Astronomy Observatory, "Recent Advances in Astrophysics."

FLORIDA GROUP REORGANIZES

Members of the Choctaw Astronomical Society have reorganized their club, naming it the Northwest Florida Astronomical Association and moving their headquarters to Ft. Walton Beach. The group has 48 seniors and seven juniors. Interested persons are invited to contact J. M. Sherlin, Flamingo Apt. 2, 156 Westview Ave., Valparaiso, Fla.

OBSERVING THE SATELLITES

DISCOVERER XV

TENTH VEHICLE to achieve orbital velocity in the U. S. Air Force's program to improve satellite technology, Discoverer XV was successfully fired from Vandenberg Air Force Base in California, at 22:13:39 Universal time on September 13th. A later launching time than that for others in the Discoverer series gave fewer daylight hours for the recovery attempt.

The 300-pound capsule was separated on schedule on the 17th revolution, but either orientation of the Agena or thrust from the capsule's retro-rocket went awry, for the re-entry package sailed over the intended recovery zone to a point near Christmas Island in the mid-Pacific Ocean. Less than 3½ hours of daylight remained when radar tracking at Kaena Point, Hawaii, placed the capsule about 1,000 miles from waiting ships and planes. By the time a searching seaplane spotted the floating payload, it was necessary to postpone recovery attempts to the following morning. When operations were resumed, the ocean was too rough to permit a seaplane to land, and the capsule was lost in the turbulent waters within a few hours.

The Agena vehicle remains in orbit as 1960μ. Its initial anomalistic period was 94.19 minutes, with perigee and apogee heights of 126 and 474 miles. The orbital plane is inclined 80.90 degrees to the earth's equator. These values have been derived from information supplied by the National Space Surveillance Control Center (Space Track).

The payload of Discoverer XV was designed to monitor performance during the recovery sequence and to make un-


specified measurements of the space environment. Later Discoverers are expected to carry biological specimens.

MORE ABOUT SPUTNIK V


INFORMATION has been received from Soviet scientists about the recovery of the dog-carrying capsule from Sputnik V (October issue, page 201). When the capsule was ejected, 1960λ1 was traveling at only 485 miles per hour, far less than orbital velocity. Thus, the spaceship must have been slowed down prior to the ejection so that the capsule would not suffer the effects of re-entry heating. The final descent to earth was by parachute.

The accompanying photographs show some of the details of the life-support system. The capsule was fully pressurized, with an oxygen content held between 20 and 25 per cent and a carbon-dioxide level below one per cent, according to Soviet accounts. The miniature biological museum was designed to answer several fundamental questions relating to space-environment effects on physiology and subsequent reproductive ability of the specimens.

Chlorella, a seaweed, was carried aloft for a special reason. Sputnik V was described as the first satellite experiment to investigate the use of photosynthesis for removing carbon dioxide and generating oxygen and possibly food during space voyages of long duration.



In this cutaway view of the payload of Sputnik V, the following parts are identified: 1, air-supply tank; 2, ejection firing device; 3, radio beacon; 4, storage battery for heating test tubes containing microbes; 5, storage battery; 6, unspecified instrumentation; 7, container; 8, movement transducer; 9, animals' pressurized capsule; 10, microphone; 11, radio aerial; 12, intake and exhaust valves; 13, television camera; 14, mirror; 15, ventilator; 16, automatic food dispenser. The photographs on this page are courtesy U.S.S.R. Embassy in Washington, D. C.



The recovery capsule of Sputnik V as exhibited by Soviet scientists. The pressurized animal container, here seen nestled in the center of the capsule, is shown from the opposite side in the cutaway view above.

THIRD YEAR OF THE SPACE AGE

THE TABLE opposite provides a listing of all known satellite and space-probe launchings during the past year, and is a continuation of a similar rundown that was presented for the first two years of the space age on page 13 of the November, 1959, issue of SKY AND TELESCOPE.

To bring the earlier list up to date, only a few changes are needed: The solar-powered radio in Sputnik III functioned until the satellite ended its orbital career on April 6, 1960; Explorer IV came down on October 23, 1959; radio transmission from Explorer VI ceased on October 6, 1959; that from Vanguard III on December 11, 1959; and Discoverer VI descended on October 20, 1959. The current table begins with the present status of Lunik III.

The new tabulation, together with the earlier one amended above, gives a complete report as of October 4, 1960, for the known launch attempts.

SATELLITE AND PROBE LAUNCHINGS DURING THE THIRD YEAR OF THE SPACE AGE

	Launched	Site	Name and Designation	Status	Notes
1959	Oct. 4	S	Lunik III — 1959 θ	Down April, 1960	Transmission of moon pictures ended before Oct. 18, 1959. Rocket probably in orbit.
	Oct. 13	C	Explorer VII — 1959 ϵ 1	In orbit	Radio operating. Rocket, 1959 ϵ 2, also in orbit.
	Nov. 7	V	Discoverer VII — 1959 κ	Down Nov. 26, 1959	Fault in power supply prevented capsule recovery.
	Nov. 20	V	Discoverer VIII — 1959 λ	Down Mar. 8, 1960	Capsule separated but overshot recovery area.
	Nov. 26	C	Atlas-Able IV (moon orbiter)	No orbit	Shroud protecting payload came off during Atlas thrust.
1960	Feb. 4	V	Discoverer IX	No orbit	Fuel-loading equipment failure damaged vehicle.
	Feb. 19	V	Discoverer X	No orbit	Thor veered off course. Destroyed by safety officer.
	Feb. 26	C	Midas I	No orbit	Second-stage Agena failed to separate from Atlas.
	Mar. 11	C	Pioneer V — 1960 α	Solar orbit	311.64-day period. Jodrell Bank held radio contact to 22,462,115 miles. Rocket also in orbit.
	Mar. 23	C	Explorer	No orbit	Telemetry lost shortly after Jupiter burnout; one of the Sergeant-powered stages failed.
	Apr. 1	C	Tiros I — 1960 β 2	In orbit	Cloud pictures ended June 17, 1960. Tracking transmitter operating. Rocket, 1960 β 1, in orbit.
	Apr. 13	C	Transit I-B — 1960 γ 2	In orbit	Radios silent June 19 and July 12, 1960. Able-Star rocket, 1960 γ 1, and separated object, 1960 γ 3, also in orbit.
	Apr. 15	V	Discoverer XI — 1960 δ	Down Apr. 26, 1960	Capsule separated but was not observed.
	May 13	C	Echo	No orbit	Failure of orientation control during second-stage coasting.
	May 15*	S	Sputnik IV — 1960 ϵ 1	In orbit	Radio silent July 2, 1960. Rocket, 1960 ϵ 2, down July 17, 1960. Separated parts of spaceship, 1960 ϵ 3 through 9, also in orbit; except 7, down Sept. 24, 1960; possibly others down.
	May 24	C	Midas II — 1960 ζ 1	In orbit	Telemetry of infrared data ended May 26, 1960. Fairing, 1960 ζ 2, in orbit.
	June 22	C	Transit II-A — 1960 η 1	In orbit	Radios operating. Separate payload, 1960 η 2, also functioning. Able-Star rocket, 1960 η 3, in orbit.
	June 29	V	Discoverer XII	No orbit	Failure of orientation control during second-stage firing.
	Aug. 10	V	Discoverer XIII — 1960 θ	In orbit	Capsule recovered Aug. 12, 1960.
	Aug. 12	C	Echo I — 1960 ι 1	In orbit	Radios operating. Rocket, 1960 ι 2; two hemispheres, 1960 ι 3 and 5; and piece of aluminized mylar, 1960 ι 4, also in orbit.
	Aug. 18	V	Discoverer XIV — 1960 κ	Down Sept. 16, 1960	Capsule recovered Aug. 19, 1960.
	Aug. 18	C	Courier I-A	No orbit	First-stage Thor exploded.
	Aug. 19	S	Sputnik V — 1960 λ 1	Recovered Aug. 20, 1960	Capsule and spaceship recovered. Rocket, 1960 λ 2, down Sept. 23, 1960.
	Sept. 13	V	Discoverer XV — 1960 μ	In orbit	Capsule sighted but lost at sea.
	Sept. 25	C	Pioneer VI (moon orbiter)	No orbit	Second-stage malfunction.
	Oct. 4	C	Courier I-B — 1960 ν 1	In orbit	Radios operating. Able-Star rocket, 1960 ν 2, in orbit.

*May have been May 14th UT.

This table is compiled largely from information supplied by the National Aeronautics and Space Administration and by the National Space Surveillance Control Center. Under *Site*, C and V designate American launchings from Cape Canaveral, Florida, and from Vandenberg Air Force Base, California, respectively; S, Soviet launchings from unspecified sites. The U. S. S. R. has not released information on unsuccessful attempts. Dates are in Universal time.

Of these there have been 64 in all, 26 of them in the year under review. The total includes eight announced Soviet firings, three of them in the interval covered by this table.

The United States has fired 56 times toward space during the triennium, achieving 29 successes. These numbers are 23 and 14, respectively, for the year ending October 4, 1960. Thus, just over half of the American tries have been successful, a good performance ratio when the enormous complexity of the missiles is considered.

A typical launching vehicle must perform over a hundred separate automatic functions within close tolerances in order to place an artificial satellite into a chosen orbit. Malfunctioning of such a minor component as a relay or a rate gyro can render the mission abortive.

Of the Soviet launchings, three may be counted as deep-space shots: Lunik I or Mechta, which sailed past the moon to become an artificial planet moving around

the sun; Lunik II, which struck the moon (see page 265); and Lunik III, which photographed the moon's far side. Only the last of these shots is within the year under review.

The United States has aimed for deep space eight times in all. Four of the American attempts for the moon failed, two of them in the current year. But two earlier shots returned valuable information from high trajectories, and Pioneer IV passed the moon to continue in orbital motion around the sun.

The lone American success achieved in this category during the third year of the space age was a notable one. Pioneer V was sent into an orbit between those of the earth and Venus, thus becoming the third artificial planet. This probe attained the highest velocity of any launched thus far, and its transmissions could be followed out to the unprecedented distance of 22 million miles.

Telemetered data from Pioneer V con-

tinue to be analyzed. Already results of great importance are beginning to appear, and will be reviewed later.



COURIER I-B

AS this issue was going to press, the Advanced Research Projects Agency announced its most recent launching. Courier I-B, sent up from Cape Canaveral on October 4th at 17:50:07 Universal time, is a 51-inch sphere weighing about 475 pounds, with complex equipment for storing and retransmitting radio messages. Known as 1960 ν 1, it was accompanied into orbit by the second-stage vehicle, 1960 ν 2.

According to early figures from Space Track, both bodies circle the earth in orbits inclined 28.3 degrees to the equator. For 1960 ν 1, whose nodal period is 106.9 minutes, the perigee and apogee heights are 587 and 771 miles, respectively.

MARSHALL MELIN

Research Station for Satellite Observation
P. O. Box 4, Cambridge 38, Mass.



GOTO MEDIUM-SIZE PLANETARIUM

The Museum of Art, Science & Industry, Bridgeport, Connecticut is the first institution to install Goto's M-1 medium-size planetarium in its dome. It will open to the public in the fall of this year. The planetarium has won high acclaim when displayed at the 4th annual United States World Trade Fair in May 1960.



GOTO OPTICAL MFG. CO.

1-115 Shimmachi, Setagaya-ku, Tokyo, Japan
Cable Address: GOTOPTIC TOKYO


GETTING ACQUAINTED WITH ASTRONOMY

THE PLANETS — MARS — I

NEARLY all amateur astronomers, whether they are skilled observers or armchair students, will agree that Mars has a fascination shared with no other planet. Among professionals, Mars has been given more attention with large observatory telescopes than Jupiter or Venus.

One reason for this is that Mars is the only celestial body, other than the moon, whose true surface can be studied in detail. Venus, Jupiter, and Saturn are all shrouded in dense atmospheres masking the solid planetary globes, and the only markings seen with certainty are transitory cloud effects. But the thin Martian atmosphere allows us to view a complex pattern of surface features, some of which are recognizable on drawings made three centuries ago. In addition, Mars is the planet whose surface conditions most closely resemble the earth's; of the other bodies in the solar system, it seems the most likely to be an abode of life.

Requiring 687 days to travel once around the sun at an average distance of 141 million miles from it, Mars is a slight-



Above: Oppositions of Mars from 1956 to 1971 are shown in this chart of its orbit and the earth's. The distances are in miles.

Left: An excellent idea of Mars' appearance in a large telescope under favorable conditions is given by this E. M. Antoniadi drawing, made on November 5, 1909, with the 33-inch refractor of Meudon Observatory. Antoniadi (1870-1944) was one of the greatest of all Mars observers. Greek by birth, he did most of his work in France.



ly flattened globe with a diameter only 0.53 that of the earth. It rotates once in 24 hours 37 minutes 23 seconds, around an axis that extends to the celestial sphere not far from Alpha Cephei, which is thus the north star for Mars.

In many respects, the red planet might be called a poor relation of the earth: smaller, less dense, and farther from the sun's light and heat, with a thin atmosphere having very little oxygen or moisture. Perhaps the arid Tibetan plateau is the nearest terrestrial approximation to a Martian landscape.

Since Mars takes about 22½ months to complete one revolution around the

sun, while the earth in its interior orbit requires 12, our planet overtakes the other at intervals of 26 months. When the earth passes Mars, the latter is said to be in opposition. The last date of opposition was November 16, 1958, when the planet's distance from Earth was 45 million miles. On October 30, 1959, Mars was at superior conjunction, on the far side of the sun, and 240 million miles distant. The next opposition date will be December 30th this year, with the planet 57 million miles away.


These figures help explain why useful observations of Mars are limited to periods of a few months, around the times

of opposition. But because of the marked eccentricity of the Martian orbit, not all such occasions are equally favorable. When the planet is at opposition while passing through the perihelion point of its orbit, its distance from us is only 34.5 million miles, but at an aphelion opposition it can be 63 million miles away from the earth.

Especially favorable oppositions occur in a cycle of 15 or 17 years: 1892, 1909, 1924, 1939, and 1956. At such times, the angular diameter of the planet's disk is as great as 25 seconds of arc (about the apparent size of the lunar crater Aristarchus). Such oppositions always occur in August or September, when Mars is well south of the celestial equator. For this reason, perihelion oppositions are best seen from Southern Hemisphere observatories.

The 1960 opposition is relatively unfavorable because of the great distance of the planet, and the maximum apparent diameter of the disk will be only 15.4 seconds. For observers in the United States and Europe, a compensating factor will be the great northerly declination (+27°), so Mars will be high in the sky where telescopic seeing is best.

The minimum distance from Earth will be reached on December 25th, five days earlier than opposition. The cause of this difference is that the motions through space of Mars and the earth in late December are not parallel but slightly divergent. Mars passed perihelion on May



These six photographs of Mars in 1956 were taken by J. H. Botham with the 9-inch refractor of Union Observatory, Johannesburg, South Africa. The south polar cap is conspicuous in the first view, August 25th. In the bottom row, the large dark triangle, near the lowest part of the disk, is Syrtis Major. In the last two exposures, taken two hours apart on September 7th, note how the rotation of the planet has carried Syrtis Major leftward.

26th, and hence its distance from the sun continues to increase.

When Mars is viewed with an adequate telescope under favorable conditions, it presents a small, bright, ochre disk, mottled with darker, greenish shadings, and usually with a brilliant white polar cap. Actually the dark areas are not green, but appear so only by contrast with the bright reddish regions, according to the recent work of G. P. Kuiper.

The first map of the Martian surface that approximates modern standards was drawn by G. V. Schiaparelli, from his observations at Milan during the favorable opposition of 1877. In this map, the Italian astronomer introduced a system of nomenclature still in current use. Dark areas he named after terrestrial seas (for example, Mare Tyrrhenum), bright regions after lands (Hellas), while elongated dark markings — the canals — were called after rivers (Indus). His designations were all drawn from classical geography and mythology.

Many hundreds of names have since been assigned to Martian surface features, real and nonexistent, so the nomenclature has recently been revised by the International Astronomical Union. Since 1958, names have been retained for only about 130 important formations, while lesser features are to be designated by their Martian longitudes and latitudes. The IAU system of nomenclature was presented, with maps, in *SKY AND TELESCOPE* for November, 1958, page 23.

In small telescopes, the Martian maria seem to form a nearly continuous girdle around the southern hemisphere of the planet. But large instruments, under the best seeing conditions, resolve the dark markings into highly intricate tiny

mottlings, as is well shown in the maps accompanying H. Camichel's article on page 600 of the September, 1959, issue.

According to Camichel and other expert observers, most canals can be similarly resolved into finer detail, though some seem to be merely boundaries between areas of unlike brightness. In instruments of lower resolving power, canals of both sorts would be seen, if at all, as dark lines. The manner in which an observer depicts detail at the limit of vision depends strongly on how his mind interprets what his eye barely sees.

Because Mars' equator is tipped 25 degrees from its orbital plane, the red planet has seasons analogous to terrestrial ones. In the Martian northern hemisphere, winter set in on July 2, 1960; spring starts on December 8th, and summer begins on June 26, 1961. During this interval, the north polar cap changes markedly in size; largest near the commencement of spring, it will become very small by late Martian summer.

Meanwhile, conspicuous changes occur in the tint of the dark markings. Near the end of winter, a wave of darkening spreads equatorward, crossing it before mid-spring, and extending into the other hemisphere. This remarkable seasonal variation recurs with much regularity each Martian year. First recognized by Percival Lowell, it has recently been closely studied by G. de Vaucouleurs.

In the planet's southern hemisphere, of course, the seasons are reversed. At the 1960 opposition, Mars' north pole will be slightly tipped toward the earth, and hence the phenomena of the north polar cap will be better visible than those of the south cap.

The amateur who wishes an introduc-

tory book about the red planet should read Patrick Moore's *Guide to Mars* (1956). A thorough survey of modern findings is provided by de Vaucouleurs in his *Physics of the Planet Mars* (1954). For detailed topographical information, the standard authority is E. M. Antoniadi, *La Planète Mars* (1930). Among the *SKY AND TELESCOPE* articles on Mars not mentioned above are: June, 1954, pages 251 and 265; July, 1955, page 360; April, 1956, 256; April, 1957, 266; and July, 1959, 484.

(To be continued)

QUESTIONS... FROM THE S+T MAILBAG

Q. What is the reflectivity of a freshly aluminized mirror in the visible region of the spectrum?

A. For violet light (4000 angstroms) it is 92 per cent, for red (7000) 96 per cent.

Q. What is the *Bonner Durchmusterung*?

A. This is a famous star catalogue, compiled at Bonn Observatory by F. W. Argelander and his associates a century ago. It lists approximate positions of 324,198 stars brighter than nominal magnitude 9.5 (but actually magnitude 11) and north of declination -2° . It is accompanied by an atlas. The catalogue and charts were later extended to -23° by E. Schönfeld of Bonn. The *Bonner Durchmusterung* (BD) catalogue and atlas remain today standard astronomical reference works, and are still in print.

Q. Who holds the record for number of variable stars discovered?

A. By means of photographs Cuno Hoffmeister, of Sonneberg Observatory in East Germany, has found about 5,000 new variables during the past three decades.

Q. What is the greatest number of comets discovered in a single year?

A. In 1947 a total of 14 comets was found.

Q. What is the north polar distance of a star?

A. Its angular distance from the north celestial pole. Thus, a star with an NPD of 70° has a declination of $+20^\circ$, and NPD 110° corresponds to -20° . Some older catalogues, in particular the NGC, used this co-ordinate instead of declination.

Q. When will the asteroid Icarus make its next close approach to the earth?

A. In 1968, when it will pass 3.9 million miles from us. Its minimum possible distance is 3.5 million miles — about 15 times as far as the moon.

Q. How bright would the sun appear if seen from the nearest star and from the Andromeda galaxy, M31?

A. It would be a star of magnitude 0 from Alpha Centauri, and about magnitude $+29$ from M31.

W. E. S.



★ Questar
NEW HOPE • PENNSYLVANIA

A WOMAN WRITES OF HER QUESTAR

"Since receiving Questar, I have been writing you mentally every day to say what you no doubt have heard many times, that Questar is an instrument of such jewellike quality that I fail to find words to express my feelings adequately. It is a pleasure just to lift it out of its case. It so out-modes other telescopes that it is like a leap from the Stone Age to the present time. I cannot praise it highly enough, and I begrudge every cloudy or foggy day that keeps it safely cased.

"To the men who designed this instrument my undying thanks, and that of many others, for Questar is truly progress, and way ahead of its time." (Miss) Julia Wightman, —, New York

STAR-TESTED QUESTARS ARE USUALLY IN STOCK READY FOR IMMEDIATE DELIVERY, FROM \$995. TERMS ARE AVAILABLE. LET US SEND YOU THE QUESTAR BOOKLET.

November, 1960, SKY AND TELESCOPE 279

Planetarium Notes

(Most planetariums give group and special showings by appointment.)

CALIFORNIA

LOS ANGELES
Griffith Observatory and Planetarium
Griffith Park, P. O. Box 27787, Los Feliz Station (27). NO 4-1191. Director, C. H. Clemmishaw.
SCHEDULE: Daily (except Monday), 3 and 8:30 p.m.; also 1:30 and 4:30 p.m. Saturday and Sunday. Zeiss projector.

SAN FRANCISCO
Morrison Planetarium
California Academy of Sciences, Golden Gate Park (18). BA 1-5100. Curator, George W. Buntin.
SCHEDULE: Daily (except Monday and Tuesday), 3:30 and 8:30 p.m.; also 2 p.m. Saturday and Sunday. Tuesday, 3:30 p.m. Academy projector.

SAN JOSE
Rosierian Planetarium and Science Museum
Park and Naglee Aves. Director, Rodman R. Clayton.
SCHEDULE: Sunday and Wednesday, 2 and 3:30 p.m. Spitz projector.

SANTA BARBARA
Gladwin Planetarium
Museum of Natural History, 2559 Puesta del Sol Rd. WO 6-6720. Lecturer, C. Adair.
SCHEDULE: 1st and 3rd Monday, 3 p.m.; 2nd and 4th Thursday, 8 p.m. Admission free. Spitz projector.

COLORADO

COLORADO SPRINGS
Academy Planetarium
U. S. Air Force Academy. GR 2-2779. Director, Maj. Richard J. Pfarr.
SCHEDULE: (Closed until December 10th) Wednesday, 8 p.m.; Saturday, 2:30 p.m.; Sunday, 2 and 3:15 p.m. Admission free. Spitz Model B projector.

DENVER
Denver Museum of Natural History Planetarium
City Park. EA 2-1808. Curator, Robert E. Samples.
SCHEDULE: Daily, 3:30 p.m.; also 12:30, 1:30, and 2:30 p.m. Saturday and Sunday. Spitz projector.

CONNECTICUT

MYSTIC SEAPORT
Seaport Planetarium
Marine Historical Association. JE 6-2631. Director, Frederic W. Keator.
SCHEDULE: Daily, 2 and 4 p.m.; also noon Saturday and Sunday. Spitz projector.

STAMFORD
Edgerton Planetarium
Stamford Museum and Nature Center. DA 2-1646. Director, Ernest T. Lubbe.
SCHEDULE: Saturday, 11 a.m. and 3:15 p.m.; Sunday, 4 p.m. Spitz projector.

FLORIDA

ST. PETERSBURG
St. Petersburg Junior College Planetarium
St. Petersburg Junior College (10). Director, Elizabeth James.
SCHEDULE: Monday, 7:30 p.m. (except school holidays). Spitz projector.

ILLINOIS

CHICAGO
Adler Planetarium
900 E. Acheson Bond Dr. (5). WA 2-1428. Director, Robert L. Johnson.
SCHEDULE: Monday through Saturday, 11 a.m. and 3 p.m.; also 8 p.m. Tuesday and Friday, 10 a.m. Tuesday through Friday (special school program). Sunday, 2 and 3:30 p.m. Zeiss projector.

INDIANA

EVANSVILLE
Koch Planetarium
Evansville Museum of Arts and Sciences, Sunset

Park. HA 5-2406. Curator of education, James Gilliam.

SCHEDULE: Sunday, 2, 3, and 4 p.m. Spitz projector.

INDIANAPOLIS
Holoomb Planetarium
Butler University (7). AT 3-1340. Director, Harry E. Crull.
SCHEDULE: Saturday and Sunday, 4 and 8 p.m. Spitz projector.

IOWA

CHEROKEE
Sanford Museum and Planetarium
117 E. Willow. CA 5-3922. Director, W. D. Frankforter.
SCHEDULE: By appointment daily (except Sunday) during museum hours. Spitz projector.

WATERLOO
Theater of the Stars
Grout Historical Museum, Park Ave. at South St. AD 4-6357. Director, Genevieve Woodbridge.
SCHEDULE: Saturday, 2:30 p.m. Admission free. Spitz projector.

MARYLAND

BALTIMORE
Davis Planetarium
Maryland Academy of Sciences, Enoch Pratt Library Bldg., 400 Cathedral St. (1). MU 5-2370. Director, Paul S. Watson.
SCHEDULE: (Sept.-June) Thursday, 7:15, 7:45, 9 p.m.; Saturday, 2 and 3 p.m. Admission free. Spitz projector.

MASSACHUSETTS

BOSTON
Charles Hayden Planetarium
Museum of Science, Science Park (14). RI 2-1410. Director, John Patterson.
SCHEDULE: Tuesday through Friday, 11 a.m. and 3 p.m.; Friday, 8 p.m.; Saturday, 11 a.m., 2 and 3:30 p.m.; Sunday, 1:30, 2:45, and 4 p.m. Korkosz projector.

SPRINGFIELD
Seymour Planetarium
Museum of Natural History (3). Director, F. D. Korkosz.
SCHEDULE: Tuesday, Thursday, Saturday, and Sunday, 3 p.m.; also 8:30 p.m. Tuesday; 2 p.m. Saturday, special star stories for children. Admission free. Korkosz projector.

MICHIGAN

BLOOMFIELD HILLS
McMath Planetarium
Cranbrook Institute of Science. In charge, James A. Fowler.
SCHEDULE: Wednesday, 4 p.m.; Saturday and Sunday, 2:30 and 3:30 p.m. Spitz projector.

FLINT
Robert T. Longway Planetarium
Flint Junior College, 1310 E. Kearsley St. (3). CE 8-1631. Director, Maurice G. Moore.
SCHEDULE: Tuesday, Thursday, Friday, and Saturday, 8 p.m.; Saturday and Sunday, 2 p.m. Spitz Model B projector.

MINNESOTA

MINNEAPOLIS
Science Museum
Minneapolis Public Library, 300 Nicollet Ave. (1). Planetarium director, Mrs. Maxine B. Haarstick.
SCHEDULE: (Beginning December) Saturday and Sunday, 2, 3, and 4 p.m.; also 10 and 11 a.m. Saturday. Admission free. Spitz Model C projector.

MISSOURI

KANSAS CITY
Kansas City Museum Planetarium
3218 Gladstone Blvd. (23). HU 3-8000. Director, Wilber E. Phillips.
SCHEDULE: Saturday and Sunday, 3 p.m. Spitz projector.

LAQUEY
Tarbell Planetarium
Inca Cave Park. Director, E. D. Tarbell.
SCHEDULE: Daily, 10 a.m.-6 p.m., continuous. Spitz projector.

NEBRASKA

HASTINGS
McDonald Planetarium
Hastings Museum. Supervisor, Lawrence E. Brown.
SCHEDULE: Daily, 4 p.m.; also 2 p.m. Saturday and Sunday. Spitz projector.

NEW JERSEY

NEWARK
Newark Museum Planetarium
49 Washington St. (1). MI 2-0011. Supervisor, Raymond J. Stein.
SCHEDULE: Saturday, Sunday (except 1st Sunday of month), and holidays, 2:30 and 3:30 p.m. Admission free. Spitz projector.

NEW YORK

NEW YORK CITY
American Museum-Hayden Planetarium
81st St. and Central Park West (24). TR 3-1300. Chairman, J. M. Chamberlain.
SCHEDULE: Monday, 2 and 3:30 p.m.; Tuesday through Friday, 2, 3:30, and 8:30 p.m.; Saturday, 11 a.m., 1, 2, 3, 4, 5, and 8:30 p.m.; Sunday and holidays, 1, 2, 3, 4, 5, and 8:30 p.m. Zeiss projector.

NORTH CAROLINA

CHAPEL HILL
Morehead Planetarium
University of North Carolina. Director, A. F. Jenzano.
SCHEDULE: Daily, 8:30 p.m.; also 11 a.m. and 3 p.m. Saturday, 3 and 4 p.m. Sunday. Zeiss projector.

OREGON

PORTLAND
Oregon Museum of Science and Industry
4015 S. W. Canyon Rd. (1). CA 6-4518. Planetarium director, Norman C. Smales.
SCHEDULE: Monday through Thursday, 2:30 p.m.; Saturday, 2 and 3:30 p.m.; Sunday, 2, 3, and 4 p.m. Spitz projector.

PENNSYLVANIA

LANCASTER
North Museum and Planetarium
Franklin and Marshall College. Curator, John W. Price.
SCHEDULE: Tuesday and Thursday, 8 p.m.; Saturday and Sunday, 2 and 3 p.m.; Monday, Wednesday, and Friday, 9:15 and 10:15 a.m. (special school program). Admission free. Spitz projector.

PHILADELPHIA
Fels Planetarium
Franklin Institute, 20th St. at Benjamin Franklin Parkway (3). LO 4-3600. Director, I. M. Levitt.
SCHEDULE: Tuesday through Sunday, 3 p.m.; also Wednesday and Friday, 8 p.m.; Saturday, 11 a.m.; Saturday, Sunday, and holidays, 2 p.m. Zeiss projector.

PITTSBURGH
Buhl Planetarium and Institute of Popular Science
Federal and W. Ohio Sts. (12). FA 1-4300. Program director, Arthur L. Draper.
SCHEDULE: Daily, 2:15 and 8:30 p.m.; also 11:15 a.m. Saturday, 4:15 p.m. Sunday. Zeiss projector.

RHODE ISLAND

PROVIDENCE
Roger Williams Planetarium
Roger Williams Park Museum (5). WI 1-5640. Director, Maribelle Cormack.
SCHEDULE: (Oct.-May) Saturday, 3:30 p.m.; Sunday and holidays, 3 and 4 p.m. Admission free. Spitz projector.

TENNESSEE

CHATTANOOGA
Clarence T. Jones Observatory
University of Chattanooga, Brainerd Rd. MA 2-5733. Astronomer in charge, Karel Huijter.
SCHEDULE: Friday, 8 p.m. Admission free. Jones projector.

MEMPHIS
Memphis Museum
233 Tilton Rd. at Central Ave., Chickasaw Gardens (11). GL 2-4732. Director, Mrs. Ruth C. Bush.
SCHEDULE: Saturday and Sunday, 2 and 2:45 p.m.; also 3:30 p.m. Sunday. Admission free. Spitz projector.

NASHVILLE
Sudekum Planetarium
Children's Museum, 724 2nd Ave. S. (10). CH 2-1858. Director, Jacqueline Avenet.
SCHEDULE: Sunday, 2:45, 3:30, and 4:15 p.m. Spitz projector.

TEXAS

DALLAS
Dallas Planetarium
Dallas Health and Science Museum, Fair Park (10). HA 8-8351. Director, M. L. McDonald.
SCHEDULE: Saturday and Sunday, 3 p.m. Spitz projector.

FT. WORTH
Charlie M. Noble Planetarium
Ft. Worth Children's Museum, 1501 Montgomery. FE 2-1461. Supervisor, Norman C. Cole.
SCHEDULE: Saturday and Sunday, 2:30 and 3:30 p.m.; also 11 a.m. Saturday. Spitz projector.

WEST VIRGINIA


CHARLESTON
Hillis Townsend Planetarium
Children's Museum, Public Library Bldg. Director, Mrs. R. L. Sullivan.
SCHEDULE: Saturday, 11 a.m. Admission free. Spitz projector.



The latest
Spitz MODEL A-2
PLANETARIUM

has been installed at
Danbury State College
Danbury, Connecticut

★
Spitz Laboratories
Inc.
YORKLYN, DELAWARE
Phone: CEder 9-5212



The Many Faces of a Telescope Maker

Tinsley Laboratories was founded 31 years ago to build reliable, high quality telescopes for amateur and professional astronomers. Today, the emblems you see here represent Tinsley's growth—company divisions engaged in significant scientific, industrial, military and educational applications of optics.

Reading down, these divisions are: **TELESCOPES**, including professional observatory instruments up to 60 inches; school and advanced amateur 'scopes such as the new 12-inch; and terrestrial telescopes for missile tracking and guidance... **WIND TUNNEL AIR FLOW VISUALIZATION**. Since 1942, Tinsley has designed and built complete Schlieren, Interferometer and Shadowgraph systems for the aerospace industries... **OPTICAL SURFACES**: mirrors up to 72 inches diameter, prisms and lenses... **OPTICAL TOOLING**, the development of sensitive measuring instruments, such as portable autocollimators, for product quality control.

These divisions add up to a firm with a wide spectrum of capabilities in research, design, engineering and manufacturing of optics: **TINSLEY LABORATORIES, INC.** 2526 Grove Street, Berkeley 4, California. For a copy of our new Facilities Folder, write Dept. R.

EYES TO THE SKY, Tinsley's new 16mm full-color film on telescope manufacturing is available for showings • Address requests to the Audio-Visual Department

OBSERVER'S PAGE

Universal time (UT) is used unless otherwise noted.

THE SEPTEMBER PARTIAL ECLIPSE OF THE SUN

SEPTEMBER was a banner month for amateur eclipse observers in North America, the moon being totally obscured on the 5th and the sun partly covered September 20th. Not until the year after next will another solar eclipse be visible from the United States — a partial event for Californians at sunset on February 4, 1962, followed by another for Floridians at dawn on July 31, 1962.

For this year's partial solar eclipse, scattered areas of cloudy weather somewhat hindered the efforts of North American amateurs on the afternoon of September 20th, but those in the south-

western and western states were generally rewarded by a good view. In California and Texas, especially, numerous drawings and photographs were obtained.

In the Middle West, where the eclipse was a late afternoon event, high thin cirrus clouds obscured the view shortly before sunset for amateurs at O'Fallon, Missouri. However, five minutes earlier a good bright projected image was furnished by an 8-inch reflector, the sharp edge of the moon appearing clearly, as well as many sunspots.

Equipment used for pictures that were sent to this magazine ranged from simple



Star Instrument's*

Highly Portable, Fully Equipped
8-inch f/6 Reflector.....\$589.45
(8" instruments from \$311.00)

TF OPTICAL LABORATORIES

P. O. Box 481

Huntington Beach, Calif.

*Trade Mark TF Optical Laboratories

Deep-Sky Observers



An extremely fine, imported Zuiho binocular with a wide field of view — 10½°!

\$35.00
Tax Paid

Complete with beautiful case. Delivered anywhere in the Continental United States.

AMERICAN TYPE WIDE ANGLE
7 x 35 Central Focus

SPECIFICATIONS

Magnification, 7x. Objective aperture, 35 mm. Eyepiece aperture, 24 mm. Exit pupil, 5 mm. Field of view at 1,000 yds., 551 ft. Field-of-view angle, 10.5°. Height, 128 mm. Breadth, 195 mm. Net weight, 950 grams. Weight of pigskin case, 400 grams.

We have sold hundreds of pairs of these binoculars with the utmost satisfaction. They are unconditionally guaranteed. If not pleased, return them and your money will be immediately refunded.


Before you buy or trade a telescope or radio amateur gear, see **WARD W2FEU** for the best deal.




Established 1936

ADIRONDACK RADIO SUPPLY

185-191 West Main St.
P. O. Box 88, Amsterdam, N. Y.
Ward J. Hinkle, W2FEU — Owner



To show the progress of the eclipse at Ft. Worth, Texas, Burton G. Ford made an exposure every five minutes, from 5:30 p.m. Central standard time.



This 1/500-second exposure by Donald E. Lazenby was made at the prime focus of his homemade 10-inch reflector in North Hollywood, California. He reduced the sun's light with a 2½-inch aperture stop and a dense neutral filter. The film was Adox KB-14, developed in X-22. Sunspots, bright faculae, and solar limb darkening are all shown, while the irregularities in the edge of the moon are conspicuous against the disk of the sun.

cameras to the elaborate instruments of the Lockheed solar observatory in Burbank, California. For most places in the United States, the moon obscured between 15 and 25 per cent of the sun's diameter at maximum eclipse. At Ft. Worth, Texas, Burton G. Ford timed the maximum at 6:07 p.m. Central standard time, while Don Lammon, San Bruno,

California, noted it at 3:40 p.m. PST, for his location. The latter recorded an obscuration of about 25 per cent.

Two major precautions to dissipate excess light were taken by Jack Eastman, Jr., of Manhattan Beach, California, when photographing the eclipse directly with his 12½-inch reflector. He replaced the aluminized diagonal with an objective



One excellent way to observe a solar eclipse (and the transit of Mercury this month) is being demonstrated by Charles Rothermich to his daughter, Mary, at O'Fallon, Missouri. The solar image is here projected through the eyepiece of an 8-inch Cassegrainian onto a white screen, allowing observation by several people at one time. This method has the advantage that no filters are necessary. Photograph by Robert E. Cox.

Astronomy Teaching Aids

ESSCO Publications

Teachers of astronomy, both amateur and professional, will find the following ESSCO study materials useful in their classrooms. The test charts are challenging fun for the amateur astronomer, too!

- SC1 Equatorial constellation chart — with star designations
- SC1T Test equatorial chart — without star or constellation names
- SC2 Circumpolar constellation chart — with star designations (new large size)
- SC2T Test circumpolar chart — without star or constellation names (new large size)
- SS08A Ecliptic-based star map — with equatorial grid and names
- SS08 Ecliptic-based star map — with equatorial grid, without names
- SS08B Ecliptic star map list — positions and magnitudes for 224 stars
- SS05 Nine-inch protractor on paper — for planet orbit drawings
- SS11 Inner planet chart — orbits of Mercury, Venus, Earth, Mars
- SS12 Outer planet chart — orbits of Mercury to Saturn
- SS01A Special rectangular co-ordinate paper — for star maps
- SS02 Polar co-ordinate paper — for circumpolar star maps

Price for each item listed above: 1 to 9 sheets, 10 cents each; 10 to 24 sheets, 8 cents each; 25 to 99 sheets, 6 cents each; 100 or more, 5 cents each.

From Stetson's *Manual of Laboratory Astronomy*, the chapter "Star Chart Construction" is available as a separate booklet, at 50 cents per copy.

SPECIAL: DC5 — Large wall chart of the Draper spectral classes, printed in four colors and mounted on canvas. Limited quantity. (Slightly damaged along edges.) **\$14.00 each**

SKY PUBLISHING CORPORATION

Cambridge 38, Massachusetts



Celestial prints and wall murals.

Write for free brochure #1

astro murals

33 West 60th Street, New York 23, N. Y.

GUARANTEED FINEST . . . OR YOUR MONEY BACK



Pat. Pend.

SKY-MICRO REGULAR 1.25" mount with exclusive MICRO-VERNIER focusing • 360° rack with hobbled pinion • Stops on both tubes • Phenolic drag • Large aluminum knobs • Contoured base (specify tube diam. or "flat" base) • Adjustable hex bushings • Complete with screws, instructions...\$19.95 ppd.

SKY-MICRO GIANT (new) 2" mount (same features as above) includes 1.25" adapter...\$36.95 ppd.
SPACEKING 1.25" spiral focus mount...\$9.95 ppd.

HEX-LAP MATS

Custom die-cut for each mirror size. Makes a perfect pitch lap in minutes. Gives superior results in polishing and figuring. Finest quality rubber. 4.25" or 6" \$2.50 ppd.
8" or 10" \$3.50 ppd.
12.5" \$4.00 ppd.



G. M. LABORATORY

(Dept. H) 1359 Romulus Drive, Glendale 5, Calif.



RADIOMETER

American made. Balanced four-arm vane, in a 3"-diameter evacuated glass globe, turns upon exposure to light, any light, even a burning match. Highly decorative, interesting, educational, instructive. A conversation piece. We guarantee yours to operate.

\$3.00 ppd.; 2 for \$5.00 ppd.

Jupiter BLUE Filter

A useful accessory for your telescope, especially for viewing the Red Spot on Jupiter. Improves color contrast, reduces glare. Helps you to see detail in the belts of this planet. 35-mm. diameter, 1-mm. thick. Polished plane parallel. Famous manufacturer. Unmounted.

Only \$1.50 each, ppd.

HARRY ROSS

Scientific & Lab Apparatus
61-H Reade St., New York 7, N. Y.

NEW FALL ITEMS

THE EXPLORATION OF SPACE

Edited by Robert Jastrow. This book covers the symposium on space physics, sponsored by the National Academy of Sciences, the National Aeronautics and Space Administration, and the American Physical Society, on April 29-30, 1959. The aspects of space exploration are covered very thoroughly. Up to date in the very truest sense. Note these authors: F. L. Whipple, T. Gold, E. Parker, J. A. Van Allen, N. C. Christofilos, H. E. Newell, G. P. Kuiper, H. C. Urey, G. de Vaucouleurs, H. Friedman, L. Goldberg, J. W. Townsend, Jr., and R. Jastrow.

Only \$5.50

PHOTOGRAPHIC LUNAR ATLAS

Edited by Gerard P. Kuiper. 230 sheets, 15" by 20", showing 280 views of the moon, taken at Mount Wilson, Lick, Pic du Midi, McDonald, and Yerkes observatories. 24-page booklet included. The cost is pennies per picture.

Only \$30.00

MOON MAPS

By H. P. Wilkins. 65 pages, including a map of the far side of the moon. Spiral bound.

Only \$6.00

"SEE-THRU" SKY MAPS

12 8 1/2" charts on clear plastic with luminous paint. Hold one of the maps the right distance from your face to match the stars in the sky. 1960 co-ordinates. Instructions and identifications, so that you can find the right map instantly for any section of the sky and time of the year. Orientation diagram included. New supply — order now — they go quickly!

Only \$5.25

THE SOLARSCOPE

Imported from England. Two rotary disks give you instant information about the sun, moon, and planets — at a twist of the wrist.

Only \$2.25

It's not too early to think about Christmas gifts! Write for new catalogue just printed.

ASTRONOMY CHARTED

33 Winfield St., Worcester 10, Mass., U.S.A.
Phone: PL 5-6992




This eclipse photograph in red hydrogen-alpha light (turned left-right from the other pictures in this article) was taken through a narrow-band birefringent filter at 4:00:30 p.m., PST, by J. W. Harvey, at the Lockheed solar physics observatory, in Burbank, California.

prism mounted as a Herschel wedge, and then put a No. 11 welder's glass between the wedge and the camera. The wedge passed about four per cent of the incident light, and the filter reduced this by a factor of around 30,000. With this reduction, he could use an exposure time of 1/25 second in direct photography.

Other reports and photographs of the eclipse were received from: John Harvey, Burbank, Calif.; Lewis Chilton, Los An-

geles, Calif.; John Shea and Earl Bartsch, Bell, Calif.; Ken Hoover, Long Beach, Calif.; James Williams, Woodland Hills, Calif.; Fred Larsen, Los Angeles, Calif.; D. Colin Wyatt, Victoria, British Columbia; Clyde Honstein, Jr., El Paso, Tex.; William Vance, Kingsville, Tex.; John Marshall, Elmhurst, Ill.; Maurice Moore, Flint, Mich.; Donald Lazenby, North Hollywood, Calif.; Roderick Dunphy, Monta Vista, Calif.; G. Vlahakis, Jr., Salem, Wis.; and David Dunham, Berkeley, Calif.




A camera held at the eyepiece of his 4 1/2-inch reflector gave Jack Eastman, Manhattan, California, this result.

OPTICAL FLATS

Pitch Polished, Beral Coated

Rectangular shape

1 3/8" x 1 7/8"
1/2 wave \$1.75 each
1/4 wave \$4.50 each
Postpaid.



Elliptical shape

Heavy edges to minimize temperature effects.
1 1/4" minor axis.
1/2 wave \$4.00 each
1/4 wave \$6.50 each
Postpaid.



BERAL COATINGS — same optical characteristics as aluminum — mechanically more durable — not over-coated — may be removed without harming glass surface. Prices for Beral coating telescope mirrors: 4" diameter — \$3.50, 6" — \$5.00, 8" — \$6.50, 10" — \$8.50, and 12 1/2" — \$12.50 each, f.o.b. Skokie.

LEROY M. E. CLAUSING

8038 Monticello Ave.

Skokie, Ill.

OBSERVE MARS THIS YEAR AT CHRISTMAS — with your New ASTROLA Reflecting Telescope



Mars on October 13, 1958, as seen through an 8" f/7 De Luxe ASTROLA Reflector. The drawing was made when the Martian longitude of the central meridian was 210°-215°, and the disk was 17.2 seconds of arc in diameter. The power used was 360-450, with seeing fairly good, but transparency only fair.

AMERICAN MADE — AVAILABLE NATIONALLY ON TIME PAYMENTS

ON CHRISTMAS EVE, next month, Mars will be closest to the Earth, and if you have just received your new ASTROLA, you will be doubly thrilled observing the planet with an instrument of truly adequate aperture and resolution. Like the many hundreds of present ASTROLA owners, you will be able to see all the dark maria and other deserts, and marvel at the changing sizes of the polar caps. When seeing is really good, the Martian canals and a multitude of fine maria details will flash into view. You will see all this and more as only the superb definition of an ASTROLA can reveal. Mars will be a source of fine observing for weeks before and after Christmas, too.

When the planet finally becomes too distant to observe, you can discover minute features on the moon, obtain brilliant views of other planets, and examine thousands of deep-sky wonders with your ASTROLA Reflector. REMEMBER — ASTROLAS are manufactured and sold by Cave Optical Co. and are not imported telescopes.

There is an ASTROLA for every budget, from our 6" STUDENT or STANDARD to our massive 12½" OBSERVATORY telescope. On special order, we also manufacture larger custom instruments. All ASTROLA models include a large finder and three or more Brandon oculars. De Luxe models include clock drive, setting circles, and rotating tube. Truly ASTROLA IS AMERICA'S MOST WANTED REFLECTING TELESCOPE.

Christmas shoppers: Portable-model ASTROLAS ordered by November 10th will be shipped for delivery by December 25th.

PRICES FOR COMPLETE ASTROLA TELESCOPES*

	Standard	De Luxe
6" STUDENT, f/7, f/8, or f/9	\$194.50
6" MODEL "A", f/7 or f/8	\$325.00	\$500.00
8" MODEL "B", f/6, f/7, or f/8	\$390.00	\$590.00
10" MODEL "C", f/6, f/7, or f/8	\$490.00	\$750.00
12½" MODEL "D", TRANSPORTABLE, f/6 or f/7	\$1150.00
12½" MODEL "D", OBSERVATORY, f/6, f/7, or f/8	\$1780.00

*Slight additional charge for packing and crating.

BRANDON ORTHOSCOPIC OCULARS: We are the sole West Coast distributor for these eyepieces. Available in 32-mm., 24-mm., 16-mm., 12-mm., 8-mm., 6-mm., and 4-mm. effective focal lengths. Price, \$15.95 each, ppd.

CAVE ACHROMATIC and COATED 3X BARLOW, in mount \$20.00 ppd.

PRICES FOR CAVE'S NEW PARABOLIC MIRRORS MADE OF PYREX-BRAND GLASS AND FOR REFIGURING IMPERFECT MIRRORS

Our finished mirrors of highest quality include elliptical diagonal and aluminized. Refiguring imperfect mirrors is our specialty.

	New Mirror	Refiguring
6"	\$60.00	\$35.00
8"	\$92.50	\$50.00
10"	\$160.00	\$75.00
12½"	\$250.00	\$120.00


RAPID DELIVERY on new 6", 8", and 10" mirrors.

Send for new 1961 catalogue in color.

CAVE OPTICAL COMPANY

4137 E. Anaheim St., Long Beach 4, Calif.

Phone: GENEVA 4-2613



The Completely New De Luxe Model "C" 10-inch ASTROLA, f/7, complete with rotating tube, clock drive, setting circles, and 4 oculars (74x, 111x, 222x, 444x, or 4 of your choice).
\$750.00

Astronomy Films

16-mm. sound, 400-foot reels
I THE SUN; II THE MOON;
III SOLAR SYSTEM; IV MILKY WAY;
V EXTERIOR GALAXIES.

2 x 2 SLIDES

35-mm. STRIPS OF SLIDES

THROUGH 200-INCH AND
OTHER GREAT TELESCOPES

Catalogues on request.

**International
Screen Organization**

1445 18th Ave. North, St. Petersburg 4, Fla.


LUNAR RILLS IN EASTERN MARE TRANQUILLITATIS

AMATEUR observers of the moon seem to have devoted surprisingly little study to the very extensive system of rills in the eastern part of Mare Tranquillitatis. Perhaps the most important paper devoted to it is that by the French selenographer C. M. Gaudibert (*L'Astronomie*, 7, 182, 1888), although he describes only a little of what is known at the present time. There have been many short notes about parts of the rill system, for example, that by R. Barker in the British periodical, *The Moon* (Vol. 3, No. 1, page 7, 1954). He was, however, concerned only with the southern rills.

Certainly there has not been the atten-

tion given to the Mare Tranquillitatis rills that has been given to, say, the famous system just west of the crater Triesnecker.

The accompanying chart summarizes my observations from 1955 to 1960, seven of them with a 7-inch reflector, two with a 12½-inch reflector, and one with the 18-inch refractor of the University of London Observatory. Most of the finer details were found with this last instrument.



This map should be compared with H. P. Wilkins', on page 68 of the 1955 book, *The Moon*, by him and Patrick Moore. Rills 2 and 7-13 are omitted by him. In addition, 3 and 4 are actually not connected, nor are 5 and 6. On my chart some of the rills are not linear. There is a tendency among observers always to draw rills as regular objects, whereas in fact many of them meander irregularly, for example, 3. The deviations are usually visible only in large telescopes.


Two other interesting features, believed to be new, are the long, broad, shallow valley that runs southward from Arago, and the row of *elongated* craterlets west of Manners.

I have prepared a chart of the rills near Plato, which is to appear in the *Journal of the British Astronomical Association*, and it shows that many of them are radial to Plato and others to the crater Plato D. Similar examples may be seen in the Mare Tranquillitatis system, where numbers of rills appear to diverge from Ritter and from Dionysius.

Dotted regions on the present map represent domes that I am sure exist. Many small hills and ridges have been omitted for clarity.

BRIAN WARNER

2 Diamond Cottages
Crawley Down, Sussex, England




Magnusson Telescopes and Accessories

- ★ Mountings as low as \$29.00 and \$59.00.
- ★ Low-priced clock drives and other accessories.

Parts are sold separately. Write for prices.

SETTING CIRCLES

Made of choice aluminum or brass, machined and polished all over. Hour circles, machine-scribed with hour, half-hour, and five-minute marks. Declination circles scribed in degrees 0-90-0-90. Numbers stamped with ⅛" dies. Holes reamed standard sizes. State sizes.




	Aluminum	Brass
5" circles, set of two	\$15.00	\$19.50
6" circles, set of two	\$17.00	\$22.10
8" circles, set of two	\$24.00	\$31.20

O. MAGNUSSON

14570 West 52nd Ave.
Arvada, Colorado

Attention: TELESCOPE BUILDERS



TELE-SCHMIDT

The new aspheric correcting lens whose precise contours are shown by the interferogram — is available to the amateur telescope builder at a price beginners can afford.

You can now make your **first** telescope an **8-inch f/7**, the size general-

ly considered as ideal for amateur astronomers, with complete assurance of satisfying success, because **TELE-SCHMIDT eliminates the need for parabolizing the primary mirror** — it is made spherical.

Furthermore, your instrument will be of the very latest catadioptric design, having a wider corrected field and a closed tube. Price? Only \$62.50.

Write now for complete specifications and special introductory offer. Group purchase discount is available — dealer inquiries are invited.

OPTRON LABORATORY

Box 25, D. V. Station
Dayton 6, Ohio

U.S. SURPLUS BARGAINS

IN HIGH-QUALITY OPTICAL PARTS AND EQUIPMENT




**BAUSCH & LOMB f/6
AERO-TESSAR LENS**

Focal length 24". Completely mounted with iris and shutter. Excellent for wide-field telescope. All in excellent condition. Price **\$25.00**



**WIDE-ANGLE ERFLE EYEPIECE
(68° Field)**

Coated optics. 3 achromatic lenses. 1 1/4" focal length, 1-13/16" aperture. Price **\$12.50**



**16-mm. AERIAL GUN
CAMERA, MODEL AN-N6**

Takes standard 16-mm. magazine load (50'). Operates at 16, 32, or 64 frames per second. Complete with Wollensak type V lens, 1 1/4" (35-mm.), f/3.5. Excellent condition. Operates on 24-volt d.c. Price **\$22.50**




METAL PARABOLOIDAL MIRROR

These are made of hastelloy hard metal. 12" diameter. 5 1/4" focal length. Manufactured by Bausch & Lomb. Approx. weight, 3 1/2 lbs. Brand-new condition. Price **\$10.00** each

**LENS ASSEMBLY
48" FOCAL LENGTH — f/6.3**

This was used with the K40 aerial camera, which takes a 9" x 18" plate. Manufactured by Eastman Kodak. Approx. dimensions, 14" diam., 26" long. Contains built-in filters and iris. Lenses mounted in aluminum housing. Approx. weight of unit, 125 lbs. All in original crates. Approx. shipping weight, 200 lbs. Price **\$150.00**



**BELL & HOWELL
f/8 TELEPHOTO LENS**

Coated lenses. Focal length 36". Completely mounted with iris and shutter. Approx. weight 25 lbs. Excellent condition. Price **\$49.50**

8-DAY AIRCRAFT CLOCK



24-hr. dial and civil-date indicator. Center sweep second hand. Luminous figures and hands on black face. Case is made of black plastic. 3 1/2" mounting. Manufactured by Waltham Watch Co. Price **\$20.95** postpaid

8-POWER WIDE-FIELD ELBOW TELESCOPE



COATED LENSES

Field of view 8° 45'. Large focusing Erfle eyepiece. Eye lens diameter 2-1/16" with diopter scale +2 to -4. 4 built-in filters — clear, neutral, red, amber. Length 15 1/2", width 6 3/4", height 7 3/4", weight approx. 9 1/2 lbs. Excellent unused condition. Price **\$39.50**

8-POWER ELBOW TELESCOPE

2" objective; Kellner eyepiece; Amici erecting prism; 4 built-in filters; reticle illumination; field is 6° (325 ft. at 1,000 yds.). The focusing 28-mm. eyepiece allows focus from 15 ft. to infinity. Weight 5 lbs.



These were used by the government for anti-aircraft. Government cost over \$200.00. Price **\$12.50**

**5" SCHMIDT
ULTRA-HIGH-SPEED
OBJECTIVE LENS SYSTEM**


Eastman Kodak infrared receiver, formerly known as U. S. Navy Metascope. Type B. 7" long with 5" Schmidt ultra-high-speed objective lens (approx. f/0.5). Elaborate optical system, many coated lenses. Uses two penlight batteries. Government cost, approx. \$300.00. Factory-new. Shipping wt. 9 lbs. Price **\$19.95**



Waterproof carrying case extra. Shipping wt. 3 lbs. Price **\$3.00**


**NAVY INFRARED RECEIVER,
TYPE C-3**

Produces visible image from infrared image in complete darkness. For seeing in dark with infrared light source. Nothing to wear out. Operates on two flashlight batteries. Has high-voltage power supply and image tube (1P23) and elaborate optical system. Weighs only 6 1/4 pounds and is only 11" long, 8" high, 5 1/2" wide. Manufactured by R.C.A. Objective f/0.9 Schmidt system, 2.4" focal length, 3 1/4" aperture. Excellent condition, with waterproof carrying case. Price **\$35.00**



INFRARED FILTERS

5 1/2" diam. x 1/8" thick. Price **\$2.95**



16-mm. SINGLE-FRAME CAMERA

Manufactured by Bell & Howell (200 series). 50' magazine load. Operates on 12-volt d.c. Has rotary solenoid, standard C mount, lens and view finder. Excellent for photographing records, nature study, and so forth. Price **\$35.00**

**K24 AERIAL CAMERA
WITH 7" f/2.5 KODAK
AERO-EKTAR LENS**



Camera weighs 26 lbs. Size: 10" x 11" x 16". Removable magazine. Picture size: 5" x 5". Diaphragm adjustment from f/2.5 to f/16. Film may be advanced manually or by 24-volt d.c. motor.

Used, in good condition Price **\$50.00**
New Condition Price **\$79.50**

KODAK AERO-EKTAR

f/2.5,


6" f.l.



Mounted in barrel. Hard-coated lenses, color corrected. Complete with dust caps. Red and yellow filters. Price **\$25.00**

**BAUSCH & LOMB f/6, 24" f.l.
AERO-TESSAR
LENS SET**


The clear diameter of front lens is 4"; that of rear lens is 3 3/4". The set consists of two metal-mounted elements. Price **\$20.00**



**AERO-EKTAR
f/2.5, 12" f.l.**

Completely mounted with shutter. Originally manufactured for night-photography aerial camera. Film size, 9" x 9". Excellent condition. Price **\$50.00**

**OBSERVATION INSTRUMENT THAT
CONTAINS ELBOW TELESCOPE M3**



May be varied from 10 to 20 power. Field of view at 20 power, 3° 5'; at 10 power, 6° 9'. Objective lens, 2 1/4" in diameter. Approx. size, 17" x 14 1/2". Approx. weight, 14 lbs. Uses penta prism and erecting lens. Telescope altazimuth mounted. Horizontal 360° scale calibrated in 1/10 mil; altitude 90° scale calibrated in mils; precise adjustments made by turning knob. Azimuth circle and telescoping tripod contain spirit levels. Government cost over \$1,425.00. Approx. shipping wt. 115 lbs. Price **\$125.00**

Send full amount with order. All prices, except as noted, net f.o.b. Pasadena, Calif.
No C.O.D.'s, please.

C & H SALES CO.


2176 East Colorado St.
Pasadena 8, Calif. MURRAY 1-7393

THANKS TO YOU . . . DEMAND FOR OUR NEW GALAXY SERIES IS SO GREAT WE ARE ABLE TO REDUCE OUR PRICES!

Now you can buy the **FINEST** at the greatest savings ever!

Buy up to or trade up to a **TRECKERSCOPE!**

Now, as in the past, Coast Instrument's high quality is the standard of comparison! The American telescope company that provides 1/20th-wave optics for amateurs and professionals alike again is way out in front with the GALAXY. No other telescope manufacturer today offers as many outstanding features. These superb instruments are guaranteed to resolve to, or surpass, Dawes' limit on nights of good seeing. The GALAXY optics are hand-figured and, therefore, our instruments are able to produce results far better than expectations. All of the GALAXY telescopes include these outstanding features — rock-steady equatorial mount; quickly detachable legs for easy portability (only 70 seconds to assemble or disassemble entire scope for field trips); hydro-smooth rotating tube with built-in ocular rack; remote-assistance light; Opti-Set finderscope (only requires 3" movement of head from finder to eyepiece, which is directly in line); Hydro-Glide (ultra-smooth, no backlash) focusing device; fiberglass tube; your choice of any three of the finest-quality oculars (Ortho-Stars), 7 mm. — 10 mm. — 16 mm. — 20 mm. — 27 mm., or any two oculars and a Coast Achromatic Barlow Lens; synchro-smooth clock drive with manual flex-line control for precise tracking when no 110-volt power is available; wheels for ease in moving from location to storage area (wheels snap out for viewing). Leveling screws are provided for any minor difference in ground level. All in all, your dollar cannot purchase better viewing. All of our instruments are American made, and you are protected by our 20-year unconditional guarantee.



GALAXY 6 INCH

GALAXY 6"	\$ 495.00
GALAXY 8"	\$ 595.00
GALAXY 10"	\$ 795.00
GALAXY 12.5"	\$1095.00
STANDARD 6"	\$ 325.00
STANDARD 8"	\$ 425.00
STANDARD 10"	\$ 545.00
6" PATHFINDER	\$ 189.50
10" CASSEGRAIN	
SKY GIANT	\$1795.00

**OUR EXTENDED
PAYMENT PLAN IS
OFFERED TO ALL
TAKE MONTHS TO PAY**
6 — 12 — 18 — 24
IF DESIRED . . .

**WRITE FOR OUR LAB
SPECIFICATION SHEET**

PARABOLIC MIRRORS, GUARANTEED UP TO 1/20TH WAVE OR BETTER FROM COAST'S OWN OPTICAL SHOP, ALUMINIZED AND QUARTZ OVERCOATED

	6" f/10	6" f/8	8" f/7.5 or f/6	10" f/7.5 or f/6	12 1/2" f/7.5 or f/6
1/20th WAVE	\$75.00	\$105.00	\$195.00	\$295.00	\$235.00
1/10th WAVE	\$55.00	\$85.00	\$150.00	\$235.00	\$235.00
1/8th WAVE	\$40.00				

REFIGURING, NO CHANGE IN YOUR OWN FOCAL LENGTH

1/20th WAVE	\$47.50	\$65.00	\$110.00	\$175.00
1/10th WAVE	\$37.50	\$55.00	\$80.00	\$125.00

ALL PRICES ABOVE INCLUDE MATCHING ELLIPTICAL DIAGONALS AND ALUMINIZING WITH QUARTZ OVERCOATING.

TREAT YOURSELF TO THE FINEST — Words alone cannot express the superb quality of COAST mirrors — they will positively amaze you with their "out-of-this-world" resolving power. This we guarantee you! Dawes' limit is easily reached.

TELESCOPE OPERATING MANUAL by Raymond Dudley

Explains amateur telescope operation, including setting circles, sidereal time, alignment, orientation. Shows how to find 52 most beautiful objects in sky as well as general "how-to-observe" instructions. 44 pages of information every amateur needs.

\$1.00 postpaid

HELAJUST FOCUSING MOUNT



Now at a new low price, but unexcelled in construction. Both push-pull for coarse adjustment, and helical movement for fine, precise focusing. Beautifully finished in black crackle and satin chrome, with a total extension of main tube and drawtube of 6", and engineered to fit all sizes of tubes without modification. Has single-vane-spider holding boss for those desiring this method of using diagonal. Guaranteed far superior to any other low-priced focusing mount, or money back. (Takes 1 1/4" eyepieces.) \$9.95



ORTHO-STAR OCULARS


**DUST SEALED — ALL COATED
BEAUTIFUL FINISH**

Guaranteed to be the finest you ever used — or return for full refund! Outstanding features: wide flat field, sharp to the very edge; extra-long eye relief; parabolized for easy change of power; sealed-in optics, never need interior cleaning; hard coated, magnesium fluoride; boldly marked for easy identification; striking chrome and black-velvet finish, beautifully machined, 1 1/4" O.D.

ORTHO-STAR oculars are available in the following focal lengths, giving, for example, the indicated powers when used in conjunction with an 8" f/8 mirror: 26.6-mm.—61x; 20-mm.—81x; 16-mm.—102x; 10-mm.—163x; 6.6-mm.—233x.

\$19.50 each ppd.

NOW AVAILABLE: 20-MM. ERFLE ORTHO-STAR OCULAR, 66" apparent field. \$29.50 each ppd.



TRECKER-PATHFINDER

EQUATORIAL MOUNT


\$89.50

complete

This mount will accommodate 4-inch to 8-inch telescopes. Specify your tube size when ordering.

Standard 36-inch height — massive 1 1/2-inch steel shafting.

This amazing EQUATORIAL MOUNT is just what the doctor ordered for mounting that homemade telescope you labored so hard to finish. Now you can purchase a beautifully constructed, highly rigid equatorial mount, COMPLETE, for your own telescope as economically as if you had built it yourself. This terrific mount is made entirely of metal; all of the moving equatorial parts are polished to work with maximum ease. Legs, head, and counterweight are all removable for easy storing. The saddle allows complete rotation of your tube. One of the more important features in this mount is that the polar axle is extended for ease in attaching a clock drive and/or setting circles, which may be added at any time. The TRECKER-PATHFINDER mount also has a beautiful, chip-resistant finish. Taking all of these unusual features into consideration, this is truly one of the best DOLLAR-FOR-DOLLAR values.



HYDRO-GLIDE

**Precision
Focusing Device**

Coast Instrument's own "Hydro-Glide" (type formerly referred to as "rack and pinion"). Now you can have WHISPER-SMOOTH control with absolutely no high and low spots as geared units have. This has never before been available in any focusing mount. Standard in appearance, yet utilizing an entirely new concept. We guarantee you will be astonished at its unbelievable superiority. If this isn't the absolute "smoothest," return it for full refund.

\$18.50 patent pending



MIRROR CELLS

Skeleton type

6"	\$6.95	10"	\$17.95
8"	\$10.95	12 1/2"	\$21.00




4-VANE SPIDERS

6"	\$11.50	10"	\$14.95
8"	\$11.50	12 1/2"	\$16.95

TRECKER FINDER

7x, 50-mm. objective, helical focusing, with mounts and crosshairs. Same as used on TRECKERSCOPES. \$18.50



Send for Coast Instrument's Professional MIRROR CLEANING KIT. \$2.35 ppd.



SEND FOR OUR NEW 68-PAGE CATALOG showing many of the other telescope items manufactured or distributed by COAST INSTRUMENT. It includes our treatise, "WHAT YOU SHOULD KNOW, LOOK FOR, AND DEMAND, BEFORE BUYING ANY TELESCOPE," an impartial, informative article that will give you information you've probably been looking for to help make up your mind as to just what telescope you should buy!

All prices, unless otherwise indicated, are at our warehouse in Long Beach, California, from where shipping charges will be added, and are subject to change without notice. Mechanical specifications are also subject to change. Nominal crating charge added for all telescopes and mounts. California residents: Add 4% sales tax to all prices.

TRECKERSCOPES ARE CUSTOM MADE IN AMERICA BY PROFESSIONAL OPTICIANS AND ENGINEERS. NOW APPOINTING DEALERSHIPS . . . INQUIRIES INVITED.

COAST INSTRUMENT, INC.
"IN OPTICS SINCE 1933"

4811 Long Beach Blvd., Long Beach 5, Calif.

Phone: GARfield 2-3411 or NEVada 6-7683

THREE CURRENT COMETS

FOR several years, Los Angeles amateur Alan McClure has used his portable 7-inch $f/7$ photographic refractor in a program of attempting to rediscover periodic comets on their predicted returns. On the night of September 4-5, 1960, he took this instrument to the summit of 8,828-foot Mt. Pinos, California, in an effort to recover Comet Borrelly, last observed in 1953.

It was a windy but exceptionally clear night, and Mr. McClure waited for the scheduled lunar eclipse to attain totality, giving a sky dark enough for photography. Reproduced below is the 12-minute exposure he obtained, showing the comet as a barely visible 15th-magnitude object. Another picture, exposed for 21 minutes, also revealed it, but was somewhat fogged from twilight.

The identification of the object as the sought-for Comet Borrelly was established by three facts: It was located close to the predicted spot in Leo; it was moving in



Comet Borrelly's very faint image lies between the broad gray marks. Alan McClure took this 12-minute exposure at 11:36 UT, September 5, 1960.

the right direction at approximately the expected rate; a later check revealed no galaxy in the location.

Discovered in 1905, Borrelly's comet has a period of 7.02 years. It passed through the perihelion point of its orbit on June 12th of this year. Hence it is fading steadily as it recedes from both sun and earth.

On the same night, just before he recorded Comet Borrelly, Mr. McClure also took advantage of the eclipse darkness to obtain the 20-minute exposure of Comet Finlay shown here. It is likewise a periodic comet, requiring 6.90 years to complete one orbital revolution. The first observation of its 1960 return was Mr. McClure's photograph of June 20th, but he could not find it on the plate until a later Lowell Observatory position by R. Burnham, Jr., told him where to look.

Comet Finlay was considerably brighter than the other. Elizabeth Roemer, at the Flagstaff, Arizona, station of the U. S. Naval Observatory, saw it in a 5-inch finder on September 26th, and estimated



Mr. McClure's September 5th photograph of Finlay's comet shows a coma about 2.7 minutes of arc in diameter.

the total magnitude as about 11½. In November it will be about magnitude 13, and the following predicted positions (1950 co-ordinates) are taken from the *Handbook of the British Astronomical Association*:

November 2, 7^h 08^m.0, +25° 31'; 12, 7^h 06^m.3, +26° 35'; 22, 6^h 58^m.0, +27° 41'. December 2, 6^h 44^m.4, +28° 38'.

Periodic Comet Encke will return to perihelion on February 5th next year. It was first observed in 1660 on August 18th, by Dr. Roemer with the 40-inch reflector. Then a 19th-magnitude object, the comet was still extremely faint and nearly starlike when she recently obtained the image shown in the picture below.



Periodic Comet Encke is centered in this 30-minute exposure obtained by Elizabeth Roemer on September 26th. Official U. S. Navy photograph.

TRIGARTH TURRET and Eyepiece Attachment with Rack and Pinion

Just turn the Trigarth Turret and easily improve the performance of your telescope. It holds three eyepieces of standard 1¼" O.D. The Trigarth Turret sells for \$15.95 postpaid.



Also priced at \$15.95 postpaid, the Eyepiece Attachment with Rack and Pinion takes standard 1¼" O.D. eyepieces. The rack and pinion is machined from solid aluminum castings, precisely fitted for smooth performance. Main tube, 1¼" long; sliding tube adds 2"; total movement, 3¼". Choice of gray or black crinkle finish.

Both Turret and Eyepiece Attachment. \$31.90.

Satisfaction Guaranteed.

Write for Free Catalog. Instructions, 10¢.

GARTH OPTICAL CO.

P. O. Box 991, Springfield 1, Mass.

THE ASTRONOMER KIT

A complete telescope mirror making kit.

- ★ FOUCAULT TESTER, in its entirety, with light source, knife-edge, measuring scale
- PYREX-brand glass MIRROR BLANK
- ★ NEW, NONVITREOUS TOOL, faster, smoother, more accurate
- SEVEN ABRASIVES for greater safety and speed
- CERUM OXIDE for fast, clean polishing
- SPECIAL OPTICAL PITCH to minimize danger of turned-down edge
- MAGNIFYING LENS, essential in mirror grinding
- ★ DETAILED, ILLUSTRATED INSTRUCTIONS, "Mirror Making in the Kitchen"

(Starred items exclusive with The Astronomer Kit)

6-inch kit \$15.00 ppd.

4¼-inch \$11.50 ppd.; 8-inch \$22.50 ppd.

Foucault tester only \$ 6.00 ppd.

Money-back guarantee. Write for catalogue.

ASTRONOMY, INC.

Suite 305, American National Bank Bldg.

Denver 2, Colorado

TELESCOPE MIRRORS

Better than 1/8 wave.

Guaranteed to resolve to Dawes' limit.

6-inch \$52.50

8-inch \$79.90

10-inch \$165.00

No C.O.D.'s, please. Pennsylvania residents add 4% sales tax.

H. HUNTER LARGE
Optics

995 North Franklin Street
Pottstown, Pennsylvania

DE LUXE

SKYSCOPE

As illustrated — with new plastic tube, 6x finder, heavy all-metal tripod, equatorial mounting, and 60x eyepiece.

Price **\$44.75**

Unconditionally guaranteed.

100% American made.

Write for free brochure showing our standard model which has been sold world-wide for over 20 years. Sky-scope still remains outstanding in value by any comparison.

THE SKYSCOPE CO., INC.

P. O. Box 555, Brooklyn 28, N. Y.



DEEP-SKY WONDER

NOT ALL of the Messier objects are grand and impressive. A rather sparse one is M34 (NGC 1039) in Perseus, at right ascension $2^h 38^m.8$, declination $+42^\circ 34'$ (1950 co-ordinates). It is



The open cluster M34, photographed with Palomar Observatory's 48-inch Schmidt camera. South is toward the top in this view, and the width of the picture is one degree. This object is 1,500 light-years away, and it contains 81 stars brighter than magnitude 12.7, according to A. Wallenquist.

easily found between Algol and Gamma Andromedae, a little nearer the former.

Unlike most clusters, M34 is not more spectacular in large telescopes, as it does not seem to have the needed fainter stars to buttress the view. Rather, I feel that

15 x 65 binoculars give the best impression, though this is from tests made in the hazy air of the Connecticut River valley. M34 looks much like the Palomar 48-inch Schmidt picture reproduced here — something not quite applicable to most of the clusters we have discussed this year. More magnification merely spreads out the few bright stars that the binoculars show perfectly well.

H. Shapley lists the cluster's diameter as $18'$, although the exact boundaries are not clear, and the count of 80 stars in the catalogues leads the observer to expect more than is to be seen. Nevertheless, the total magnitude of M34 is 6.

As this was my first night of observing in Connecticut, I naturally turned to the objects that I used in Kansas to test the seeing. The Andromeda galaxy M31 did not seem to suffer from poor seeing, for it could be traced out to over three degrees in the binoculars. M33 in Triangulum was out of reach of the naked eye, and in binoculars gave the illusion of hanging between me and the starry background, an effect I have seldom noticed on nebulae. The Double Cluster in Perseus looked washed out and lacked its usual sparkle.

WALTER SCOTT HOUSTON
36 Lawn Ave.
Middletown, Conn.

SKY-GAZERS EXCHANGE

Classified advertising costs 30 cents a word, including address; minimum charge, \$4.00 per ad. Only one for sale ad per issue for each advertiser. Remittance must accompany order. Insertion is guaranteed only on copy received by the 20th of the second month before publication; otherwise, insertion will be made in next issue. We cannot acknowledge classified ad orders. Sky Publishing Corporation assumes no responsibility for statements made in classified ads, nor for the quality of merchandise advertised. Write Ad Dept., *Sky and Telescope*, Harvard Observatory, Cambridge 38, Mass.

EQUATORIAL MOUNTS: Seven models finished or semifinished. Marvin Stell, 600 E. 5th, Coleman, Tex.

ALUMINUM TUBING: Sun diagonal for Japanese refractors, \$6.50. Peninsula Scientific, 2421 El Camino Real, Palo Alto, Calif.

FOR SALE: Like-new de luxe 4" Dynascope with drive, \$65.00. Charles E. Herfurth, 2828 E. 20th St., Tucson, Ariz.

FOR SALE: 208 color slides of astronomical, planet, and symbolic paintings. Catalogue and sample, 25¢. Morris Dollens, 4372 Coolidge Ave., Los Angeles 66, Calif.

BALL BEARINGS for superb mounts. 6"-inside-diameter, 48-ball; cost government \$32.00; each \$8.25 postpaid. R. E. Kyrilach, 3005 Graceland Dr. N.E., Albuquerque, N. M.

6" REFRACTOR, Tinsley Saturn, almost brand new. Cost \$600.00. Sell \$375.00. Alton Walker, Box 65, Pebble Beach, Calif.

164-PAGE photographic bargain catalogue, listing thousands of photographic bargains. Send 25¢ for your copy, credited on first order. Dept. 26-C11, Central Camera Co., 230 S. Wabash Ave., Chicago 4, Ill.

PYREX MIRRORS, 1/20 wave, 6" f/8, \$69.00. Other sizes on request. Astro-Research Laboratories, 4601 70th St., La Mesa, Calif.

6" REFLECTOR, equatorial, like new, electric drive, circles, \$140.00. James B. Thompson, 616 Du Maine St., New Orleans 16, La.

METEORITES: Visitors from space. Willamette, \$1.00 and \$2.00; Arizona crater, \$10.00, \$18.00, and \$20.00. Scientific Laboratory, 2846 Oakley Ave., Baltimore 15, Md.

AT LAST: Fiberglass mirror cells. Great strength, lightweight, modern design, blackened forever. Low thermal conductivity. Another one of our scientific advances. 6" cell, \$7.50; 8" cell, \$12.50. Also our superior fiberglass tubes. W. R. Parks, 20942 S. LaSalle, Torrance, Calif.

FOR SALE: Amateur-ground 8" paraboloidal mirror, f/8, good figure. Sacrifice for \$45.00. Bradley Stark, 101 Yale Station, New Haven, Conn.

WANTED: Questar in excellent condition. Give details and price. C. Bight, C.P.O. Box 435, Kingstons, N. Y.

4" UNITRON altazimuth refractor, year old, like new, \$350.00. H. G. Perry, 220 W. 98th St., New York 25, N. Y.

CASSEGRAINS, GREGORIANS, and other unusual optical systems made to order. Worlds, Box 366, Farmington, Conn.

MOUNTED first-quality 6" telescope objectives, \$350.00. Correspondence invited. 24 years of experience. Earl Witherspoon, Sumter, S. C.

87 ISSUES of *Sky and Telescope*, August, 1950, to December, 1957, \$30.00. Frank Simon, 5924 Southern Ave., S.E., Washington 20, D. C.

SOUTHWESTERN OBSERVERS: Finest refractor telescopes supplied in 2.4", 3", and 4" altazimuth and equatorial models. Details on request. Melton Industries, 1901 Levee St., Dallas 7, Tex. Phone: RI 8-4769.

INTERESTED in astronomy as a career? *Vocational and Professional Monographs: Astronomy*, by Dr. Freeman D. Miller, describes personal qualifications, scholastic training, and job opportunities. \$1.00 postpaid. Send to Box B, *Sky and Telescope*, 49 Bay State Rd., Cambridge 38, Mass.

FOR SALE: M6A1 wide-field elbow telescope, 8" 45' field, 2-1/16 eye-lens aperture, four filters, unused, \$27.50. Model C3 infrared receiver, filter, good condition, \$29.00. Free list. Thermoelectric Devices, 302 Massachusetts Ave., Cambridge 39, Mass.

UNITRON 4" photo-equatorial, all accessories, plus extras, tripod and new metal pier. Value approximately \$1,600.00. All for \$750.00. Joseph Lampel, 8141 Sycamore Rd., Indianapolis 44, Ind. Telephone: CL 5-1272 or ME 5-9531.

6" FECKER catadioptric Celestar. New condition. All original accessories. Prefer local sale or pick-up. \$425.00. Stephen Zuzze, 189-02 64th Ave., Fresh Meadows, New York 65, N. Y.

FOR SALE: Right-angle-prism telescope, 20x; color filters of clear, amber, red, neutral; 60-mm. lens; \$25.00. Rifle-range telescope, 50-mm. lens, 33x, \$20.00. Vario-power telescope, 15x to 40x, case, \$12.00. 20x 80-mm. prism telescope, case, tripod, new, \$25.00. Dr. F. N. Solsem, Spicer, Minn.

QUESTAR DE LUXE, practically unused. List price, \$995.00. Will accept \$825.00, including shipping cost anywhere in continental United States. Mirakel Optical Co., Inc., 14 W. First St., Mt. Vernon, N. Y.

SUNSPOT NUMBERS

The following American sunspot numbers for July have been derived by Martin Nason, National Bureau of Standards, Boulder, Colorado, from AAVSO Solar Division observations.

July 1, 144; 2, 152; 3, 153; 4, 158; 5, 141; 6, 117; 7, 119; 8, 117; 9, 115; 10, 90; 11, 72; 12, 73; 13, 87; 14, 91; 15, 108; 16, 117; 17, 118; 18, 125; 19, 127; 20, 126; 21, 127; 22, 123; 23, 106; 24, 97; 25, 90; 26, 82; 27, 77; 28, 75; 29, 82; 30, 64; 31, 63. Mean for July, 107.6.

The following American sunspot numbers for August have been derived by Dr. Sarah J. Hill, Whitin Observatory, Wellesley College, from AAVSO Solar Division observations.

August 1, 60; 2, 34; 3, 21; 4, 19; 5, 17; 6, 19; 7, 39; 8, 41; 9, 50; 10, 88; 11, 167; 12, 198; 13, 195; 14, 237; 15, 228; 16, 234; 17, 243; 18, 231; 19, 208; 20, 126; 21, 148; 22, 141; 23, 96; 24, 95; 25, 126; 26, 133; 27, 107; 28, 113; 29, 97; 30, 99; 31, 111. Mean for August, 120.0.

Below are provisional mean relative sunspot numbers for September by Dr. M. Waldmeier, director of Zurich Observatory, from observations there and at its stations in Locarno and Arosa.

September 1, 103; 2, 105; 3, 80; 4, 75; 5, 83; 6, 100; 7, 108; 8, 107; 9, 129; 10, 162; 11, 149; 12, 143; 13, 156; 14, 157; 15, 159; 16, 115; 17, 114; 18, 141; 19, 156; 20, 171; 21, 177; 22, 189; 23, 168; 24, 157; 25, 141; 26, 114; 27, 92; 28, 89; 29, 74; 30, 44. Mean for September, 125.3.

Astronomy Teaching Aids ★ ★ ★ ★ ★ ★ ★ ★




DeLuxe Celestial Globes

No. 740. Our finest 16" globe, designed as a visual aid by teaching astronomers. All stars of the first six magnitudes are shown on a pale blue background, with the first three magnitudes having bright yellow, circular backgrounds. Brighter stars are named, and constellation boundaries indicated. Declination and right-ascension circles are printed in black. Magellanic Clouds and Milky Way stand out in white, and important star clusters are indicated. Various bases available. Weighted table base illustrated. **\$67.50**

No. 742. 12" Celestial Globe. Shows stars down to 5th magnitude on a light blue background. Constellations named. On weighted table base with movable meridian and horizon. **\$35.25**


No. 745. 30" Illuminated Rotating Celestial Globe. A translucent plastic globe illuminated by electric light. Shows stars down to 10th magnitude. Background includes brightly colored figures from celestial mythology. Base houses a motor to produce one complete rotation of the globe every 1½ minutes. Attractive modern base supports globe. **\$1,185.00**



Trippensee Planetarium

No. 708. This demonstration device shows the relative positions and motions of the Sun, Earth, Moon, and Venus. Simplifies teaching the rotation of the earth, the annual revolution around the Sun, day and night, change of seasons, and the cause of eclipses. This is an orrery-type device and not a projection planetarium. Constructed of brass with chain-drive gears. Arm length is 16½", overall height 16". Hand operated. **\$96.50**

Spitz Shadow Orrery



No. 712. An ingenious classroom device to demonstrate in a minute the intricate movements of the six major planets. One may view them as from outer space, all moving in the same direction around the Sun. Then, in an instant, the motion of the planets can be seen as from the moving Earth. The inferior motions of Mercury and Venus from one side of the Sun to the other can be clearly demonstrated, as can the forward and retrograde apparent motions of Mars, Jupiter, and Saturn in their exterior orbits. **\$295.00**



Transparent Globes

No. 736. A combination globe device consisting of an outside celestial globe showing stars and constellations and an inside brightly colored 6" terrestrial globe. Unique star pointer simulates Sun's motion. Demonstrates many Earth-sky-Sun relationships.

No. 736 — 12" Globe **\$49.50**

No. 736 — 20" Globe **\$115.00**



Practical Telescopes

No. 755. 4.5" Reflector. Designed for students and teachers, approved under NDEA for high schools. Optical system: 4.5" mirror, f/11, spherically corrected with surface accurate to 1/8 wave. Three eyepieces: Kellner 50x, Ramsden 100x, Ramsden 200x. Mounting: equatorial and adjustable for latitude. Tube: seamless aluminum, 45" long. Pedestal: heavy-wall steel tube with collapsible legs, free from wobble and torque. **\$98.50**

No. 757. 4.5" Electra. Same optical system as above, but with synchronous electric motor drive and slow-motion control in declination. **\$149.50**

★ ★ ★ ★ ★ ★ ★ ★

Science Associates, Inc.

Instruments/Weather — Astronomy/Teaching Aids

Write for complete catalog

194 Nassau Street

P. O. Box 216

Princeton, N. J.

★ ★ ★ ★ ★ ★ ★ ★



OUR LENSES ARE UNEQUALED

So say the many purchasers at universities, laboratories, and government institutions. Prices and quality defy comparison.


ASTRONOMICAL OBJECTIVES

... Air-Spaced ...

Specifically designed for those who demand an objective lens for genuine astronomical observation.

WHY BUY FROM US?

On the premise that the objective lens is the heart of the telescope, any prospective builder would certainly expect one that will perform to exacting specifications. Some may ask, "Why a refractor over a reflector telescope?" In the first place, a refractor is much easier to build and cheaper to maintain over the years. A reflector contains many small parts that are subject to wear or loosening, while the refractor is generally more compact, easy to store or carry, and needs little attention. When exploring the possibility of building an astronomical refractor telescope, the question is, "Which objective lens is best?" The answer is simple, and attested to by thousands of satisfied people in the know. "A Jaegers Astronomical Objective" is the only answer. Examine these pertinent facts:



Each lens is thoroughly tested and guaranteed to resolve to Dawes' limit. They are corrected for both the C and F lines (secondary chromatic aberration). The zonal spherical aberration and the chromatic variation of spherical aberration are negligible. The cells are machined to close tolerances so that they fit directly over or into our standard aluminum tubing, eliminating any mounting problems. Test a lens, or have any qualified person test it; we are certain that you will be satisfied. If not, take advantage of our money-back guarantee. We offer the lowest-priced, hand-corrected, precision, American-made astronomical objective, mounted in an aluminum cell. Our reputation for high-quality lenses has established us as the most reliable source in the industry.

HARD COATED ON 4 SURFACES

3 1/4" APERTURE f/15 — 48" F. L.	Cat. No. S1447	Mounted and Not Coated	\$28.00 ppd.
	Cat. No. S1471	Mounted and Coated	32.00 ppd.
4 1/8" APERTURE f/15 — 62" F. L.	Cat. No. S1472	Mounted and Not Coated	\$60.00 ppd.
	Cat. No. S1473	Mounted and Coated	69.00 ppd.
6" APERTURE f/10 — 60" F. L.	Cat. No. S1861	Unmounted and Coated	\$150.00 ppd.
	Cat. No. S1860	Mounted and Coated	175.00 ppd.
6" APERTURE f/15 — 90" F. L.	Cat. No. S1863	Unmounted and Coated	\$150.00 ppd.
	Cat. No. S1862	Mounted and Coated	175.00 ppd.

"BIG" ACHROMATIC TELESCOPE OBJECTIVES

We have the largest selection of diameters and focal lengths in the United States available for immediate delivery. Perfect magnesium-fluoride coated and cemented astronomical lenses made of finest crown and flint optical glass. Not mounted. Each lens is guaranteed to resolve to the theoretical limit.

● ALUMINUM TUBING and CELLS are in stock. ●

Cat. No.	Diam.	F.L.	ppd.	Cat. No.	Diam.	F.L.	ppd.	Cat. No.	Diam.	F.L.	ppd.
S1476	(2 1/8")	(10")	\$12.50	S1520	(2 1/8")	(50")	\$12.50	S-957	(3 1/4")	(40")	\$30.00
S1112	(2 1/8")	(11 3/4")	12.50	S-851	(3-1/16")	(15")	21.00	S1155	(4")	(34 1/2")	60.00
S1110	(2 1/8")	(13")	12.50	S1158	(3 3/8")	(19 1/2")	28.00	S1460	(4 1/4")	(36")	60.00
S-958	(2 1/8")	(15 1/2")	9.75	S-822	(3-3/16")	(24 1/2")	22.50	S1159*	(4 3/8")	(42")	60.00
S1145	(2 1/8")	(20")	12.50	S1092	(3 3/4")	(26")	28.00	S1225	(4 3/8")	(42")	67.50
S-952	(2 1/8")	(23 1/2")	12.50	S1093	(3 3/4")	(28")	28.00	S1474*	(5-1/16")	(24 3/4")	75.00
S1431	(2 1/8")	(30")	12.50	S1139	(3 3/4")	(30")	28.00	S1475	(5-1/16")	(24 3/4")	85.00
S1432	(2 1/8")	(40")	12.50	S-955	(3 3/4")	(34 1/2")	28.00				

*Not coated.

Astronomical Mirror Grinding Kits



● FREE OFFER ●

Now you can save many tedious hours of grinding time. Tool and blank of No. S2053 and No. S2054 have a generated f/8 curve. Take advantage of this extra special offer at no additional cost.

NOTE: For those who desire to grind a curve to their own specifications, the standard blanks are also offered as manufactured, without the generated curve.

EACH KIT CONTAINS: Mirror blank, plate-glass tool, eyepiece lenses, first-surface diagonal mirror, assorted abrasives, tempered pitch, and inspection magnifying lens.

Cat. No.	Mirror Diam.	Thickness	ppd.
S2053	4 1/4" (f/8 curve)	3/4"	\$ 7.50
S2093	4 1/4" (flat)	3/4"	7.50
S2054	6" (f/8 curve)	1"	11.95
S2094	6" (flat)	1"	11.95
S2055	8" (flat)	1 3/8"	19.50
S2056	10" (flat)	1 3/4"	30.75
S2057	12" (flat)	2 1/8"	54.75

We pay POSTAGE IN U. S. — C.O.D.'s you pay postage. Satisfaction guaranteed or money refunded in 30 days.

Equatorial Mount & Tripod



FOR TELESCOPES UP TO 4"

Constructed of cast aluminum and finished in black. The tripod is war-surplus, one of the sturdiest of its size ever built. Made of prime oak with bronze fittings, 3 adjustable locking legs with steel shods. Extended 50", closed 36". Shipping weight is 13 lbs.

Cat. No. S2128 EQUATORIAL MOUNT \$17.50 ppd.
Cat. No. S2131 TRIPOD 15.00 ppd.

SAVE MONEY!

Cat. No. S2083 Combination EQUATORIAL MOUNT and TRIPOD f.o.b. Lynbrook, N. Y., \$27.50

Reflector Telescope Kits

Excellent quality mirrors, polished, aluminized, silicon-monoxide coated. Each kit contains mirror, diagonal, and lenses for eyepiece. No metal parts. Mounting instructions included.

Cat. No.	Diam.	F.L.	ppd.
S2284	3-3/16"	42"	\$ 9.95
S2285	4 1/4"	45"	14.75
S2286	6"	60"	26.25

Astronomical Mirrors

These mirrors are of the highest quality. Aluminized with silicon-monoxide protective coating. You will be more than pleased with their performance.

Cat. No.	Diam.	F.L.	ppd.
S1429	3-3/16"	42"	\$ 8.70
S1506	4 1/4"	45"	13.50
S1507	6"	60"	25.00
S2188	6" parabolic	48"	49.50

Mirror Mounts

Cast aluminum, holds all of our mirrors with metal clips. Completely adjustable, assembled.



Cat. No. S1634 3-3/16" Mount for 4 1/2" Tubing \$4.00 ppd.
Cat. No. S1633 4 1/4" Mount for 5" Tubing 4.00 ppd.
Cat. No. S1632 6" Mount for 7" Tubing 6.50 ppd.

A. JAEGER'S
6915 MERRICK RD., LYNBROOK, N. Y.

OPTICS

COATED BINOCULARS

Beautifully styled imported binoculars. Precision made.
LOW, LOW PRICES!



American Type



"Zeiss" Type

Complete with carrying case and straps. American type offers a superior one-piece frame and clean design. Individual Focus (IF) or popular Center Focus (CF).

Cat. No.	Size	Field at 1,000 yards	Type	Price, ppd.
S1149	6 x 15 IF	360 ft.	Opera	\$12.75
S1436	6 x 30 CF	395	"Zeiss"	18.75
S1435	6 x 30 IF	395	"Zeiss"	16.75
S1438	7 x 35 CF	341	"Zeiss"	20.75
S1437	7 x 35 IF	341	"Zeiss"	17.95
S1771	7 x 35 CF	341	American	23.50
S1439	7 x 35 CF	578	American*	35.00
S2191	7 x 50 CF	530	American**	42.50
S1106	7 x 50 CF	372	"Zeiss"	24.95
S-961	7 x 50 IF	372	"Zeiss"	22.50
S1503	7 x 50 CF	372	American	32.50
S1443	8 x 30 CF	393	"Zeiss"	21.00
S1229	8 x 30 IF	393	"Zeiss"	18.25
S1108	10 x 50 IF	275	"Zeiss"	26.75
S1442	20 x 50 CF	183	"Zeiss"	33.75

*Wide-angle 11°

**Wide-angle

All prices above plus 10% Federal tax.




Monoculars

Brand new, coated optics, complete with pigskin case and neck straps.

Cat. No. S1576	6 x 30	\$10.00 ppd.
Cat. No. S1577	8 x 30	11.25 ppd.
Cat. No. S1578	7 x 35	12.50 ppd.
Cat. No. S1303	7 x 50	14.75 ppd.
Cat. No. S1579	16 x 50	17.50 ppd.
Cat. No. S1580	20 x 50	20.00 ppd.

4 1/4" Reflector Telescope

35x — 175x
4x Finder — Wood Tripod



From the mysterious craters of the moon to the fabulous rings of Saturn, this high-quality telescope will give you many pleasurable hours of crystal-clear astronomical viewing. Contains an f/6 parabolic mirror which gives a 50-per-cent brighter image than the ordinary f/10. Two coated eyepieces, 35x and 175x; rack-and-pinion focusing; sun filter; and 4x focusing finder. Its short tube, 27 3/4" long, and collapsible wood tripod (33" closed, 60" open), make for portability, stability, and compactness. Finished in white enamel with black trim. Although priced low, this is definitely not a toy. Imported. Shipping weight about 20 lbs.

FREE BOOK with Scope: DISCOVER THE STARS.

Cat. No. S2273 f.o.b. Lynbrook, N. Y., \$59.50

8-Power Elbow Telescope

This telescope has a brilliant-image 48° apparent field; actually 325 feet at 1,000 yards. Can be adjusted for focusing 15 feet to infinity. 2" achromatic objective, focusing eyepiece 28-mm. f.l., Amici erecting system. "Turret-mounted" filters: clear, red, amber, and neutral. Lamp housing to illuminate reticle for night use. Truly the biggest bargain you've ever offered. Original Gov't. cost \$200.



8 x 50

Cat. No. S1686	Not Coated	\$13.50 ppd.
Cat. No. S1975	Coated	\$17.50 ppd.

A Beginner's Guide to the Skies


R. Newton Mayall and Margaret W. Mayall

"Probably the best handbook now available for a newcomer to amateur astronomy." — *Sky and Telescope*. Contains a wealth of information about the solar system, and many lists of objects to be observed with binoculars or a telescope; includes 100 pages of detailed charts and observing data, arranged alphabetically by constellation. 184 pages. Cloth bound.

Cat. No. S2301 \$2.50 ppd.

Star Diagonals

Guaranteed to make viewing the heavens with a refractor telescope 100% easier for the observer. Stop needless crouching forever. Choose the STAR DIAGONAL that best suits your requirements. Slides into standard 1 1/4" eyepiece mounts. Chromium tubes and aluminum housing, attractive black-crackle finish.



AMICI STAR DIAGONAL WITH ERECT IMAGE

For comfortable viewing of terrestrial or astronomical objects. The Amici prism gives an erect image.

Cat. No. S2095 \$12.00 ppd.

RIGHT-ANGLE PRISM STAR DIAGONAL

Contains a high-quality aluminized right-angle prism, wide field. Practical for astronomical use.

Cat. No. S2096 \$12.00 ppd.

MIRROR STAR DIAGONAL

Diagonal houses an excellent first-surface mirror, flat to within 1/4 wave. Important accessory for refractor telescopes.

Cat. No. S2097 \$12.00 ppd.

7 x 30 Wide-Field Finderscope



Newly designed wide-field finderscope for reflector or refractor. Featured in this handsomely conceived black anodized aluminum tube are a 30-mm. coated achromatic objective, a wide-field Kellner eyepiece, and the all-important true crossline reticle for fast, exact locating. Compact and lightweight, 9" long, 1 1/2" diam., 9 oz. May also be used as a hand-held telescope, offering excellent extra-wide-field views of the sky.

Cat. No. S2193 \$9.75 ppd.

Finder Ring Mounts




Will accommodate finder above. Made of cast aluminum with black crackle finish. Each ring has three locking wing screws for adjusting. Base has two holes for mounting screws, fits any diameter tubing. Easy to mount. Inside diameter of ring mount 1-15/16".

Cat. No. S1963 Per Pair \$3.95 ppd.

New Prismatic Telescope

Coated

80-mm. Objective



Big 80-mm.-diam. (3 1/8") objective, for spotting or astronomical use. Will show stars of the 11th magnitude — 100 times fainter than the faintest visible to the naked eye. Table-top tripod included, extended 18" closed 13 3/4". Length of scope 16 1/4". Combined weight of scope and tripod is 5 1/4 lbs. All coated optics.

FIVE EYEPIECES

Power	Field at 1,000 yards	Exit pupil diam.	Relative Brightness
15x	122 ft.	5.4 mm.	29
20	122	4.0	16
30	61	2.7	7
40	49	2.0	4
60	32	1.3	1

Cat. No. S2052 \$59.50 ppd.

60-mm.-diam. Scope. Same as above but with smaller objective. Equipped with same five eyepieces — 15x, 20x, 30x, 40x, 60x. Coated. With tripod.

Cat. No. S1721 \$42.95 ppd.

Mounted Eyepieces

The buy of a lifetime at a great saving. First choice of the amateur and professional astronomer. Perfect lenses set in black anodized standard 1 1/4" O.D. aluminum mounts.



Cat. No.	F.L.	Type	ppd.
S2192	.6 mm. (1/4")	Orthoscopic	\$13.25
S1851	.6 mm. (1/4")	Ramsden	4.75
S1853	.125 mm. (1/2")	Ramsden	4.50
S1207	.125 mm. (1/2")	Symmetrical	6.00
S1251	.16 mm. (3/8")	Erfle (wide-angle)	12.50
S1257	.16 mm. (3/8")	Orthoscopic	12.50
S1209	.18 mm. (3/4")	Symmetrical	6.00
S1211	.22 mm. (27/32")	Kellner	6.00
S1835	.27 mm. (1-1/16")	Kellner	4.50
S1355	.32 mm. (1 1/4")	Plossl	12.50
S1253	.35 mm. (1 3/8")	Symmetrical	8.00
S1255	.55 mm. (2-3/16")	Kellner	6.00
S1485	.56 mm. (2 1/4")	Symmetrical	6.00

COATED LENSES 75 cents extra.

"Giant" Wide-Angle Eyepiece

Coated Lenses

Erfle eyepiece best for low-power performance. Superior six-element design. 65° field, 1 1/2" E.F.L., clear aperture 2 1/4", focusing mount and diopter scale. Combine this eyepiece with our 5" objective No. S1475 for a tremendous light-gathering Rich-Field Telescope.



Cat. No. S1405 (illustrated) \$12.50 ppd.

Cat. No. S1858 Same as above without diopter scale \$ 9.95 ppd.

FREE RUBBER EYESHIELD

With S1405

Cat. No. S1594 1 1/4"-diam. ADAPTER for eyepiece above \$ 3.95 ppd.

ADAPTER PLATE AND TUBE. Aluminum tapered tube and adapter plate, black anodized and finished in black crackle. Slides into our 5" O.D. Aluminum Tubing No. S1511. These parts are used with the eyepieces above to make a Rich-Field Telescope. Details in our catalogue.

Cat. No. S2300 \$18.00

Wide-Angle Erfle




Brand-new eyepiece with 68° field; coated. E.F.L. 1 1/4". Focusing mount, 3 perfect achromats, 1-13/16" aperture.

Cat. No. S1020 \$13.50 ppd.

Cat. No. S1593 1 1/4" diam. ADAPTER for eyepiece above 3.95 ppd.

An Economical Eyepiece

This mounted eyepiece has two magnesium-fluoride-coated achromatic lenses 29 mm. in diameter. Excellent definition. E.F.L. 1 1/4". Cell fits 1 1/4" tubing.



Cat. No. S1911 Coated \$5.90 ppd.

Cat. No. S1991 Not Coated \$2.25 ppd.

Rack-&-Pinion Eyepiece Mounts



For Refractors

Here is a wonderful opportunity for you to own a most mechanically perfect Rack-&-Pinion Focusing Eyepiece Mount with variable tension and adjustment. Will accommodate a standard 1 1/4" eyepiece, positive or negative. The body casting is made of lightweight aluminum with black-crackle paint finish, focusing tube of chrome-plated brass. Focusing tube for reflectors has a travel of 4", for reflectors 2", and will fit all size tubing.

REFRACTOR TYPE

Cat. No. S1494 for 2 1/8" I.D. Tubing \$12.95 ppd.

Cat. No. S1495 for 3 1/4" I.D. Tubing \$12.95 ppd.

Cat. No. S1496 for 4 3/8" I.D. Tubing \$12.95 ppd.

REFLECTOR TYPE

Cat. No. S1976 (less diagonal holder) \$8.50 ppd.

Cat. No. S1982 DIAGONAL HOLDER \$1.00 ppd.

FREE CATALOG

Millions of lenses, prisms, eyepieces, rack-and-pinion eyepiece mounts, cells, aluminum tubing, mirrors, binoculars, telescopes, parts, accessories, and so forth. Write today!

A. JAEGER'S
6915 MERRICK RD., LYNBROOK, N. Y.

Star Atlases and Books on Astronomy

1961 AMERICAN EPHEMERIS, limited supply, order now.....	\$4.00
New: PHOTOGRAPHIC LUNAR ATLAS, edited by G. Kuiper.....	\$30.00
New: PROPERTIES OF DOUBLE STARS, by L. Binnendijk.....	\$12.50
THE OTHER SIDE OF THE MOON.....	\$2.50
MOON MAPS, by H. P. Wilkins.....	\$6.00
AMATEUR ASTRONOMER'S HANDBOOK, by J. B. Sidgwick.....	\$12.75
OBSERVATIONAL ASTRONOMY FOR AMATEURS, by J. B. Sidgwick.....	\$10.75
SKY OBSERVER'S GUIDE.....	\$2.95
THE PLANET VENUS, by P. Moore.....	\$3.95
THE PLANET JUPITER, by B. Peek.....	\$8.95
AMATEUR TELESCOPE MAKING, Book 1, \$5.00; Book 2, \$6.00; Book 3, \$7.00	
Norton's STAR ATLAS.....	\$5.25
Beyer-Graff STAR ATLAS.....	\$15.00
BONNER DURCHMUSTERUNG.....	\$100.00

Pocketbooks and paperbacks on astronomy — write for listing. Books on telescope making and optical glassworking.

HERBERT A. LUFT

69-11 229th St., Oakland Gardens 64, N. Y.

ATOMIC CHART

of the ELEMENTS

A new approach to the periodic table, from the standpoint of atomic structure.

Arranged in circular form for easy reference, the chart includes atomic number, symbol, name, and mass of all known elements. 17" x 24", printed on heavy paper in two colors.

\$1.00 postpaid

THE WILLIAM RICHMOND PETERS OBSERVATORY AND RESEARCH GROUP

115 East Aspen St., Flagstaff, Ariz.



BOOKS AND THE SKY



ARABISCHE STERNNAMEN IN EUROPA

Paul Kunitzsch. Verlag Otto Harrassowitz, Taunusstr. 5, Wiesbaden, West Germany, 1959. 240 pages. DM 28, paper bound.

THIS great work in German covers such a large field and is so important to the student of star names that a reviewer finds it extremely difficult to determine whether or not he has adequately evaluated the book. Nevertheless, I have had considerable pleasure in reviewing it.

To the title, *Arabic Star Names in Europe*, Paul Kunitzsch might very well have added "and the many corrupt forms thereof." Such forms outnumber the legitimate names many times over. The work is undoubtedly the finest on Arabic star names that has ever been published.

One hundred and fifty years ago the student of stellar nomenclature looked to C. L. Ideler for many of the Arabic names found in the works of T. Hyde and al-Qazwini. But Hyde did not have any authentic manuscripts of as-Sufi; and one could not be sure in some instances what stars were being designated by at-Tizini because of errors in right ascension and declination.

Dr. Kunitzsch's title states that he is concerned with the Arabic star names in Europe. He is, however, too modest, for

I believe that he has not restricted himself to any one continent or any one hemisphere. The collection of sources is so thorough that the work could not have been written without the benefit of the great libraries of Europe, Africa, and Asia. And one author who resides in the United States has been mentioned or quoted several times.

The chapter on astrolabes is most excellent and fills a long-felt need. Although we have heard about them in the literature from time to time, there was always much to be desired, and we were never certain that the names were copied correctly.

The history and description of 210 star names, corrupt and otherwise, comprise the most important section of the book. Here, a tremendous amount of information about the Arabic names, including the non-Arabic one Albireo, is given. Unfortunately, in many cases the material is presented in a way that is sure to confuse the reader who is not an accomplished linguist, and who cannot differentiate between the original, most ancient authorities and those authors who may look as important and learned, but whose knowledge of Arabic is not competent. The qualified student, on the other hand, is bound to be most grateful for the many excellent references.

In one important respect Dr. Kunitzsch's work is very disappointing: Without any previous discussion of the possible meanings or interpretations of a given star name he condemns descriptions by other authors as "artificial" or "ridiculous." This procedure reminds one somewhat of the decision of the "Qualifiers" that Galileo's proposition of the immobility of the sun was "foolish and absurd." The reader feels left in a void; with his great knowledge of Arabic, the author should have indicated the reasons for his denunciations or at least given what he believes to be the true definitions and interpretations.

In another respect Dr. Kunitzsch leaves us with a very peculiar sense of the value and meaning of scholarly research. He gives the impression that, unless the exact meaning of a star name is found in an Arabic star text, it is practically sacrilegious for one to attempt to explain it from his own knowledge of Arabic and on the general principles of modern criticism. It is as though it were taboo for scholars such as Sethe, Breasted, or Speleers to write on the monuments of ancient Egypt, or for Kramer, Langdon, or Jastrow to give us the benefit of their great knowledge of Sumerian and Akkadian, or for West, Müller, or Darmesteter to attempt to translate or interpret the difficult passages in Avestan, Pahlavi, and Persian writing.

Dr. Kunitzsch forgets that in connection with all ancient languages covering

YOUR TELESCOPE IS NO BETTER THAN YOUR ABILITY TO FIND THE THINGS YOU WANT TO SEE

Now — for the first time



An entirely practical solution to the amateur's biggest problem. Now you can find celestial objects with an ease, speed, and accuracy never before approached.

To find any object you must know the sidereal time, and ACCUSTAR will give it to you instantly. Having already set the outer disk to your longitude, turn the inner disk until your clock (standard) time mark coincides with the date. The sidereal time is then indicated on the inner dial. It's as simple as that. Consistently accurate within two minutes or less.

ACCUSTAR will do many things: tell precisely where any star or constellation will be at any time or place, determine longitude, tell the solar time with one simple observation easily made with ACCUSTAR.

ACCUSTAR and the book, *Find the Stars*

\$9.75

A precision instrument, blue and gold in color, 10 by 12 inches, plastic coated, metal backed, weighs less than a pound. Nothing to get out of order. The 56-page book *Find the Stars* has 23 sky maps and needed information on over 250 celestial objects of prime interest to the amateur.

ACCUSTAR will be your indispensable guide to the heavens and multiply your pleasure in exploring them. Place your order now.

ACCUSTAR

P. O. Box 2806, Detroit 31, Michigan

long periods of time, certain significant bits of knowledge become lost. Even the meaning of the Sphinx had been forgotten or misunderstood by the time of the Empire or New Kingdom of Egypt. The author, I must admit, has a right to be skeptical at times concerning the knowledge of some moderns who feel the urge to write on star names; only a few years ago a well-known observatory published under its own aegis a list of Arabic star names with their meanings. In this list one of the names of Alpha Hydrae was translated as "The Brave Man's Neck." Such stellar erudition is hardly to be appreciated.

The beautiful modern Oriental system of transliteration has been used throughout, and the author and his publisher are to be congratulated on this score, as I have not yet found any typographical errors. It is unfortunate, however, that a book so important does not have a more durable and attractive binding.

The student of star names has waited for this book for many years; and from now on there will be no excuse, including even ignorance of Arabic, for writers on the constellations to make the same mistakes generation after generation.

GEORGE A. DAVIS, JR.
Buffalo, N. Y.

THE SEARCH FOR ORDER

Cecil J. Schneer. Harper and Bros., New York, 1960. 398 pages. \$6.00.

THE AUTHOR of this book, a comparatively young geologist-crystallographer, must be considered a rather remarkable scholar. In an age when specialization makes conversational strangers even of fellow faculty members in one department of a college, he can write lucidly and entertainingly about every field of science from botany through relativity. Dr. Schneer is a well-nigh perfect example of what he would like to have every serious science student become before he undertakes the ever-narrowing journey to proficiency in his chosen specialty.

As stated in the preface, "there can be a sense of excitement and compulsion to science," and the author has pretty well conveyed this to the reader. But the apology for the historical order of his development is not as well expressed as the one of Dr. Henry Margenau in the book's foreword: "It is commonly supposed that discussion of method makes the teaching of science more difficult. I doubt if this is true, but if it is, a compensating emphasis will make it easy. This is the accent on *history* of science . . . for it exposes the existential bond which holds scientific thought together. Historical incidents furthermore enliven the teaching of science in an anecdotal way, and liveliness is surely needed in our elementary science courses."

Dr. Schneer's classes, the reviewer be-


lieves, must be quite lively with anecdotal material, much of which he has included in the book. Sometimes this leads to a slight looseness in expression of the scientific principles involved, but that is usually well taken care of later on. In a way, the author does not let either the personal anecdote or the critical scientific point stand in the way of his objective — to point up the search for order. He confesses what so many of us fail to realize, that we cannot define the scientific method in so many words or even in so many volumes, but we can, by a critical and thorough review of the march of science, have a certain sure feeling for it.

The young scientist is likely to find the chapter on "Classical and Nonclassical Science" very revealing and even, in spots, rather startling. The scientist who is three decades or more beyond his own student days will similarly be disturbed by having science and truth, science and history, and science and so many other catchwords of our civilization, so keenly discussed in a way that many teachers should envy.

Dr. Schneer often plays the role of the die-hard antagonist very well, never scoffing at an outmoded way of thinking except when it has rested upon poor foundations. He has been very gentle with the obstructionists, defending them for their sincerity and thus attempting to pass along to us a lesson that all of

ELEMENTARY Astronomy

by
OTTO STRUVE,
Director,
National
Radio
Astronomy
Observatory,
with
BEVERLY
LYNDS
and
HELEN
PILLANS



AUTHORITATIVE AND UP TO DATE, this absorbing account of modern astronomy stresses the basic ideas and laws of physics as they apply throughout the universe. Otto Struve, an eminent authority on astronomy and regular contributor to *Sky and Telescope*, considers at length such topics as universal gravitation; the origin and evolution of the stars, the sun, the planets, and the comets; nuclear processes in stars; the dynamics of the galaxy. Ideal as a manual for the amateur astronomer at home or in school, the book contains a superb collection of illustrations, including 110 photographs, 181 line drawings, a color spectrum, and star charts. 1959. 404 pages. \$7.00.

OXFORD UNIVERSITY PRESS NEW YORK 16

THE MONTHLY EVENING SKY MAP

Have you seen it yet? THE MONTHLY EVENING SKY MAP . . . the exciting new version of the magazine that has "brought the stars down to earth" for many thousands of readers for more than half a century. Now . . . in its bright and expanded format, the SKY MAP is bigger and better than ever before.

FREE TRIAL!!

Receive Nov.-Dec. issue FREE by sending order now . . . your subscription will begin with Jan.-Feb., 1961, issue. Money refunded if not satisfied with first issue!

One year \$3.00 Two years \$5.00

- ★ Articles of popular interest.
- ★ Guide for users of small telescopes.
- ★ Detailed star and planet charts for each month.
- ★ Regular features by such writers as Tom Cave, Walter Houston, Leon Campbell, Jr.
- ★ Group projects for students and amateurs.

Sample copy 50 cents (Bimonthly)

A wonderful Christmas gift . . . just add \$2.00 to your subscription for each additional gift. Recipient receives card and bonus issue!

THE MONTHLY EVENING SKY MAP

Sky Map Publications, Inc.
P. O. Box 213-U, St. Louis 5, Mo.

With this fascinating, precision-built globe combination you can follow the exciting events of an entire day in the heavens. You read time by observing the earth's relation to the sun. There are two globes, a 6-inch terrestrial and a 12-inch celestial. The earth globe rotates by an electric motor once every 24 hours. The sun's noontime position is shown for any point on the earth's surface, and because the movable ecliptic ring is connected to the twilight ring, sunrise and sunset and the duration of twilight are shown for all places.

Only two daily settings are necessary — simply move the sun indicator eastward to the next day's date on the ecliptic circle, and turn the earth globe one degree to correspond to the actual rotation of the earth (about 361° in 24 hours).


The movable ecliptic (sun) ring is also marked for the moon's phase positions and the maximum greatest elongations of Mercury and Venus.

Model ST-12-E.C. — 12-inch diameter \$97.50

Write for free brochure.

FARQUHAR TRANSPARENT GLOBES
3724 Irving St., Philadelphia 4, Pa.
Telephone: EVergreen 2-2833

FARQUHAR EARTH CLOCK



us may, at some time or other, find hard to learn.

There are remarkably few outright errors in the book: Pluto's date of discovery is given as 1934 on page 123, the sun's distance from the center of the galaxy as 50,000 light-years on page 162; Galileo heard reports of sunspots on page 59 and discovered companions to Saturn (with no further explanation) on page 60; and there are a small handful of other mistakes. But they certainly do not detract from the fine quality of the writing and the clearness of expression in the book taken as a whole. The choice of illustrations is good, and the mechanical aspects of the volume are conducive to reading large sections at one sitting.

The reasoning is rather rigid in many passages, requiring considerable concentration and even, on occasion, a fast reprise of an earlier section for full comprehension, but the end result is rewarding. The reviewer wonders how well the average student of the liberal arts could follow some of the exposition with the one casual reading he would presumably want to give it.

Certainly this is not to be considered a "popular" book in the usual sense, but few really good works are popular. For the serious student of science, this volume should be required reading. For the serious citizen who wishes to find out what is happening to his comfortable early 20th-

century world, this book contains many of the answers. At the conclusion of almost every chapter, Dr. Schneer's paragraphs, not of summary but of impression and impact, in themselves could stand among the most quotable expressions of social consciousness of our times.

A beautiful blend of science, philosophy, and humanism has been achieved in this book. As a reference volume for a lively series of lectures to beginning students in liberal arts colleges, Dr. Schneer's method could project that "sense of excitement and compulsion" that would persuade more young people to embark on careers of science.

ROY K. MARSHALL
New York, N. Y.

PHYSICS AND MEDICINE OF THE ATMOSPHERE AND SPACE

Otis O. Benson, Jr., and Hubertus Strughold, editors. John Wiley and Sons, Inc., New York, 1960. 645 pages. \$12.50.

THE PROCEEDINGS of the second international symposium on physics and medicine of the atmosphere and space, held in San Antonio, Texas, in November, 1958, are reported in this volume, making it a must for everyone concerned professionally with the problems of space. Furthermore, it provides the nonspecialist with authoritative and comprehensible information.

The editors are pioneers in space medicine who arranged the first symposium on the topic in 1952. The balanced format and wide scope of this book are derived from their experience.

At the May, 1958, IGY symposium on the use of satellites for biological research, there were difficulties in the semantics of communication between physicists and biologists. However, a remarkable improvement in mutual comprehension is demonstrated in the present volume, which contains 44 papers presented by internationally known scientists.

Eight of these reports deal with the physical composition of the upper atmosphere and the contents of interplanetary space. H. K. Kallman of the Rand Corp. reports that rocket and satellite data disclose higher densities in the upper atmosphere than had previously been assumed. Large day and nighttime variations in atmospheric density occur, as well as changes with latitude. F. L. Whipple assesses the amount of meteoric material in space, and the hazards of meteoric impact in space vehicles and operations.

J. A. Van Allen discusses how the radiation belts that bear his name may influence spaceflight. The theory that the earth's magnetic field is trapping and storing solar corpuscular radiation is explored by S. F. Singer in his discussion of interplanetary dust and radiation.

M. Nicolet, University of Brussels, gives



**SATELLITE
3-INCH**

\$29.95

Completely assembled 3-inch reflector * All-metal construction with seamless aluminum tube * Standard 1 1/4-inch rack-and-pinion focusing eyepiece mount * Corrected optics for fine definition * Collapsible self-locking metal tripod * Peep-sight finder * 60-power eyepiece * (120-power eyepiece available, \$4.00) * With altazimuth mounting, \$29.95 * With equatorial mounting, \$39.95 * f.o.b. Pottstown, Pa.

BOOKLET, "Instructions To Establish Your Telescope and To Set and Use Your Setting Circles." 21 pages. **\$1.00 ppd.**



**SPACE K-4
DELUXE**


**\$79.95
KIT**

**\$98.50
ASSEMBLED**

OUR SPACE K-4 DELUXE . . . nationally recognized by astronomers and schools for its outstanding performance. Purchased by schools under NDEA for teaching astro-sciences, the SPACE K-4 DELUXE is available completely assembled for \$98.50, or as a kit with assembled optics for \$79.95, f.o.b. Pottstown, Pa. Prices include 3 eyepieces: 50x, 100x, and 200x, standard 1 1/4"; 6 x 25 achromatic finder; adjustable equatorial mount; pedestal; and collapsible legs. This model also available with clock drive and setting circles.

All prices subject to change.
DEALER INQUIRIES INVITED

SPACEK INSTRUMENT CO.



© 1959

PLANETARIUM SHADE

Attractively styled lamp shade provides the finishing touch for anyone interested in astronomy and space science. Constructed of durable translucent stock and printed in color. Stars appear white with outline of 37 constellations on deep blue background. Constellations of spring, summer, fall, and winter accurately scaled and identified by name, with Milky Way, zodiac, and equator outlined. Size, 12" diameter, 8 1/2" high. Supplied with clip or washer-type mount (specify which when ordering). Ideal gift. Practical and educational. **\$5.98 ppd.**

FAVORITE CONSTELLATIONS OF THE NORTHERN HEMISPHERE

© 1959

A photographic atlas of the sky from declination -30° to the North Pole. This set of **33 slides** on 35-mm. film is made up of actual **unretouched photographs** requiring almost three years to obtain using special equipment. Set includes 39 slides — one title, five star maps, and 33 constellations. Each slide mounted and identified. Suitable for projection with any 35-mm. equipment on screen and ceiling for lectures and teaching. This is the finest set of slides we have ever seen; each constellation is sharp and clear with excellent contrast. Instructions included. **\$16.00 ppd.**

Inquiries from schools, clubs, dealers, and planetariums invited.

1130 Sembling Ave.
Pottstown, Pa.
Telephone: FAculity 3-2825



There's a Sky Publication for every one on your Christmas list

SKY SETS I. 24 beautiful halftone prints of objects in the solar system and in the Milky Way. 8½" by 11¾" on paper 12½" by 16¾". **\$4.00 per set**

SKY SETS II. 20 halftone prints of striking nebulae in the Milky Way and portraits of other galaxies, and four drawings of the 200-inch telescope. 8½" by 11¾" on paper 12½" by 16¾". **\$4.00 per set**


MOON SETS. 18 halftone prints forming a large-scale photographic map of the entire visible face of the moon, made from Lick Observatory negatives of the first and last quarters. **\$3.00 per set**

LUNAR CRESCENT SETS. 10 halftone pictures from Lick Observatory that form a matching series to Moon Sets, but for the waxing crescent 4½ days after new moon, and the waning crescent about five days before new moon. **\$2.50 per set**

ELGER'S MAP OF THE MOON. The detailed moon chart itself is 18" high by 17½" wide and identifies all important lunar features. Under it are extensive notes by H. P. Wilkins on 146 regions. **\$3.00**


PHILIPS' PLANISPHERE. Suitable for use within 10 degrees of latitude 42° north. Printed in dark blue and gold, it identifies the principal constellations and stars above the observer's horizon for any time of the night throughout the year. **\$3.00**

COLOR MAP OF THE NORTHERN HEAVENS. Shows the positions and apparent magnitudes of celestial objects; distances, sizes, and spectral types of many bright stars. The northern sky to -43° is plotted on a polar projection; stars brighter than magnitude 5.1 included. 30" by 34½". **\$1.00**



COLOR CHARTS OF THE MOON

Two striking color charts of the first- and last-quarter moon. Each lunar half is an impressive 27" in diameter; the over-all size of a chart, 23" by 33". Here are two genuinely useful and informative maps that will be welcomed as a decoration in any observatory, den, or clubroom. Printed in predominantly gold and black on heavy paper. **\$2.00 per set of 2**



LAROUSSE ENCYCLOPEDIA OF ASTRONOMY.

By L. Rudaux and G. de Vaucouleurs. "By far the best general survey of its kind for anyone who has graduated from the ordinary popularizations and wants a taste of the real business of astronomy. . . . The lucid and interesting text is supported by a most effective and intelligently planned collection of diagrams, photographs, and charts." — *Scientific American*. Many full-color renditions. 506 pages, 818 illustrations. **\$12.50**

NORTON'S STAR ATLAS.

Covers the whole heavens, showing over 9,000 objects: stars to magnitude 6½, nebulae, and clusters. Descriptive lists of 500 subjects for viewing with small telescopes, and data for observing the sun, moon, and planets. 108 pages. Here is one astronomical gift that will be cherished by any amateur. Latest, 14th edition. **\$5.25**

POPULAR STAR ATLAS.

Simpler version of Norton's *Star Atlas*, with 16 bound charts to magnitude 5½. 38 pages. **\$2.00**

THE HISTORY OF THE TELESCOPE.

By Henry C. King. A complete account of the development of the telescope from crude instruments of the ancients to the giants of today. 456 pages, 196 illustrations. **\$7.50**

DE LUXE ATLAS OF THE HEAVENS.

This is the *Atlas Coeli 1950.0*, by A. Becvar. 16 bound charts cover the entire sky to magnitude 7.75, and show over 35,000 objects. Printed in many colors, each map is 16½" by 23". Transparent coordinate grid overlay included. **\$9.75**

PHOTOGRAPHIC LUNAR ATLAS.

Edited by G. P. Kuiper. Contains 281 of the finest photographs of the moon in existence, taken with large telescopes at Mount Wilson, Lick, Pic du Midi, McDonald, and Yerkes observatories. 230 sheets, 15½" by 19", loose leaf in sturdy box, thumb-indexed and numbered. (Orders accepted from the United States and possessions only.)

\$30.00 each; 2 for \$50.00

FIELD ATLAS OF THE HEAVENS.

Each chart is reduced from the original *Atlas Coeli*; stars are white on black background. The 16 charts are unbound, each 18" by 12¼", on heavy paper. **\$4.00 each; 2 for \$7.50**

ATLAS COELI CATALOGUE.

Listed, with descriptive data, are the 6,362 stars brighter than magnitude 6.26, with their right ascensions and declinations, magnitudes, and much other useful information; extensive lists of star clusters, nebulae, and galaxies. Cloth bound, 367 pages. **\$8.75**

MAKING YOUR OWN TELESCOPE.

By Allyn J. Thompson. Complete step-by-step directions for making and mounting your own 6-inch reflector at low cost. In easy-to-understand chapters, you learn how to grind, polish, and figure the mirror, and how to make an equatorial mount that will provide a sturdy, solid support for your optics. 211 pages, 104 illustrations. **\$4.00**

SPLENDORS OF THE SKY.

Celestial picture booklet with short and easy-to-understand captions. 36 pages, 66 illustrations. 8½" by 11½". **50c**

Write for descriptive catalogue. All items sent postpaid. Please enclose your check or money order.

SKY PUBLISHING CORPORATION

Harvard Observatory • Cambridge 38, Massachusetts

an extended presentation of atmospheric chemical reactions. Oxygen and nitrogen, the principal gases of the atmosphere, are affected by solar radiation, which causes great variability of composition at higher levels.

Among the seven articles on technical phases of space travel is W. von Braun's discussion of timing in satellite launchings. Firing without careful prelaunching study might place a satellite in continuous sunlight, where its internal temperature could rise so high as to injure the apparatus and affect the success of the mission.

Sixteen papers presented next examine biological aspects of spaceflight. H. G. Clamann reports the physiological effects of ozone, especially in connection with high-flying aircraft and balloons. At 100,000 feet, concentrations of this gas reach four to six parts per million (p.p.m.).

Exposure to the inhalation of two p.p.m. of ozone for 16 hours a day depressed the growth rate of rats. Irritable behavior is manifested by human subjects

in a concentration of only one p.p.m. Exposure to six p.p.m. for an hour led to a marked decrease in the vital capacity of one subject, whose lung alveoli became partially filled with fluid.

Dr. Clamann also reviews some metabolic problems of a sealed cabin environment: oxygen requirements and processes of carbon dioxide elimination. Although rapid developments have taken place in this field since 1958, the article is still pertinent. J. Meyers, pioneer in algae physiology and related photosynthesis, presents the use of algae for gas exchange in a closed ecological system of the sealed space-cabin type.

Psychological problems of spaceflight indicated by laboratory experiments with sensory deprivation are well covered by G. T. Hauty. The foremost authorities on weightlessness, S. J. Gerathewohl and J. E. Ward, give an excellent account of its effects as simulated in parabolic flights. They consider spatial orientation, neuromuscular control, and other physiological functions.

Problems of visual perception, accommodation, and reaction time under space-flight conditions are described by H. Rose. Vision in the higher atmosphere and in space differs from that at sea level. Because of the gradual decline of light-scattering air particles, the sky becomes darker with increasing altitude and stars are visible by day. Outside the atmosphere light intensities are increased about 40 per cent.

Rescue operations from space vehicles are the topics of six papers, and the volume is concluded with a six-paper discussion of the physics and environment of the sun and planets.

This book seems to be the most important treatise on space biology thus far published in any language.

HOWARD LASKEY, M.D.
Carolina, R. I.

NEW BOOKS RECEIVED

THE UNIVERSE AROUND US, *Sir James Jeans*, 1960, Cambridge University Press. 297 pages. \$1.95, paper bound.

First published in 1929, this classic by an outstanding British astronomer has been reissued as a reprint of the fourth (1944) edition. The nature and origin of the stars and galaxies are lucidly described for the informed general reader. Much of the discussion of evolutionary problems, however, has been outmoded by recent work.

THE PLANETARY EQUATORIUM, *E. S. Kennedy*, translator, 1960, Princeton University Press. 267 pages. \$7.50.

The facsimile reproduction of a Persian astronomical manuscript, together with a line-for-line translation and an extended commentary, make up this volume. The text is largely devoted to a description of an astronomical computing device invented in the 15th century by Kashi, who worked at Ulugh Beg's observatory in Samarkand. The device was used to predict solar, lunar, and planetary positions, and the details of eclipses.

APPARENT PLACES OF FUNDAMENTAL STARS — 1961, 1960, *Astronomisches Rechen-Institut*, Mönchhofstr. 12-14, Heidelberg, West Germany. 511 pages.

The 21st of a series, this annual volume is prepared under the auspices of the International Astronomical Union. It lists apparent positions of 1,535 bright stars, giving right ascensions to 0.001 second and declinations to 0".01, at 10-day intervals.

In the United States, order from Cincinnati Observatory, Cincinnati 8, Ohio. Advance payment of \$6.50 per copy is required (covering postage and packaging); make checks payable to Cincinnati Observatory APFS. European orders should be sent to Verlag G. Braun, Karl Friedrich Str. 14, Karlsruhe, West Germany.

AN INTRODUCTORY TREATISE ON THE LUNAR THEORY, *Ernest W. Brown*, 1960, Dover. 292 pages. \$2.00, paper bound.

Although first published in 1896, this remains the standard English-language text on the moon's orbital motion. Written by a master of celestial mechanics, it traces the methods of Laplace, de Pontécoulant, Hansen, Delaunay, and the author. The highly technical treatment is intended for readers with an extensive mathematical background.

Read This Advertisement

Here is a combination of a Barlow and a particular ocular which gives outstanding results. It consists of our new Barlow and our 16.3-mm. (3/4" focal length) Erfle eyepiece. While the Barlow was not specifically designed to work with this eyepiece, it does so to an astonishing degree. All images are sharp and hard to the very edge of the field.

The Barlow gives magnification up to slightly over three times that of the ocular alone. It is achromatic, coated, and mounted to the U. S. standard size of 1.250 inches.

The modified Erfle eyepiece has a field of 75 degrees with excellent eye relief. The combination gives an equivalent focal length of slightly under 6 millimeters. Many users state it is far superior to any shorter focal length ocular of equivalent magnification.

The Barlow sells for \$16.00 postpaid, and the Erfle for \$15.30 postpaid. Both are guaranteed to perform as stated above or money refunded.

ORTHOSCOPIC OCULARS—All hard coated, standard 1 1/4-inch outside diameter.

28-mm.	\$15.60	10.5-mm.	\$16.80	4-mm.	\$17.70
16.3-mm. (Erfle)	\$15.30	7-mm.	\$17.70	Barlow 3x	\$16.00

Telescopes

6565 Romaine Street
Los Angeles 38, Calif.

Warranted to equal or surpass any oculars obtainable anywhere or money refunded.

Finished mirrors, mirror kits, spiders, elliptical flats, focusing devices, aluminizing.

Send for catalogue.

THE
NEW
SPITZ



MODEL C

PLANETARIUM

FOR THE SCIENCE MUSEUM and PUBLIC LIBRARY
of MINNEAPOLIS



PLANETARIUM DIVISION • SPITZ LABORATORIES, INC. • YORKLYN, DELAWARE

ENGINEERED FOR THE LAST WORD IN BREATHTAKING PERFORMANCE . . . PRICED FOR FIRST CHOICE IN VALUE . . . FULLY EQUIPPED WITH MANY EXTRA FEATURES!

You'll Marvel At How
The Superb Optics Of
This Portable RV-6

6-inch DYNASCOPE®

Gives The Same Exquisite Definition
As Far More Expensive Instruments!

This new addition to the Dynascope line has won immediate recognition from schools, colleges, and professionals, as an outstanding achievement in a 6-inch telescope. Although it was only recently introduced, our files are already filled with complimentary letters from excited amateurs and professionals all over the country. Each one is truly amazed at the superior optical performance of this new RV-6 6-inch Dynascope! Here is large aperture in a quality instrument at a price that compares with many 4-inch telescopes. *And this low cost includes such exclusive extra features as electric drive (patented), setting circles, and rotating tube!* There are no "extras" to run up your cost!

The superb optical system resolves difficult objects with definition that is absolutely breathtaking. The close tolerances of the precision construction assure an accuracy and smoothness of operation once associated only with the finest custom models. The heavy-duty mount, complete with electric drive, provides the stability so essential for satisfactory viewing, yet there is easy portability because in a matter of minutes the entire telescope can be dismantled into three easy-to-handle sections.

Only Criterion's engineering ingenuity, coupled with volume production and modern manufacturing methods, makes this handsome 6-inch model available at such reasonable cost. You can order it with complete confidence that it will live up to your expectations in every way, for this assurance is *guaranteed* under our full-refund warranty. Send your check or money order today. Or use our liberal time-payment plan and take months to pay.

YOU COULD PAY \$100 MORE WITHOUT GETTING ALL THESE SUPERIOR FEATURES (Except on Another Dynascope)

1. **EXQUISITE OPTICAL SYSTEM** INCLUDES F/8 6-INCH PARABOLIC MIRROR made of PYREX brand glass, accurate to better than $\frac{1}{8}$ wave, zircon-quartz coated, and guaranteed to reach all theoretical limits of resolution and definition. Teamed with elliptical diagonal, mounted in 4-vane adjustable spider.
2. **NEW DYN-O-MATIC ELECTRIC DRIVE** with smooth self-acting clutch that engages and disengages automatically as you seek different objects. Will not interfere when manual operation is desired. Entire drive housed in trim aluminum case. Plugs into ordinary household socket.
3. **CHOICE OF ANY 3 EYEPIECES** from the following: 65X-18-mm. Achromatic Ramsden; 100X-12.7-mm. Achromatic Ramsden; 150X-9-mm. Achromatic Ramsden; 220X-6-mm. Orthoscopic; 320X-4-mm. Orthoscopic. All are precision-threaded, give sharp images to extreme edges.
4. **SOLID NEW EQUATORIAL MOUNT**, extra-reinforced design to provide pillarlike stability. No annoying side play or wobble. Adjusts easily to any latitude.
5. **SETTING CIRCLES** for both right ascension and declination. Handsomely engraved and finished in fine aluminum.
6. **6 x 30 FINDERSCOPE**, ACHROMATIC, COATED, with accurate crosshairs and fine focus. Durable cast-aluminum bracket with 6 adjusting screws allows positive collimation.
7. **ROTATING TUBE FOR MAXIMUM VERSATILITY AND VIEWING COMFORT**. Solid-cast chrome-finished rings are generous 1" wide with felt lining. Newly designed construction, with over-sized knurled adjusting knobs, affords maximum rigidity and allows for quick disassembly and portability, with or without tube.
8. **STRIKINGLY HANDSOME WHITE 50" BAKELITE TUBE** with porcelainized Duralite finish, durable yet light. Walls are $\frac{1}{8}$ " thick, completely insulated and anti-reflective blackened inside.
9. **BRASS RACK-AND-PINION EYEPIECE HOLDER** has precision-cut rack and gear for easier, smoother, more positive focusing. Takes standard $1\frac{1}{4}$ " oculars, negative or positive.
10. **STRONG, VIBRATION-FREE, ALL-METAL TRIPOD** with easily removable legs. Provides sure, steady support, plus lightweight portability.

CRITERION MANUFACTURING CO.
331 Church St., Hartford 1, Conn.
Manufacturers of Quality Optical Instruments

Including . . . • **ELECTRIC DRIVE (Patented)**
• **SETTING CIRCLES** • **ROTATING TUBE**

A Complete Instrument, No Costly Accessories Needed!

Model RV-6 Complete
with Dyn-O-Matic Electric Drive
and All Features Described Below

\$194⁹⁵

f.o.b. Hartford, Conn.
Shipping Wt. 55 lbs.
Express Charges Collect
No Packing or Crating
Charges



ENJOY IT NOW FOR ONLY \$74.95 DOWN

No need to put off the thrills of owning this magnificent instrument! Send your check or money order today for only \$74.95 as full down payment . . . pay balance plus small carrying charge in your choice of 6, 12, or even 24 monthly payments. Same unconditional guarantee applies, of course. Or order today by sending your check or money order with coupon below.

Sound too good to be true?
Then read what these delighted
DYNASCOPE owners have to say:

"I never dreamed such a brilliant, clear image could be had with a telescope costing under \$200. It is one of these rare bargains that you find only once or twice in a lifetime."

B.S. — New Jersey

"A friend of mine has a 6-inch telescope without electric drive, for which he paid over \$500. He agrees that my new DYNASCOPE not only outperforms it, but is a much better-looking and better-built instrument."

J.L. — New York

"Congratulations on the excellence of your workmanship. The optics are truly amazing. I never expected such wonderful performance."

L.H.N. — Massachusetts

Criterion Manufacturing Co.
Dept. STR-16, 331 Church St., Hartford 1, Conn.

- ☐ Please send me, under your unconditional guarantee, the RV-6 6" Dynascope. Full payment of \$194.95 is enclosed.
- ☐ I prefer your easy terms! Enclosed is \$74.95 as down payment with understanding that I will pay balance (plus small carrying charge) over 6, 12, 24 months (check choice).
- ☐ Send FREE ILLUSTRATED LITERATURE describing the RV-6 6" Dynascope and all the telescopes in the Dynascope line.

Name
Address
City State



3-inch Astronomical Reflector

60 to 180 Power
An Unusual Buy!

Assembled — ready to use! See Saturn's rings, the planet Mars, huge craters on the moon, star clusters, moons of Jupiter, double stars, nebulae, and galaxies! Equatorial-type mounting with locks on both axes. Aluminized and over-coated 3"-diameter f/10 primary mirror, ventilated cell. Telescope comes equipped with a 60x eyepiece and a mounted Barlow lens, giving you 60 to 180 power. A finder telescope, always so essential, included. Sturdy, hardwood, portable tripod.

FREE with Scope: Valuable **STAR CHART** plus 272-page "**HANDBOOK OF THE HEAVENS**" plus the book "**HOW TO USE YOUR TELESCOPE**."

Stock #85,050-Y.....\$29.95 ppd.

4 1/4" ASTRONOMICAL TELESCOPE

UP TO 255 POWER

New Vibration-Free Metal Pedestal Mount



With this scope you can see everything described above, but with greater power. Also, it will split closer double stars. Mirror has twice the light-gathering power. Rack-and-pinion focusing, real equatorial mounting — only one motion needed to follow the stars! Aluminum tube, 6-power finder telescope, 2 mounted-size eyepieces and mounted Barlow lens give you powers of 40x, 90x, 120x, and 255x. Low-cost accessory eyepiece available for higher powers. FREE with Scope: Valuable **STAR CHART** plus 272-page "**HANDBOOK OF THE HEAVENS**" plus the book "**HOW TO USE YOUR TELESCOPE**." Shipping wt. 25 lbs.

Stock #85,105-Y.....\$79.50 f.o.b. Barrington, N. J.

Same telescope as described above but equipped with Electric Clock Drive.


Stock #85,107-Y.....\$109.00 f.o.b. Barrington, N. J.

NOW! A FULL SELECTION OF ORTHOSCOPIC EYEPIECES

Wide, flat field — better correction under high magnification — excellent eye relief.

The orthoscopic eyepiece is one of the most important and best corrected eyepieces for astronomical work. These are of a four-element design, with coated lenses, and are standard 1 1/4" outer diameter, precision made of chrome-plated brass and aluminum.

Stock #30,364-Y.....	4 mm.....	\$14.50 ppd.
Stock #30,404-Y.....	6 mm.....	14.50 ppd.
Stock #30,405-Y.....	12.5 mm.....	14.50 ppd.
Stock #30,406-Y.....	18 mm.....	14.50 ppd.
Stock #30,407-Y.....	25 mm.....	14.50 ppd.



Rack-&-Pinion Eyepiece Mounts



Real rack-and-pinion focusing with variable tension adjustment; tube accommodates standard 1 1/4" eyepieces and accessory equipment; lightweight aluminum body casting (not cast iron); focusing tube and rack of chrome-plated brass; body finished in black wrinkle paint. No. 50,077-Y is for reflecting telescopes, has focus travel of over 2", and is made to fit any diameter or type tubing by attaching through small holes in the base. Nos. 50,103-Y and 50,108-Y are for refractors and have focus travel of over 4". Will fit our 2 7/8" I.D. and our 3 7/8" I.D. aluminum tubes, respectively.

For Reflectors

Stock #50,077-Y...(less diagonal holder)...\$8.50 ppd.
Stock #60,049-Y...(diagonal holder only)...1.00 ppd.

For Refractors

Stock #50,103-Y...(for 2 7/8" I.D. tubing)...12.95 ppd.
Stock #50,108-Y...(for 3 7/8" I.D. tubing)...13.95 ppd.

SCIENCE TREASURE CHESTS

For Boys — Girls — Adults!

Excellent "Science Fair" Materials



Science Treasure Chest — Extra-powerful magnets, polarizing filters, compass, one-way-mirror film, prism, diffraction grating, and lots of other items for hundreds of thrilling experiments, plus a Ten-Lens Kit for making telescopes, microscopes, etc. Full instructions included.

Stock #70,342-Y.....\$5.00 ppd.

Science Treasure Chest De Luxe — Everything in chest above plus exciting additional items for more advanced experiments including crystal-growing kit, electric motor, molecular models set, first-surface mirrors, and lots more.

Stock #70,343-Y.....\$10.00 ppd.

BINOCULAR-TO-CAMERA HOLDER

For Exciting Telephoto Shots

Will fit any camera



Bring distant objects 7 times nearer with a 35-mm. camera, 7 x 50 binocular, and our **BINOCULAR-TO-CAMERA HOLDER**. Ideal for photographing the constellations, star clusters, the moon, as well as cloud formations, wildlife, vistas.

Camera and binocular attach easily. Use any binocular or monocular — any camera, still or movie. Take color or black-and-white. Attractive gray crinkle and bright chrome finish, 10" long. Full directions for making telephoto pictures included.

Stock #70,223-Y.....\$11.50 ppd.

Take Pictures Through

Your Telescope with the EDMUND CAMERA HOLDER FOR TELESCOPES



Bracket attaches permanently to your reflecting or refracting telescope. Removable rod with adjustable bracket holds your camera over scope's eyepiece and you're ready to take exciting pictures of the moon. You can also take terrestrial telephoto shots of distant objects. Opens up new fields of picture taking!



SUN PROJECTION SCREEN INCLUDED

White metal screen is easily attached to holder and placed behind eyepiece. Point scope at sun, move screen to focus — and you can see sunspots!

All for the low, low price of \$9.95

Includes brackets, 28 3/4" rod, projection screen, screws, and directions. Aluminum; brackets black crinkle painted.

Stock #70,162-Y.....\$9.95 ppd.

PRISM STAR DIAGONAL

For comfortable viewing of the stars near the zenith or high overhead with refracting telescopes using standard-size (1 1/4" O.D.) eyepieces, or you can make an adapter for substandard refractors. Contains an excellent high-quality aluminized right-angle prism. The tubes are satin chrome-plated brass. Body is black wrinkle cast aluminum. Optical path of the system is about 3 1/2".


Stock #70,077-Y.....\$12.00 ppd.



AMICI-PRISM STAR DIAGONAL

Same as above except contains Amici roof prism instead of usual right-angle prism. Thus your image is correct as to top-bottom, making it excellent for terrestrial viewing.

Stock #50,247-Y.....\$12.00 ppd.



NOW! LENS ERECTOR FOR TERRESTRIAL VIEWING WITH YOUR REFLECTOR

This Edmund development adds real convenience to viewing objects on the earth. Just put the lens erector in your eyepiece holder, insert eyepiece, and focus normally. You see everything right side up and correct as to left and right. Made of polished chrome-finish brass telescoping tubing that will fit any standard 1 1/4" eyepiece holder. Tubing is 9 1/2" long and slides 3" into eyepiece holder. Erecting system consists of two coated achromats.

Stock #50,276-Y.....\$9.95 ppd.

GIANT-SIZE FULL-COLOR ASTRONOMICAL POSTCARDS

Ideal for Mailing to Your Amateur Friends . . .

For Framing as Room Decorations

You'll be thrilled when you see these new astronomical postcards by Edmund. They consist of dramatic full-color renderings of Mars, Jupiter, Saturn, a sunspot, and a solar prominence, all done by astronomical artist Hugh Stevenson. Size, 5 1/2" x 7".

Stock #40,398-Y.....Set of 5.....50c ppd.

"MAKE-YOUR-OWN" 4 1/4" MIRROR KIT

The same fine mirror as used in our telescopes; polished and aluminized, lenses for eyepieces, and diagonal. No metal parts.

Stock #50,074-Y.....\$16.25 ppd.

OFFSPRING OF SCIENCE . . . REALLY BEAUTIFUL!



CIRCULAR DIFFRACTION- GRATING JEWELRY

A Dazzling Rainbow of Color!

Shimmering rainbows of gemlike color in jewelry of exquisite beauty — made with **CIRCULAR DIFFRACTION-GRATING REPLICA**.

Just as a prism breaks up light into its full range of individual colors, so does the diffraction grating. Promises to become a rage in current fashion.

Stock #30,349-Y...Earrings.....\$2.75 ppd.

Stock #30,350-Y...Cuff Links.....\$2.75 ppd.

Stock #30,372-Y...Pendant.....\$2.75 ppd.

Stock #30,390-Y...Tie Clasp.....\$2.75 ppd.

WAR-SURPLUS TELESCOPE EYEPIECE

Mounted Kellner eyepiece, type 3. Two achromats, focal length 28 mm., eye relief 22 mm. An extension added, O.D. 1 1/4", standard for most types of telescopes. Gov't. cost \$26.50.

Stock #5223-Y.....\$7.95 ppd.


6X FINDER TELESCOPE



Has crosshairs for exact locating. You focus by sliding objective mount in and out. Base fits any diameter tube — an important advantage. Has 2 holders each with 3 centering screws for aligning with main telescope. 20-mm.-diameter objective. Weighs less than 1/2 pound.

Stock #50,121-Y.....\$8.00 ppd.

STANDARD 1 1/4" EYEPIECE HOLDER



Here is an economical plastic slide-focus eyepiece holder for 1 1/4" O.D. eyepieces. Unit includes 3"-long chrome-plated tube into which your eyepiece fits for focusing. Diagonal holder in illustration is extra and is not included.

Stock #60,067-Y.....\$2.50 ppd.

Stock #60,049-Y
Diagonal holder.....\$1.00 ppd.

EDMUND SCIENTIFIC CO.


THE ORIGINAL GOODWIN RESOLVING POWER BARLOW



For many years, this achromatic coated Barlow lens has been the pride and joy of serious astronomers. Due to the death of the designer, Mr. Frank Goodwin, it has not been available for the past two years. NOW, we can offer these lenses, in exact accordance with the original specifications. Remember, this Barlow is achromatic, coated, mounted in a blackened tube, and as optically perfect as only precision craftsmen can make it. Complete with instructions, in a 4"-long adapter tube for standard 1 1/4" eyepieces.

Stock #60,122-Y.....\$23.50 ppd.

DOUBLE AND TRIPLE YOUR TELESCOPE'S POWER WITH A BARLOW LENS



WHAT IS A BARLOW? A Barlow lens is a negative lens used to increase the power of a telescope without resorting to short focal length eyepieces, and without the need for long, cumbersome telescope tubes. Referring to the diagram above, a Barlow is placed the distance P inside the primary focus of the mirror or objective. The Barlow diverges the beam to a distance Q. This focus is observed with the eyepiece in the usual manner. Thus, a Barlow may be mounted in the same tube that holds the eyepiece, making it very easy to achieve the extra power. The new power of the telescope is not, as you might suppose, due to the extra focal length given the objective by the difference between P and Q. It is defined as the original power of the telescope times the quotient of P divided into Q.



Beautiful chrome mount. We now have our Barlow lens mounted in chrome-plated brass tubing, with special spacer rings that enable you easily to vary the power by sliding split rings out one end and placing them in other end. Comes to you ready to use. Just slide our mounted lens into your 1 1/4" I.D. tubing, then slide your 1 1/4" O.D. eyepieces into our chrome-plated tubing. Barlow lens is nonachromatic.


Stock #30,200-Y...Mounted Barlow lens...\$8.00 ppd.

UNMOUNTED 3X BARLOW LENS

These lenses are made for telescopes that have smaller-diameter eyepieces than the standard 1 1/4" size. Mount one between the eyepiece and objective, and triple your power. Instructions included. Single-element lens, focal length -1.5/16", unmounted.

Stock #30,185-Y...0.932" diam.....\$3.50 ppd.
Stock #30,328-Y...0.912" diam.....\$2.50 ppd.

3X ADJUSTABLE-DIAMETER BARLOW LENS



For telescopes with eyepieces smaller than the standard 1 1/4" outer-diameter size. Prongs on mount can be opened or closed to fit tubes from 13/16" to 1" outer diameter. Directions for using included.

Stock #30,339-Y.....\$5.00 ppd.

MORE POWER FROM YOUR JAPANESE TELESCOPE

Mounted Barlow for Japanese Telescopes

By inserting this single-element lens in the eyepiece end of your Japanese telescope, and putting your regular eyepiece in the end of the Barlow tube, you can increase your telescope's power up to three times. Thus, instead of 60x, you will get 120x or 180x. Barlow is mounted in two pieces of telescoping brass tubing each 4" long, satin chrome plated on the outside. Inner diameter of large tube and outer diameter of small tube are 0.965", the standard size for most Japanese telescopes. Measure yours before ordering. 0.965" is approximately 31/32" or 24.5 mm.

Stock #30,370-Y.....\$6.00 ppd.

SPECIAL! VARIABLE 10- TO 20-POWER U. S. ARMY OBSERVATION TELESCOPE



This is the most versatile and complete government - surplus telescope unit we have ever offered you. USES: spotting scope, surveying, balloon and satellite tracking, all general observation. THE TELESCOPE: Elbow-type, 10 to 20 power continuously variable. Field of view 6° 9' to 3° 5'. 2 1/4" objective lens. Focusing eyepiece. THE MOUNTING: This is really something! Has completely enclosed 8"-diam. circle scale reading directly to 1/10 mil or approx. 20 seconds. (6400 mils equal 360°.) Clutch release allows rapid rotation. Rotates 90° horizontal to vertical with direct reading to 1 mil. Has crossed levels, precision adjustment, and many other exciting features. THE TRIPOD: Made of solid oak with brass fittings. Has cast brass head with four leveling screws. Legs adjustable from 30" to 52" excluding mount height. Extremely sturdy and rigidly cross braced. Instruments used but in good condition. Shipping wt. approx. 100 lbs.

Stock #85,117-Y.....\$89.50 f.o.b. Chicago

CLOCK-DRIVEN EQUATORIAL MOUNT ON PEDESTAL BASE IDEAL FOR YOUR 6" OR 8" REFLECTOR

Accurate electric clock drive and heavy-duty mounting. Operates on household current. Follows stars smoothly. Pedestal is 24" high. Polar-axis shaft diameter 1".

Stock #85,111-Y.....\$74.50 f.o.b. Barrington, N. J.

Same mount and clock drive on 32" hardwood tripod.

Stock #85,081-Y.....\$69.00 f.o.b. Barrington, N. J.

Same mount on metal pedestal, no clock drive.

Stock #85,108-Y.....\$45.00 f.o.b. Barrington, N. J.

Same mount on tripod, no clock drive.

Stock #85,023-Y.....\$39.50 f.o.b. Barrington, N. J.



STARS THAT GLOW IN THE DARK



Transform your den or bedroom into a realistic night sky. Schools, too, can set up authentic sky scenes showing constellations, planets in a dark room for daytime lectures or explanations of earth and space science. Equally attractive as a means of making your children's room a fairyland, where their canopy of stars will make them welcome rather than fight "lights out." Set consists of

125 luminous star decals made of a safe, nonradioactive phosphor. Stars glow after exposure to strong light.

Stock #9298-Y.....\$1.00 ppd.

Mounted Ramsden Eyepieces

Standard 1 1/4" Diameter

Our economy model, standard-size (1 1/4" O.D.) eyepiece. We mounted two excellent quality plano-convex lenses in black anodized aluminum barrels instead of chrome-plated brass to save you money. The clear image you get with these will surprise you. Directions for using short focal length eyepieces are included with both the 1/4" and 1/2" models.

Stock #30,204-Y.....1/4" focal length.....\$4.75 ppd.

Stock #30,203-Y.....1/2" focal length.....\$4.50 ppd.



STAR AND SPACE MAPS

Three giant maps: 1. THE SOLAR SYSTEM, 50" x 38", showing planet statistics; detailed moon map; 12 telephoto pictures of moon, sun, planets. 2. WORLD STAR CHART, 50" x 38"; locates stars for any time of year, any position on earth. 3. MAP OF THE SOLAR SYSTEM, 35" x 48"; shows planets, zodiac, and so on.

Stock #9245-Y....Set of 3 maps.....\$1.25 ppd.

8" SETTING-CIRCLE SET

Stock #50,133-Y...Complete set.....\$3.00 ppd.

Stock #60,078-Y...360° declination circle only \$1.60 ppd.

Stock #60,079-Y...24-hour right-ascension circle only \$1.60 ppd.

5 1/4" SETTING-CIRCLE SET

Stock #50,190-Y...Complete set.....\$2.50 ppd.

Stock #60,080-Y...360° declination circle only \$1.35 ppd.

Stock #60,081-Y...24-hour right-ascension circle only \$1.35 ppd.

LARGE ERFLE EYEPIECE 1 1/2" F.L.

War-Surplus Bargain — Gov't. Cost Approx. \$100

Large telescopes should have one of these for low-power viewing. Apparent field of view 65°. Also use with the 24"-focal-length Aerial Camera lens to make a 16-power wide-field telescope or a 27-power scope with one of the 40"-focal-length Aerial Camera lenses. Low-reflection-coated, 5-element lens system. Field lens of Eastman Kodak's rare-earth glass for better aberration correction. Has diopter scale. Smooth focusing 3/8" movement. Outside diameter of attaching threads, 3" — 32 threads per inch. Clear aperture of eye lens 2", field lens 1-25/32". Weight 3 1/2 lbs.

Stock #50,091-Y.....\$9.95 ppd.

ADAPT GIANT ERFLE TO STANDARD TELESCOPE EYEPIECE SIZE

This adapter lets you use our Giant Erfle with any telescope that has a standard 1 1/4"-diameter eyepiece holder. Of course, you won't get the full field of the Erfle, but you will get a wider field than with a regular eyepiece. Adapter is made of black anodized aluminum tubing approx. 2 3/4" long and 3 3/8" wide.

Stock #50,358-Y.....\$3.95 ppd.

Attention! For Beginners, Junior Astronomy Clubs, Boy and Girl Scouts, School Science Clubs

MAKE 8X ASTRONOMICAL SCOPE with Low-Cost Beginner's Kit!

Every boy and girl of today dreams of being a part of the exploration of outer space. That desire makes building his own telescope a real thrill. Now anyone of Cub Scout or Brownie age on up can make his own astronomical 8-power telescope in one evening, without tools or machinery.



Here is an ideal, attractive group project for scouts, junior astronomy clubs, or similar groups. (See special quantity prices.) Scope is powerful enough to show craters of the moon, Jupiter's satellites, and many stars not visible to the naked eye. Kit includes objective lens, field lens, eye lens, glare stops, kraftboard tubes, cadmium-plated metal ferrules, and other parts to build an 18"-long, 1 1/4"-diameter telescope of 8 power.

Stock #70,238-Y.....\$3.00 ppd.

SPECIAL QUANTITY PRICES

Stock #70,239-Y.... 5 kits.....\$12.50 ppd.

Stock #70,240-Y.... 10 kits.....\$22.50 ppd.

FREE CATALOG "Y"

144 Pages! Over 1000 Bargains!

Fantastic variety — rarely before have so many lenses, prisms, optical instruments, and components been offered from one source. Probably the greatest assembly of bargains in all America. Imported! War Surplus! Hundreds of other hard-to-get optical items. Many science and math learning and teaching aids.

Write for Free Catalog "Y".



ORDER BY STOCK NUMBER . . . SEND CHECK OR MONEY ORDER . . . SATISFACTION GUARANTEED!

BARRINGTON • NEW JERSEY

Precision Diagonals

You will get the best possible performance from your telescope with one of our clear **FUSED QUARTZ** diagonals. Accuracy guaranteed 1/20 wave.

Ellipse 1.25" x 1.77" . . . \$12.00
Ellipse 1.5" x 2.12" . . . \$15.00

PYREX-brand glass diagonals, 1/8-wave accuracy.

Ellipse 1.25" x 1.77" . . . \$ 6.00
Ellipse 1.5" x 2.12" . . . \$ 9.00

Without aluminum coating, deduct \$1.00.

Send for our complete list of supplies, quartz mirrors, blanks, oculars, coatings, and accessories.

E & W OPTICAL CO.

2406 E. Hennepin Ave.
Minneapolis 13, Minn.

THE 4¼" VERNOSCOPE

\$69.50 complete


- ★ 4¼" f/10 **PYREX**-brand glass parabolic primary
- ★ Precision-machined **PAR, Jr.**, Equatorial Mount and portable steel tripod
- ★ Seamless aluminum tube with 3-D space-blue finish
- ★ 4x achromatic finder with crosshairs
- ★ 1¼" helical focusing unit

Free 5" x 7" glossy photo and descriptive brochure available.

VERNOSCOPE & Co.

Candor, New York

TELESCOPE HOBBYISTS



**TEN DAY
MONEY-BACK GUARANTEE**

HOME INSPECTION
of the finest
TELESCOPE PARTS AND KITS

MIRROR MAKING KITS

Build a powerful precision telescope. Quality supplies and instructions. **\$450 UP**

6" MIRROR KIT
with **PYREX**-brand blank **\$9.50**

HARD OVER-COATED ALUMINIZING

6" **\$350** 8" **\$450**

Superior reflecting surface. Guaranteed not to peel or blister.
YOUR MIRROR TESTED FREE ON REQUEST.

Complete Instructions, 10c. Free Catalog.

Precision Optical Supply Co.

1001-T E. 163 St., New York 59, N. Y.

GLEANINGS FOR ATM's

CONDUCTED BY ROBERT E. COX

A PRECISION-MOUNTED SMALL REFRACTOR


SOME TELESCOPE MAKERS build complete instruments including the optics, while others purchase the optics but make all the mechanical parts themselves. A fine instrument with commercial lenses has been made by L. R. Wottring, 974 Mt. Curve Ave., Altadena, Calif., a research and development engineer whose hobby is precision machine work.

A 4½-inch five-element photographic lens of 36" focus forms the main objective. Since it has diaphragm stops of f/8 through f/22, it can be used for astronomical or telephoto-terrestrial photography. In the construction, aluminum was used throughout, except for the studs and bolts, which are stainless steel. The knurled rings on the mounting were especially made to match the one on the main lens of the 35-mm. reflex camera.

A special tool had to be built to form these parts, but Mr. Wottring spared no effort to make his instrument uniform in appearance.

The reflex camera, an Exakta VX IIa, is interchangeable with an adapter for large-field and standard 1¼"-diameter eyepieces. Provision was also made for using an achromatic Barlow lens and either the camera or eyepiece, the highest power being with the Barlow and a ¼-inch eyepiece. Rapid focus is by means of a rack and pinion, while a fine-thread arrangement permits delicate focusing at the eyepiece.

The finder mounted on the main tube has a 2½-inch objective of 16" focus, and with a ¾-inch eyepiece gives a good field for locating celestial objects. Its reticle is illuminated by a red light in order to preserve the dark adaptation of the ob-



L. R. Wottring's carefully constructed mounting has an extra counterweight to balance the heavy objective lens, which has five elements.

server's eye. Quick finding and centering are done with the two ring sights seen just to the left of the finder in the picture. These are coated with luminous paint, which is activated with a flashlight at the start of an evening's observation of the sky.

The polar axle is hollow, containing a built-in optical system and reticle for rapidly aligning the mount on the north celestial pole. The crosshairs are also red-lit, and a penta prism is used as a diagonal to bring the image to a convenient viewing position. In this mounting, latitude adjustment from 0° to 50° is possible. There are five-minute divisions on the hour circle, and the index mark is illuminated by red light. The small oblong box just below the polar-adjustment eyepiece contains a rheostat for varying the intensity of the red lights. On the polar-axis housing is a black knurled knob and flexible shaft for slow-motion control; a similar unit for declination can be seen extending over the main tube and finder.

The counterbalancing system is unusual but necessary. Without this addition, weight of the large lens would put the center of gravity awkwardly close to the upper end of the large tube. The arrangement here not only overcomes this condition but contributes to the balance around the polar axis. The declination circle is divided into 1° units and is also illuminated.

A variable-rate unit is incorporated in the clock drive, so it can be easily set to follow stars, planets, or the moon. The speed is controlled by a governor, adjusted by means of a micrometer screw; this unit monitors a secondary circuit to the drive motor. The drive can use power from either a 12-volt automobile battery or 115-volt alternating current, a transformer and diode rectifier being used in series for a.c. operation. The governor and control system for the drive are mounted on a shelf between the tripod legs. This shelf strengthens the assembly, preventing the legs from spreading outward or twisting when the heavy instrument is rotated.

Large aluminum tripod legs that have adjustable points at their ends to allow leveling are used, so that when the mount is aligned on the pole by means of the polar-axis boresight, the clock will track stars accurately. The north leg (left in picture) has a handle for lifting and steering the assembly when the two legs at the south end of the polar axis are placed on a small two-wheeled carriage, thus allowing easy transportation to and from storage. In actual use, the three legs sit in retaining cups permanently set in a concrete driveway.

R. E. C.

LIGHTWEIGHT BINOCULARS

Improved astronomical observations at low power may be expected in the future by using lightweight 7 x 50 binoculars that have been designed at the Frankford Arsenal by P. R. Yoder, Jr. He describes them, together with a new 6 x 20 binocular for daytime observing, in the Optical Society of America's *Journal* for May, 1960.

The 7 x 50's are of conventional optical arrangement, with Porro prisms, but they incorporate a three-element air-spaced telephoto objective and an eyepiece of three cemented doublets. The speed of f/3 thus obtained permits using small prisms and still maintaining a seven-degree field of view. The over-all optical path length and the diameter of the objective's flint element are also reduced. The general reduction in instrument size, and the use of magnesium metal wherever possible, results in a total instrument weight of only 25 ounces, whereas conventional 7 x 50's weigh 53 ounces. The new instrument is as small and light as ordinary 6 x 30's.

Mr. Yoder's prototype models are undergoing tests by the U. S. Army, with Navy and Air Force trials expected in the near future. A new approach to binocular maintenance has been adopted in their manufacture. The very small 6 x 20's will not be repaired at all — if defective or damaged they are to be thrown away. The 7 x 50's can be mended in the field by replacing one of five unit assemblies, each being preadjusted at the factory.

PARABOLIC MIRRORS OF PYREX-BRAND GLASS

Diam.	Focal Length	Price
6"	48"	\$39.95 postpaid
8"	48"	58.95 postpaid
10"	55"	109.90 postpaid

These mirrors are accurate to 1/4 wave length or better and are free of defects including chips, scratches, and pits. Aluminized reflective coating is protected with quartz overcoating. We invite you to test and compare these with mirrors selling for more. Returnable within 15 days if not pleased.

OVERSTOCKED

1 1/2"-minor-axis elliptical diagonals, aluminized and quartz overcoated. Made of extra-thick fine-annealed PYREX-brand glass. Flat to 1/4 wave or better and free of any defects. While they last. Only \$5.95 each, postpaid.

MOUNTED BARLOW LENS



This is a special simple lens that fits neatly into your 1 3/4" focuser and takes any 1 1/4" O.D. eyepiece. Easily doubles or triples power of your present telescope. No adapters or adjustments necessary. \$7.50 postpaid. Unmounted Barlow lens, 1 1/8" diam. \$3.00 postpaid.

BAUSCH & LOMB RED ROUGE

This comes only in factory-sealed cans and is highly recommended for those who want the best. Prompt shipment from stock.

Packed 5-lb. can, \$8.00 postpaid. WANTED TO BUY: Surplus or discontinued lots of lenses, eyepieces, objectives, and the like.

Write for free catalog.

NYE OPTICAL CO.

7310 Broadway, Lemon Grove, Calif.

DE LUXE

Reflecting Telescope Kits

De luxe kits contain:

- ★ 2 mirror blanks of PYREX-brand glass
- ★ 7 large metal cans of optical abrasives
- ★ Fast-polishing cerium oxide
- ★ Red rouge and pitch

Size	Thickness	Price
4 1/4"	3/4"	\$ 6.00
6"	1"	\$10.50
8"	1 3/8"	\$18.75
10"	1 3/4"	\$33.65
12 1/2"	2 1/8"	\$59.95

ADD POSTAGE: 1st, 2nd, and 3rd postal zones from Detroit, add 10%; 4th, 5th, and 6th, add 15%; 7th and 8th, add 20%; or we will ship postage collect.

Send for free catalog of supplies.

ASHDOWNE BROS.

18450 Grand River Detroit 23, Michigan

"See Thru" STAR MAPS

UNIQUE! Maps of phosphorescent stars printed on transparent plastic.

DIRECT! Look through map at the sky, then match stars for identification.

ACCURATE! All stars to 5th mag. carefully plotted on gnomonic projection.

COMPLETE! 12 maps cover entire sky, all stars designated, plus index map.



Complete set of Maps, \$4.95

BRODER MAPS

1368 McAllister St.
San Francisco 15, Calif.

BRANDON OCULARS



These orthoscopic oculars are the choice of the amateur as well as the professional astronomer. Seven sizes available: 4-mm., 6-mm., 8-mm., 12-mm., 16-mm., 24-mm., and 32-mm. focal lengths. Standard 1 1/4" outside diameter.

\$15.95 each

Brandon Oculars may be purchased at —

Cave Optical Co., 4137 E. Anaheim St., Long Beach 4, Calif.

Cleveland Astronomics, 7618 Lawn Ave., Cleveland 2, Ohio

American Lens and Photo Co., 5700 Northwest Highway, Chicago 30, Ill.

Optron Laboratory, Box 25, D.V. Station, Dayton 6, Ohio

Adler Planetarium, 900 E. Acheson Bond Dr., Chicago 5, Ill.

Polaris Telescopic Shop, 14319 Michigan Ave., Dearborn, Mich.

Chuck's Telescopic-Optical Shop, 887 Sherbourne Dr., Inkster, Mich.

—or order direct. We will ship airmail.

LIBRASCOPE P. R.

Guaynabo

Puerto Rico


Formerly BRANDON INSTRUMENTS



* LAFAYETTE * for * SCIENCE * and * TECHNOLOGY



Professionally Designed and Produced



Polaris

79⁵⁰


2.4" EQUATORIAL REFRACTOR

- 800-mm. focal length
- 62-mm. objective
- Micromotion adjustments on both axes
- Coated optics throughout
- Equatorial mount
- Eyepieces for 160x, 88x, 40x (200x available at extra cost*)

F-342 Shpg. wt. 30 lbs. 5.00 Down Net 79.50
*F-391 4-mm. orthoscopic eyepiece for 200x Net 8.95

Arcturus 3" Equatorial Refractor

269⁵⁰



- Ultra-precise helicoid focusing
- 76.2-mm. clear aperture
- 7 coated eyepieces to 227x
- Most advanced equatorial mount
- 1.6", 500-mm.-f.l. finder

F-385 Shpg. wt. 100 lbs. 10.00 Down Net 269.50

Galactic 3" Equatorial Refractor

189⁰⁰



- 910-mm. focal length
- 76.2-mm. clear aperture
- Micromotion controls for declination and right ascension
- Resolving power 1.6 seconds of arc
- 1.6", 500-mm.-f.l. finder
- 7 eyepieces to 227x

F-384 Shpg. wt. 60 lbs. 5.00 Down Net 189.00



STAR SPECTROSCOPE WITH 3 EYEPIECES

29⁹⁵

- Observe the spectra of stars of magnitudes 1 to 3
- Identify Fraunhofer lines from C to G
- Drawtube focus arrangement
- Fits threads of 25-mm. O.D. eyepieces

F-569 Shpg. wt. 8 oz. 2.00 Down Net 29.95



LAFAYETTE EXCLUSIVE ASTROGRAPHIC PLANETARIUM

9⁹⁵

- Hold the Universe in your hand!
- Completely portable!
- Self-illuminated!
- Locates and identifies all the important constellations seen in our latitudes!
- The perfect star finder!

Shpg. wt. 1 1/2 lbs.

F-500 Net 9.95

LAFAYETTE 14" TRANSPARENT CELESTIAL GLOBE

39⁵⁰

- Basic Aid for Students and Teachers of Astronomy
- Adjustable satellite orbits
- Satellites travel around earth automatically
- Adjustable sun and moon
- Adjustable lunar orbital inclination
- 80 constellations — including all stars of 1st, 2nd, 3rd, and 4th magnitudes

Shpg. wt. 15 lbs.

F-400 5.00 Down Net 39.50



USE OUR EASY PAYMENT PLAN — DOWN PAYMENTS AS LOW AS \$2.00 AND UP TO 24 MONTHS TO PAY

**LAFAYETTE
RADIO**

JAMAICA 33, N. Y.
165-08 Liberty Ave.

NEW YORK 13, N. Y.
100 6th Avenue

BRONX 58, N. Y.
542 E. Fordham Rd.

BOSTON 10, MASS.
110 Federal Street

NEWARK 2, N. J.
24 Central Avenue

PLAINFIELD, N. J.
139 W. 2nd Street

PARAMUS, N. J.
182 Route 17

Lafayette Radio, Dept. SKK-6 P.O. Box 190 Jamaica 31, N. Y.


☐ Please Send Stock #..... I Enclose \$.....

☐ Send me the FREE 324-pg. 1961 catalog 610.

Name.....

Address.....

City..... Zone..... State.....



W. T. Thomas, Jr., took this $\frac{1}{2}$ -second picture of the moon at 1:25 Universal time, August 14, 1959. The very large crater at upper right is Clavius.

AN ALTAZIMUTH REFLECTOR WITH A DRIVE


THE TYPE of altazimuth reflector pictured below has been popular at our amateur club here in Daytona Beach for a number of years. We like the tubeless or beam construction, and find that the telescope's portability makes it ideal for use at meetings and star parties. And the addition of what I call a *steadier* with a pneumatic slow-motion drive allows good photographs of the moon and planets to be taken, with exposure times ranging up to 10 seconds.

The steadier is triangular in shape and supports the telescope beam near the eye end. The result is that a telescope at 360 power will handle as easily and seem as steady as one of our earlier models at 50 power. Eyepieces can be changed at will, without losing the object under observation. The instrument is very little

affected by wind, as the beam does not have the air resistance of a bulky tube, and there can be no internal air currents to affect the seeing.

A telescope with a steadier must first be pointed in the general direction of the object to be viewed. The beam can then be moved nearly two inches either way along the steadier, and up and down a suitable distance until the object is found. For photography, one of the tripod legs is propped upward by means of a wedge-shaped block, so the east-west path of the moon or a planet parallels the direction of the slow-motion movement. It is best, of course, if the field being photographed is within about an hour of the meridian, so the tripod leg is not raised very high.

The first "clock" that I tried was of the



A very low tripod and the vertical adjustable axle in a pipe collar form the mounting of Mr. Thomas' tubeless $\frac{1}{9}$ 6-inch Newtonian reflector. A $1\frac{1}{2}$ " beam of light wood supports the primary mirror and cell at one end and the secondary mirror and eyepiece holder at the other. The "steadier" is made of $\frac{5}{8}$ " dowels, which are seen extending from the two left-hand tripod legs to the slotted upper part of the beam. The entire instrument weighs only $9\frac{1}{2}$ pounds.

TELESCOPE NEWS

VERNONscope & Co. announces — with great pride — the completion of 3", 4", and 6" refracting telescopes, available in f/10 and f/15 models.


Months of extensive research and testing of these American-made instruments have resulted in a design and performance that *you* — the professional and amateur astronomer — desire most in a telescope.

These refractors have actually been modeled to *your* specifications, and we express our sincere gratitude to all those whose suggestions and encouragement have made the construction of these instruments possible.

Even our wildest dream of a 6" refractor telescope produced for less than \$1,000 has at last been fulfilled.

Large glossy illustrations and descriptive brochures are available on request. We shall be honored to serve you.

VERNONscope & Co.
Candor, New York



Nebulosity in Orion captured on 35-mm. film in a ten-minute exposure at f/4, using

The VEGA 6" MAKUTOV

With 4 Erfle and orthoscopic eyepieces, camera adapter, solar filter, case . . . \$1675

Accessories for deep-sky photography also available.

All aluminum and stainless steel, Nylon and teflon bearings, spiral focusing, removable Barlow stage for conversion to f/12.

Length: 24 inches.

Weight: 37 pounds, including pedestal-type equatorial mount and electric drive.

VEGA
INSTRUMENT CO.
840 Lincoln Ave.
Palo Alto, Calif.



Get more telescope for your money ... build the guaranteed 6" CLEVELAND quickly and easily

If you build your own 6" reflector, Cleveland Astronomics can supply and guarantee* every part you need. You'll get three important advantages: first, every Cleveland part is a component of a balanced design, engineered to match and machined to fit; second, you won't need to search out unrelated parts, then cut and fit them to build your telescope; and third, you will own a fine instrument at a substantial saving.

C.A. products are in use throughout the U. S. and as far away as Tasmania. They have been purchased by observatories, colleges, schools, laboratories, and the U. S. Army.

FEATURES: Rotating base permits easy alignment and locking on north. Rotating tube with positive-locking tube clamp. Husky, vise-grip latitude adjustment is easy to set, stays put. Rigid, high-strength aluminum construction. No rust. Extra-long bearings and large brakes give precise control. Legs open to fixed position, close easily for carrying. Parts machined to close tolerance for smooth, trouble-free operation. Easily assembled with simple tools.



EQUATORIAL MOUNT Heavy-duty 12" saddle, tube clamp, axle bearings, rotating base, tripod top, tapered channel legs, and extra pier top are cast aluminum. Axles are 1 1/4" ground and polished steel. Bearings are 5 1/2" long. Large-area brakes, knurled bronze-aluminum alloy adjusting knobs. Brass thrust washers at all friction points. Knurled knobs lock rotating base in V groove in tripod top, release for easy transfer to permanent pier. Latitude adjustment 0° to 55°. Cadmium-plated latitude and leg bolts, nuts and washers. Weight 23 lbs. For 6" telescope **\$79.50 f.o.b.**

For 8" telescope (specify tube O.D.) **89.50 f.o.b.**

COUNTERWEIGHT 12 1/2 lbs., 1" or 1 1/4" bore **5.95 f.o.b.**

EXTRA PIER TOP for permanent mount **5.95 f.o.b.**

ALUMINUM TUBE 7" O.D. x 60" x .064" wall **14.85 f.o.b.**

6" MIRROR CELL Solid plate protects mirror. Ring housing reduces convection currents. Improved clips hold mirror without pressure. Shockproof, cushioned adjustments prevent vibration, keep mirror in collimation. No springs. Cast aluminum, machined for 7" O.D. tube. Attaching screws included **8.35 ppd.**

8" MIRROR CELL Same adjustments as 6" cell, but housing fits inside tube. Specify tube I.D. when ordering **11.95 ppd.**

DIAGONAL HOLDER Fully adjustable, shockproof design. One turn moves mirror .050". Three screw adjustments give perfect control of mirror angle, make collimating easy. For 1 1/4" x 1 1/4" elliptical diagonal mirror. Fits 7" tube **7.95 ppd.**

For tubes up to 10" diam. Specify tube I.D. **9.15 ppd.**

1 1/4" EYEPiece HOLDER Acme-threaded brass sleeve in accurately machined aluminum housing gives smooth, micrometer-sharp focusing. Only one moving part. Stays where you set it **7.95 ppd.**

SETTING CIRCLES 5" cast aluminum with machined faces. 1/2"-wide matte-white scales have accurate black graduations and large, legible figures. 1/2" to 2" bore. Attach with setscrews. Specify shaft size. Pair **12.75 ppd.**

6" MIRROR Pyrex, f/8 parabolic, hand figured to 1/10 wave or better. 48" focal length plus or minus up to 1 1/4" **62.50 ppd.**

DIAGONAL MIRROR 1.250" x 1.770", 1/10 wave **6.50 ppd.**

EYEPieces 1 1/4" Brandon orthoscopic oculars available in seven sizes: 4, 6, 8, 12, 16, 24 and 32 mm. Each **15.95 ppd.**

ALL MECHANICAL PARTS, including setting circles, and complete instructions for building the 6" Cleveland **129.75 f.o.b.**

ALL OPTICAL PARTS Mirror, diagonal, and one Brandon orthoscopic ocular of your choice of focal length **82.45 ppd.**

6" CLEVELAND TELESCOPE including one Brandon eyepiece of your choice of power, assembled, painted, collimated, ready for use and packed for shipment **244.50 f.o.b.**

*We will replace, or refund purchase price of any unsatisfactory product returned prepaid within ten days of receipt. No time limit on return of defective parts.

Add 3% sales tax on all Ohio orders.

7618 LAWN AVE.
CLEVELAND 2, OHIO

CLEVELAND ASTRONOMICS

windup type, but this made varying the speed difficult. Now I use an Austin Craft model-airplane ignition timer (originally \$1.25), which has a pneumatic cylinder and plunger, its speed being controlled by an air-bleed screw. This pneumatic clock is rigged to the steadier where the telescope beam rides on it, and the linkage of levers and wires causes the beam to turn slowly around the vertical axis of the mounting — hence the direction of viewing follows an object across the sky for the period during which the pneumatic clock operates.

By this means I have taken photographs of the moon, Jupiter, and Saturn, in the last case obtaining an 8-second exposure that shows one planetary belt and Cassini's division in the rings. Anyone who wishes detailed sketches of the steadier and drive should write to me.

WILLIAM T. THOMAS, JR.
105 N. Halifax Ave.
Daytona Beach, Fla.

SOME RULES OF THUMB

SEVERAL READERS have asked about the conventional relation between mirror and tube size, and diagonal position and size. Here are some rules of thumb that represent the general experience of amateurs over the years. They may not, however, all apply when special observing conditions are to be met.

Tube size: To allow adequate ventilation inside the tube, it is best to have an extra inch all around the mirror, using an 8" tube for a 6-inch reflector, a 10" tube for an 8-inch, and so on. Many 6-inch instruments, however, work well in 7" tubes.

Diagonal position: Two factors are involved here, the size of the diagonal and the diameter of the telescope tube. To allow for the eyepiece holder, the Newtonian focal plane should be at least 2" outside the tube, making it 7" or more from the optical axis of an 8-inch reflector. If such a mirror is f/8, its focal length is 64" and the diagonal would be placed 57" from the mirror. For a 6-inch f/9, in a 7" tube, the diagonal-to-focal-plane distance should be 5 1/2", placing the secondary 48 1/2" from the primary.

Diagonal size: The formula for this is

$$D = v + C(M - v)/F,$$

where D is the diagonal size in inches (minor axis), v is the diameter of the field of view or focal-plane image, C is the distance the flat is placed inside of focus, M is the primary-mirror diameter, and F its focal length. A good rule-of-thumb assumption is that $v = F/100$.

For an 8-inch instrument, f/8, with C made 7", v is 0.64" and the formula gives nearly 1.5" for D , indicating that the diagonal's minor diameter should be 1 1/2". For the 6-inch, f/9 example above, the result is 1.1".

R. E. C.

From Any Point of VIEWING . . . Quality Means Finer Performance

Enjoy the Professional Features and Exquisite Optics of Any of These 4" Dynascopes®

Check the continuing list of professional-type features offered by these DYNASCOPES®. . . Many exclusive in their respective price field. Notice how carefully each component is finished to the last detail to ensure longer life and a more attractive appearance. Compare the price and see

how any DYNASCOPE makes your dollars go further. All these are reasons why more serious amateurs, colleges, and clubs make Criterion instruments their first choice for superior quality and dependable performance.

4-INCH STANDARD DYNASCOPE

Complete For Only
\$49⁹⁵
f.o.b. Hartford
Shipping Wt. 16 lbs.



The advanced design of this precision-made instrument includes many professional-type features essential for superior viewing — some so new that they were not even available on our more expensive Dynascopes just a year or so ago. Constructed of the finest quality materials, each component is skillfully engineered for dependable operation and fully finished for attractive modern appearance. And, in the true Dynascope tradition, this thrifty 4-inch model brings you all the superior advantages of Criterion's superb optical system. You can order this rich-looking reflector with complete confidence that it will deliver all the performance it promises. Twice-tested to U.S. Government standards before it leaves our factory, it must also meet your test of satisfaction or you may return it for full refund. Send check or money order today for prompt delivery.

• CLASSIC 4-INCH PARABOLIC PYREX MIRROR guaranteed to perform to Dawes' limit for its size. Aluminized and coated with zircon quartz. • QUICK-ACTION EQUATORIAL AND ALTIZMUTH MOUNT with fingertip control. Simple twist locks mount instantly or sets for any drag desired. Free-moving polar axis (small setting circle for declination). • 4-POWER CROSSHAIR FINDERSCOPE with coated achromatic lenses, for greater accuracy, easier spotting. Screw-type micrometer focus. Extra-large field of view. (Supporting bracket with collimation adjustments.) • RACK-AND-PINION EYEPiece FOCUSING for smooth, precision focusing without lash or jerk. • 3 COMPOUND EYEPieces: 18-mm. Huygens, 9-mm. Achromatic Ramsden, 7-mm. Achromatic Ramsden, for 65X, 130X, 167X. Completely threaded precision units. Clear, sharp views to extreme edges. • LIFETIME 45° REINFORCED BAKELITE TUBE with handsome brass trimmings. Fits snugly into fully contoured saddle. Attractive optical-black finish. • TRIPOD OF HARDWOOD folding legs guarantees stability. Special position locks prevent collapse. Completely portable, folds up without disassembling.

SUPER DELUXE ELECTRIC 4-INCH DYNASCOPE

with features
found only in the
finest reflector
telescopes!



Model RV-4
Complete For Only
\$159⁹⁵
f.o.b. Hartford
Shipping Wt. 40 lbs.
Express Chgs. Collect

For those who want the ultimate in an electrically driven 4-inch reflector! This top-performing new DYNASCOPE is actually a 4-inch duplicate of our DeLuxe 6-inch model. You get such truly fine features as a 6 x 30 achromatic finderscope, a fully rotating tube for more comfortable viewing, and a superb optical system that includes five of our finest eyepieces — four achromatic and one orthoscopic. You won't find a finer 4-inch reflector anywhere!

• EXQUISITE HAND-CORRECTED 4-INCH PARABOLIC MIRROR accurate to better than 1/8 wave. Made of pyrex. Aluminized, overcoated. Triple tested. Mounted in removable cell. • ADVANCED-DESIGN EXTRA-HEAVY-DUTY EQUATORIAL MOUNT with oversize castings and fingertip control. Adjustable for latitude. Absolutely smooth action. Rotating base for easier orientation. • SETTING CIRCLES for both right ascension and declination assure faster location of specific object. • SLOW MOTION in declination. • 5 SHARP-VIEW EYEPieces: 18-mm. Achromatic Ramsden, 12.7-mm. Achromatic Ramsden, 9-mm. Achromatic Ramsden, 7-mm. Achromatic Ramsden and 6-mm. Orthoscopic, for 65X, 100X, 130X, 167X, and 220X. All precision-threaded, standard 1 1/4"-diameter barrels. • NEW DYN-O-MATIC ELECTRIC DRIVE, with totally enclosed self-acting clutch. • 6 X 30 ACHROMATIC FINDERSCOPE with crosshairs and extra-firm twin-post mounting. Micrometer focusing, coated achromatic objective. • FULLY ROTATING BAKELITE TUBE with double-thick wall for maximum strength. Polished rings of pure cast aluminum provide extra reinforcement for ends. Smart grey wrinkle finish. • HANDSOME STEEL AND GRAINED-HARDWOOD TRIPOD with special tapered shank ends. Includes spreaders and heavily reinforced joints. No danger of collapse, side-play, or wobble. Sets up quickly anywhere; folds easily and compactly for complete portability.


OTHER DYNASCOPES AVAILABLE IN 3-INCH TO 16-INCH MODELS — Quality engineering of precision instruments does not permit mass production. Consequently supply is limited. For early delivery send your check or money order today, or write for free brochure.

CRITERION MANUFACTURING CO.

Manufacturers of Quality Optical Instruments and Equipment
Dept. STD-60, 331 Church St., Hartford 1, Conn.

DELUXE 4-INCH DYNASCOPE ELECTRIC

with heavy-duty
mount and
new Rotation Rings



Model K-2A
Complete For Only
\$109⁹⁵
f.o.b. Hartford
Shipping Wt. 30 lbs.
Available without
Electric Drive
Model K-2B
\$89.95

A truly scientific instrument with all the features of our standard model plus extra advantages for even finer viewing. Larger and heavier mount (weighs over 12 lbs.) is fully equatorial. Saddle has felt-lined rotation rings that permit tube to be quickly positioned for accessories or to bring eyepiece into more comfortable viewing position.

• EXQUISITE 4-INCH PARABOLIC PYREX MIRROR accurate to 1/8 wave. For famed Dynascope optical superiority. • NEW DYN-O-MATIC ELECTRIC DRIVE, totally enclosed, with self-acting clutch. • TRUE EQUATORIAL AND ALTIZMUTH MOUNT with complete friction adjustments for any latitude, declination and right ascension. Rugged cast-iron construction. • 3 STANDARD 1 1/4" EYEPieces: 18-mm. Huygens, 9-mm. Achromatic Ramsden, 7-mm. Achromatic Ramsden, for 65X, 130X, and 167X. • 4X ACHROMATIC FINDERSCOPE with crosshairs, micrometer focusing and adjustable collimation. • RACK-AND-PINION DOUBLE-DRAW EYEPiece HOLDER for easy, exact adjustments. Accommodates any eyepiece, negative or positive. • PROFESSIONAL-TYPE STEEL TRIPOD with fast-fold legs for easy portability. Extra stability in vibration-free design.



One
big
reason
UNITRON
sales
are
mounting:
the mounting



Equatorial mounting of the 4" UNITRON refractor, Model 152.

Step up, friend. Sit yourself down at the controls of a UNITRON. Feel the sure response of the slow-motion controls. Note the extra-smooth handling ease. Observe the oversize components, which give UNITRON mountings strength to spare.

If it's an altazimuth model, you have micrometric adjustments for both altitude and azimuth. In the equatorial models, you'll find the mechanism more sophisticated—but the operation is even easier. Here, once you've pinpointed your target, only the single right-ascension control is needed to keep the object centered in the field of view. You'll welcome the added convenience of the new flexible cable extension which brings the right-ascension control within easy reach—a

new feature included with all UNITRON equatorials as standard equipment.

Note, too, the precise slow-motion adjustment for declination. Scrutinize the precision gears designed to deliver generations of velvet-smooth performance. Examine the large easy-reading setting circles and verniers provided on the 3" and larger models.

(What you won't see is the instrument's complex of inner workings—a blend of the arts of the artillery engineer and the Swiss watchmaker that gives you an instrument whose response will be obedient to the slightest touch of your fingers.)

And—while you're at it—don't forget to look through the refractor. UNITRON's optical excellence is the other big reason UNITRON sales are mounting.



Sturdy altazimuth mounting with slow-motion controls as well as rapid motions for altitude and azimuth.

UNITRON

INSTRUMENT COMPANY • TELESCOPE SALES DIV. — 66 NEEDHAM ST., NEWTON HIGHLANDS 61, MASS.

SEE THE BACK COVER FOR MORE ABOUT UNITRON.

Does Santa know you want a UNITRON for Xmas?

NEW ACHROMATIC AMPLIFIER NOW FREE with each UNITRON

UNITRON's new "Achromatic Amplifier" is a two-element, Barlow-type, negative amplifying lens designed especially for UNITRON Refractors. When used with a UNITRON eyepiece, it **doubles** the normal magnification of the telescope. Here is an ideal way to increase the usefulness of each eyepiece and to obtain the high magnifications that you have always wanted for planetary and lunar observations. The magnification factor of 2x has been selected as the most useful for general amateur observing under a wide variety of atmospheric conditions.

Two types of mounting cells are offered: one to fit the UNIHEN Rotary Eyepiece Selector and one for the UNITRON Star Diagonal. Either type can be inserted or removed in a jiffy. There are no cumbersome tubes to attach, nor ever any need to adjust the focus of the "Achromatic Amplifier." You merely use your regular standard or accessory UNITRON eyepieces in the UNIHEN or star diagonal in the customary manner. The unique optical design matches the UNITRON objectives and eyepieces to insure sharp definition without loss of resolution.

This useful and valuable accessory is now included as standard equipment with all UNITRON Refractors at no additional cost . . . another of the many reasons why the telescope you choose should be a UNITRON.

The new "Achromatic Amplifier" will be offered as a separate accessory to those who already own UNITRONS. Further details will be announced.

HOW TO ORDER A UNITRON

Send check or money order in full or use our Easy Payment Plan, as described below. Shipments made express collect. Send 20% deposit for C.O.D. shipment. UNITRON instruments are fully guaranteed for quality, workmanship, and performance, and must meet with your approval or your money back.

Use Our EASY PAYMENT PLAN

UNITRON's popular Easy Payment Plan is a convenient and economical way to buy your UNITRON Refractor when you do not want to disturb your savings or when you haven't the total cost of the telescope immediately available. The down payment required is 10%. The balance due is payable over a 12-month period, and there is a 6% carrying charge on the unpaid balance. Your first payment is not due until 30 days after you receive the instrument, and if you should want to pay the entire balance due at that time, the carrying charge is canceled. (6" models available on special plan.)

NOW is definitely the time to reserve a UNITRON for that very special person on your Christmas list who perhaps may be none other than yourself. Precision craftsmanship cannot be hurried and the number of UNITRONS which Santa will have available is necessarily limited. To avoid disappointment, order your UNITRON now. Full payment or the down payment may be made at the time you wish to have the telescope delivered. Remember, there is no substitute for a UNITRON.

MANY Models To Choose From!


1.6" ALTAZIMUTH (\$7.50 Down)	\$75
with eyepieces for 78x, 56x, 39x	
2" SATELLITE (\$7.50 Down) 6x, diagonal eyepiece, altazimuth mount with circles, stand	\$75
2.4" ALTAZIMUTH (\$12.50 Down)	\$125
with eyepieces for 100x, 72x, 50x, 35x	
2.4" EQUATORIAL (\$22.50 Down)	\$225
with eyepieces for 129x, 100x, 72x, 50x, 35x	
3" ALTAZIMUTH (\$26.50 Down)	\$265
with eyepieces for 171x, 131x, 96x, 67x, 48x	
3" EQUATORIAL (\$43.50 Down)	\$435
with eyepieces for 200x, 131x, 96x, 67x, 48x	
3" PHOTO-EQUATORIAL (\$55.00 Down)	\$550
with eyepieces for 200x, 171x, 131x, 96x, 67x, 48x	
4" ALTAZIMUTH (\$46.50 Down) with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x	\$465
4" EQUATORIAL (\$78.50 Down) with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x	\$785
4" PHOTO-EQUATORIAL (\$89.00 Down) with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x	\$890
4" EQUATORIAL with clock drive (\$98.50 Down), Model 160V, eyepieces as above	\$985
4" EQUATORIAL with clock drive and metal pier (\$107.50 Down), Model 166V, eyepieces as above	\$1075
4" PHOTO-EQUATORIAL with clock drive and Astro-camera (\$117.50 Down), with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x, 25x	\$1175
4" PHOTO-EQUATORIAL with clock drive, pier, Astro-camera (\$128.00 Down), with 10 eyepieces	\$1280

Note These UNITRON Features

- LENSES are FULLY CORRECTED for spherical and chromatic aberration and are COATED for maximum brilliance and clarity of image. Optics especially designed for astronomical observation. AIR-SPACED OBJECTIVES.
- REFRACTOR type of design duplicates the performance of larger telescopes of other types. No mirrored surfaces to become oxidized. Superior definition to the very edge of the field.
- EYEPIECES of the HIGHEST QUALITY: Orthoscopic, Achromatized Symmetrical, Kellner, Huygens. Three to ten eyepieces included with each instrument.
- FINEST MATERIALS used throughout. DURALUMIN TUBE. Moving parts of BRASS carefully machined to close tolerances, and finished in CHROMIUM. No surplus components used.
- MODERN DESIGN based on time-tested engineering principles. HANDSOME APPEARANCE to which no illustration can do justice.
- EQUATORIAL MODELS have slow-motion controls for both declination and right ascension as well as convenient clamps on both axes. Sturdy TRIPOD (or PIER).
- ALTAZIMUTH MODELS have slow-motion controls for both altitude and azimuth as well as clamps for both co-ordinates. Sturdy TRIPOD.
- VIEW FINDER with crosshair eyepiece.
- RACK-AND-PINION FOCUSING.
- Choice of UNIHEN Rotary Eyepiece Selector or STAR DIAGONAL and ERECTING PRISM for TERRESTRIAL OBSERVATION. The same complete range of terrestrial magnifications as for celestial observation.
- SUNGLASS for solar observation.
- ADDITIONAL ACCESSORIES available to add further to your observing pleasure.

Higher- and lower-priced eyepieces available for all models. Prices include basic accessories, tripod and mounting, fitted wooden cabinets, and operating instructions.

New UNITRON View Finders



UNITRON's popular view finders with newly designed optics and mechanical features are better than ever. From left to right: 23.5 mm., 30 mm., 42 mm.

1. VIEW FINDER (Used on UNITRON 2.4" Equatorials): 23.5-mm. (.93") achromatic objective, 6x eyepiece with crosshairs. Chromed brass tube. Mounting brackets with centering screws. **Only \$8.50 postpaid**

2. VIEW FINDER (As used on UNITRON 3" Refractors): 30-mm. (1.2") coated achromatic objective and 8x eyepiece with crosshairs. Other details as in View Finder 3. **Only \$10.75 postpaid**

3. VIEW FINDER (As used on UNITRON 4" Refractors): 42-mm. (1.6") coated achromatic air-spaced objective. 10x eyepiece with crosshairs. Duralumin tube finished in white enamel. Dewcap. Furnished with mounting brackets, centering screws for collimation, and mounting screws. This finder measures approximately 16" overall. It is light in weight, compact and small enough for use as a hand telescope furnishing spectacular wide-field views of the sky. **Only \$18.00 postpaid**

UNITRON

Only \$18.00 postpaid
INSTRUMENT COMPANY
TELESCOPE SALES DIVISION
66 NEEDHAM ST.
NEWTON HIGHLANDS 61, MASS.

Get UNITRON's FREE Observer's Guide and Catalog on ASTRONOMICAL TELESCOPES

This valuable 38-page book
is yours for the asking!

With artificial satellites already launched and space travel almost a reality, astronomy has become today's fastest growing hobby. Exploring the skies with a telescope is a relaxing diversion for father and son alike. UNITRON's handbook contains full-page illustrated articles on astronomy, observing, telescopes and accessories. It is of interest to both beginners and advanced amateurs.

Contents include—

- Observing the sun, moon, planets and wonders of the sky
- Constellation map
- Hints for observers
- Glossary of telescope terms
- How to choose a telescope
- Amateur clubs and research programs



UNITRON

INSTRUMENT COMPANY • TELESCOPE SALES DIV.
66 NEEDHAM ST., NEWTON HIGHLANDS 61, MASS.

Please rush to me, free of charge, UNITRON's new Observer's Guide and Telescope Catalog 20-H.

Name _____
Street _____
City _____ State _____

CELESTIAL CALENDAR


Universal time (UT) is used unless otherwise noted.

THE ECLIPSING VARIABLE STAR RZ CASSIOPEIAE

MANY years ago, the German astronomer K. Graff wrote: "Of all the bright variable stars of the Algol type, apart from Beta Persei itself, RZ Cassiopeiae is undoubtedly the easiest to observe. Its circumpolar location, its readily identifiable field on a line from Epsilon Cassiopeiae through Iota, its large amplitude and great maximum brightness, its short period — all these give the star a special significance, and I can scarcely name another object better suited for visual brightness estimates with small instruments."

The variability of RZ Cassiopeiae was discovered in 1906 at Potsdam Observatory by G. Müller, and it was soon found to be an eclipsing binary star with a period of 1.195 days. Most of the time it is at maximum brightness, magnitude 6.4. When primary eclipse begins, RZ Cassiopeiae takes only two hours to fade to magnitude 7.8, and two hours more to recover its full light. The general character of the changes is well illustrated by the accompanying light curve, formed from 575 visual estimates made by N. Florja during the years 1923-33. It even indicates the shallow secondary minimum, only 0.05 magnitude deep, which occurs midway between the deep primary eclipses.

A particularly interesting property of RZ Cassiopeiae is the marked variation in the length of its period. During the last decade the period has gradually shortened, and the minima are currently occurring nearly half an hour earlier than the predictions from a formula that was correct



This light curve of RZ Cassiopeiae was compiled by E. Slossim from visual observations by N. Florja. Adapted from the Soviet journal "Variable Stars," Vol. 5 (1937).

in 1953. From visual estimates of brightness, the amateur can determine the time of minimum light to within a few minutes. Hence, such determinations are useful for tracing the unpredictable fluctuations in period.

To observe RZ Cassiopeiae, no more equipment is needed than 7x50 binoculars, a finder telescope, or a 2-inch refractor with a low-power eyepiece. Beginning about one or 1½ hours before the scheduled time of minimum, estimate the magnitude of RZ Cassiopeiae in terms of the comparison stars in the accompanying chart. These estimates are to be continued at roughly 10- or 15-minute intervals until one or 1½ hours after midminimum. The time of each observation should be recorded to the nearest minute.

An important precaution in observing RZ Cassiopeiae, or any other rapidly varying star, is to avoid bias. Each estimate should be as nearly independent of those before as is possible, and the observer should not be influenced by the scheduled time of minimum (which in any case is uncertain).

The following predicted times of primary eclipses include only those occurring during hours of darkness for North America. They are expressed in Universal time, in hours and tenths. To obtain predictions of other minima, add to the times given, or subtract, multiples of 1 day 4.69 hours.

October 26, 3^h.1; 27, 7.8. November 1, 2.5; 2, 7.2; 7, 1.9; 8, 6.6; 13, 1.4; 14, 6.0; 19, 0.8; 20, 5.5; 21, 10.2; 25, 0.2; 26, 4.9; 27, 9.6. December 2, 4.3; 3, 9.1; 8, 3.8; 9, 8.5; 14, 3.2; 15, 7.9; 20, 2.6; 21, 7.3; 26, 2.0; 27, 6.7.

This ephemeris is based on new elements that satisfy the minima observed from 1953 to 1960.

Readers who make observations of RZ Cassiopeiae are invited to send a transcript of their estimates to SKY AND TELESCOPE for analysis and evaluation of the times of minimum.

TRANSIT OF MERCURY

Every observer in the Americas should circle Monday, November 7th, on his calendar. On that date Mercury will pass across the face of the sun. In Central and South America, and in the eastern and middle parts of North America, both the beginning and end of the spectacle will be visible. For watchers on the West Coast and in the Northwest, however, the transit will be in progress when the sun rises.

This five-hour event begins at 14:34 Universal time (8:34 a.m., Central standard time) and ends at 19:12 UT (1:12 p.m. CST), these times being valid to within a minute or two throughout the Western Hemisphere. The planet comes onto the sun's disk at position angle 148° (southeast quadrant) and exits at 262° (near the west point of the sun).

Mercury will appear as a round, black, very small dot, its angular diameter of 9.9 seconds of arc being only 1/200 the sun's. Though too small to be detected without optical aid, the planet in transit is readily observable in a 2- or 3-inch telescope. Adequate precautions should be taken to protect the observer's eyes from the blinding glare of the sun. One safe method is to project an enlarged image of the sun through the eyepiece onto a white screen (see page 283).

For detailed observing information, consult the article beginning on page 214 of last month's SKY AND TELESCOPE.

VARIABLE STAR MAXIMA

November 9, T Cephei, 210868, 6.0; 11, RU Cygni, 213753, 8.0; 13, V Monocerotis, 061702, 7.0; 14, R Reticuli, 043263, 7.6; 21, T Eridani, 035124, 8.0.

December 11, T Centauri, 133633, 5.5.

These predictions of variable star maxima are by the AAVSO. Stars are listed only if brighter than magnitude 8.0 at an average maximum. Some, but not all of them, are nearly as bright as maximum two or three weeks before and after the dates for their maxima. The data given include, in order, the day of the month near which the maximum should occur, the star name, the star designation number, which gives the rough right ascension (first four figures) and declination (bold face if southern), and the predicted visual magnitude.

MINIMA OF ALGOL


November 1, 17:35; 4, 14:24; 7, 11:12; 10, 8:01; 13, 4:50; 16, 1:39; 18, 22:28; 21, 19:17; 24, 16:06; 27, 12:55; 30, 9:44.

December 3, 6:33; 6, 3:22; 9, 0:11; 11, 21:00; 14, 17:49.


These minima predictions for Algol are based on the formula in the 1953 *International Supplement of the Krakow Observatory*. The times given are geocentric; they can be compared directly with observed times of the star's least brightness.

UNIVERSAL TIME (UT)

TIMES used in Celestial Calendar are Greenwich civil or Universal time, unless otherwise noted. This is 24-hour time, from midnight to midnight; times greater than 12:00 are p.m. Subtract the following hours to convert to standard times in the United States: EST, 5; CST, 6; MST, 7; PST, 8. If necessary, add 24 hours to the UT before subtracting, in which case the result is your standard time on the day preceding the Greenwich date shown. For example, 6:15 UT on the 15th of the month corresponds to 1:15 a.m. EST on the 15th, and to 10:15 p.m. PST on the 14th.



North is toward the top in this finder chart for RZ Cassiopeiae, and one degree is about 1.1 inches. Comparison stars are labeled with their magnitudes to the nearest tenth, the decimal points being omitted. In the same field is the Cepheid SU Cassiopeiae, which varies between magnitudes 5.9 and 6.4 in a period of 1.95 days.



THE SUN, MOON, AND PLANETS THIS MONTH

The sun, on the ecliptic, is shown for the beginning and end of the month. The moon's symbols give its phase roughly, with the date marked alongside. Each planet is located for the middle of the month or for other dates shown. All positions are for 0^h Universal time on the respective dates.

Mercury arrives at inferior conjunction with the sun on November 7th, its transit across the solar disk being visible in Europe, Africa, the Americas, and the Pacific Ocean (see facing page). The planet then moves into the morning sky, and reaches greatest western elongation, 20° from the sun, on the 24th. The week preceding and the week following that date are good for observing this -0.3-magnitude object, for Mercury rises about 1½ hours before the sun and may be seen low in the sky to the south of east.

Venus on the 15th is a prominent evening star of magnitude -3.5 in Sagittarius, setting about two hours after the sun. Telescopically, the planet appears as a gibbous disk 79-per-cent illuminated and 14" in diameter. Venus will overtake first Jupiter and then Saturn on the evenings of the 18th and 27th, respectively. In both cases, Venus passes some two degrees south.

Mars is stationary in right ascension on November 21st and then begins retrograde (westward) motion among the stars. At this time the reddish planet rises about three hours after sunset and is of magnitude -0.6. In a telescope its disk can be seen, 13".1 in angular diameter.

Jupiter in the middle of the month is low in the southwestern sky for nearly 2½ hours after sunset, in Sagittarius, its magnitude being -1.5. On the 15th the oblate disk has an equatorial diameter of 33".3 and a polar diameter of 31".1.

Saturn is approximately 12° east of Jupiter and sets almost 3½ hours after the sun in midmonth. Throughout November its magnitude is +0.8. In a telescope the rings are readily visible, being 35".3 across on the 15th, while the planet's equatorial diameter is 15".7. The moon will pass some 4° north of Saturn on the morning of the 22nd.

Uranus is at western quadrature on

November 22nd, when it rises about an hour before midnight, local time, and crosses the meridian about ¾ hour before sunrise. On the 15th this 6th-magnitude planet is at right ascension 9^h 52^m.2, declination +13° 41' (1950 co-ordinates), and should be easily located with binoculars 1½° northwest of Nu Leonis (magnitude 5.2).

Neptune is in conjunction with the sun on November 1st, and hence not observable this month.

WILLIAM H. GLENN

MINOR PLANET PREDICTIONS

Psyche, 16, 9.5. November 12, 4:55.1 +17:49; 22, 4:47.1 +17:31. December 2, 4:38.1 +17:14; 12, 4:29.1 +17:00; 22, 4:21.2 +16:52. January 1, 4:15.4 +16:51. Opposition on December 3rd.

Laetitia, 39, 10.1. November 12, 5:37.7 +7:53; 22, 5:31.4 +7:21. December 2, 5:23.2 +7:01; 12, 5:14.0 +6:55; 22, 5:05.1 +7:05. January 1, 4:57.4 +7:29. Opposition on December 11th.

Harmonia, 40, 10.2. November 22, 5:48.8 +21:07. December 2, 5:39.4 +21:21; 12, 5:28.1 +21:35; 22, 5:16.5 +21:49. January 1, 5:06.2 +22:02; 11, 4:58.7 +22:14. Opposition on December 14th.

After the asteroid's name are its number and the approximate visual magnitude expected at opposition. At 10-day intervals are given its right ascension and declination (1950.0) for 0^h Universal time. In each case the motion of the asteroid is retrograde. Data are supplied by the IAU Minor Planet Center at the University of Cincinnati Observatory.

MOON PHASES AND DISTANCE

Full moon	November 3, 11:58
Last quarter	November 11, 13:48
New moon	November 18, 23:47
First quarter	November 25, 15:42
Full moon	December 3, 4:25
November	Distance Diameter
Apogee 9, 9 ^h	251,600 mi. 29' 31"
Perigee 21, 4 ^h	225,700 mi. 32' 54"
December	
Apogee 7, 3 ^h	252,100 mi. 29' 27"

NOVEMBER METEORS


Two rather sparse meteor showers occur this month. The full moon will seriously interfere with observations of the Taurid shower maximum on November 5th, but watchers should be able to see some Taurid meteors for a fortnight before and after that date. The Leonid shower reaches maximum on the morning of the 16th, before new moon. At that time a single observer may see some 15 meteors per hour; about half of these are associated with the shower. W. H. G.

JUPITER'S SATELLITES


The curves in the accompanying chart show the positions of Jupiter's four bright moons, as seen in an inverting telescope, with north at the bottom and east at the right. Each horizontal line is for 0^h Universal time on the date specified, and the intersections of the line with the curves indicate the places of the satellites at that time. For other Universal times, interpolate between the 0^h lines. The double vertical lines represent the planet's disk.

The lower section is intended to aid observations of the eclipses of Jupiter's moons; *d* is the point of disappearance of the satellite in Jupiter's shadow, *r* is the point of reappearance. The chart is from *The American Ephemeris and Nautical Almanac*; for further explanation, see page 446 of *SKY AND TELESCOPE* for May, 1960.

JUPITER'S SATELLITES IN NOVEMBER



PHASES OF THE ECLIPSES




You'll find these fully professional features in every Custom DYNASCOPE®!

Exquisite optical systems, fully achromatic and corrected to meet the most critical research standards • Massive equatorial mounts that provide exceptional stability and embody the latest mechanical refinements for ease of operation • Full sets of matched orthoscopic and achromatic eyepieces with wide flat fields sharp to extreme edges • Needle bearings carefully fitted on substantial shafts to assure movements so fine that they equal the ease and precision of a fine watch • Lifetime materials throughout, for atmospheric resistance and dimensional stability • Fully rotating tubes, with counterpoise to provide perfect balance during rota-

tion • Rotary secondary supports to minimize diffraction • Exclusive Dyna-mesh rack and pinion focus for smoother, more accurate focusing action • Two-way pier tripods (Pat. Pending) in 6-inch and 8-inch models, 45-pound pier can be mounted permanently. Portable tripod is instantly removable for field trips • Highly accurate electric drives, free from backlash, with fully compensating clutch • 8 x 50 coated achromatic finderscopes with crosshairs and extra-large 6° fields • 7"-diameter setting circles spoked for easy setting. Graduated with verniers • Super sensitive slow motion flex line control • Adjustable to all latitudes.

MASTERPIECES *of precision performance!*



All sizes of these superb instruments are available for unusually prompt delivery ... including this Deluxe 12.5-inch model

CUSTOM DYNASCOPES®

in your
choice of
6"-8"-10"
12.5"-16"
models

Every Custom DYNASCOPE is designed and built to meet such high standards and exacting specifications that it is the overwhelming choice of schools, colleges, universities, and many well-known professional astronomers. Each of these magnificent instruments incorporates features that are years ahead in optical engineering, yet based on time-tested principles. Each is painstakingly produced and assembled by craftsmen who put careful workmanship before

speed. Each is required to pass, flawlessly, a long series of rigid performance and operational tests. Nothing is overlooked to make sure that these superb telescopes are the finest in their class, regardless of price. Whichever size you select, we guarantee that it *must* measure up to this standard or your purchase price will be refunded in full. You can order with absolute confidence...and with the exciting expectation of new thrills and experiences ahead.



Colorful New Brochure Yours FREE!

Send today for this fascinating new booklet that gives you the full specifications on all Custom Dynascopes, plus helpful information to guide you in the selection of your next telescope. A postcard will bring your copy FREE!

CUSTOM DYNASCOPES—PRICE AND DELIVERY SCHEDULE


CONVENIENT TIME PAYMENT PLANS AVAILABLE ON ALL MODELS									
SIX-INCH		EIGHT-INCH		10-INCH	12.5-INCH	16-INCH			
STANDARD	DELUXE	STANDARD	DELUXE	DELUXE	DELUXE	OBSERVATORY MODEL			
\$265 Complete	\$475 Complete	\$385 Complete	\$595 Complete	\$895 Complete	\$1,265 Complete	\$4,200 up			
Shipping Weight 80 lbs.	Shipping Weight 150 lbs.	Shipping Weight 145 lbs.	Shipping Weight 185 lbs.	Shipping Weight 245 lbs.	Shipping Weight 300 lbs.	Equipped To Your Specifications			
21 Day Delivery		21 Day Delivery		30 Day Delivery	30 Day Delivery				


Standard Six and Eight Inch Models Do Not Include Electric Clock Drive, Setting Circles, Slow Motion Control Or Pier
All Telescopes Are Shipped F.O.B. Hartford, Connecticut • No Additional Charge For Packing and Crating

© Copyright 1960 Criterion Mfg. Co.

® TM Registered U.S. Pat. Office

CRITERION MANUFACTURING CO. *Manufacturers of Quality Optical Instruments*
DEPT. D-43, 331 CHURCH ST., HARTFORD 1, CONN.





NISHIMURA 6-INCH, 10-INCH, and 8-INCH NEWTONIAN REFLECTORS

All Nishimura telescopes described here are manufactured to meet the exacting standards of the professional observatory. Biela Observatories, Inc., presents Nishimura astronomical instruments with the assurance that you will be satisfied with the quality and workmanship. If an instrument does not meet these specifications to your satisfaction, the full purchase price will be refunded.

OPTICAL SPECIFICATIONS

6-inch: focal length, 54 inches; $f/9$
 8-inch: focal length, 64 inches; $f/8$
 10-inch: focal length, 80 inches; $f/8$

Optical accuracy: $\frac{1}{8}$ -wave sodium light or better.

Mirror and diagonal ground on back and edges for precise positioning; both mirrors flotation mounted.

Full complement of eyepieces: 4-mm., 6-mm., 9-mm., 12.5-mm., 18-mm., 25-mm., 40-mm. Give the following magnifications: 6-inch, 33x to 333x; 8-inch, 40x to 400x; 10-inch, 50x to 500x.


Finderscope:

6-inch: 7x achromatic finder, 1.2-inch objective, 5-degree field.
 8-inch: 8x achromatic finder, 1.6-inch objective, 5-degree field.
 10-inch: 7x achromatic finder, 2-inch objective, 6-degree field.

The 3-inch refracting guide telescope is standard equipment on the 8- and 10-inch reflectors (not illustrated on 8-inch). Focal length,

45 inches, $f/15$, mounted on main tube, with prism-type star diagonal. Four eyepieces of your choice.

All eyepieces, including one universal solar filter, furnished in a neat wooden case for convenience and protection. The eyepieces fit both the reflector and auxiliary refractors.



ENGINEERING SPECIFICATIONS

Heavy cast-aluminum tripod for the 6-inch reflector; massive cast-iron pier for 8- and 10-inch, with adjustable double pier plate for precise polar alignment.

Cast-iron German-type equatorial mounting, adjustable polar axis with graduated latitude indicator.

Brass setting circles with vernier reading (no vernier on the 6-inch reflector), accurately engraved for right ascension and declination.

110-volt, 60-cycle, sidereal-rate electric clock drive with heavy-duty motor.

Manual slow-motion controls on 6- and 8-inch reflectors; electric slow-motion controls on the 10-inch. These controls operate in right ascension and declination independently of the sidereal-rate electric clock drive.

Positive clamps on both right ascension and declination axes.

Steel tube with machined cast-iron end rings, precision-machined rotation saddle.

Precision rack-and-pinion eyepiece holder with three sizes of adapter rings.

Four independent counterweights on threaded shaft for accurate balancing.

All castings and tubes are finished in high-quality enamel. Castings steel grey, tubes off-white.

All hardware chrome plated.

Metal covers provided for all optical surfaces as added protection.

Total net weight: 6-inch, 90 lbs.; 8-inch, 284 lbs.; 10-inch, 490 lbs.

Prices, f.o.b. Anaheim, Calif.: 6-inch reflector, \$495.00
 8-inch reflector, \$995.00
 10-inch reflector, \$1,495.00

Biela Observatories, Inc.

325 BERRY AVENUE, ANAHEIM, CALIFORNIA. PHONE: KEystone 3-2207


DISTRIBUTOR FOR NISHIMURA ASTRONOMICAL INSTRUMENTS

NOW...

a complete line of Astro-Science Equipment *
available from ONE source!

A complete line of Astro-Science Equipment is available from ONE source... a great advance in the history of astronomical education.

Now you can obtain from just one distributor all the major equipment necessary to establish an astronomy department in your school, college, or community. A complete line of planetarium projectors and astronomical telescopes has been added to the Astro-Dome line through our new association with the Goto Optical Manufacturing Co.



**GOTO S1 and S2
PLANETARIUM
PROJECTORS**

Designed and priced for modest installations, in domes 16 to 30 feet in diameter.

- 1 **Optical Projection.** Like their big relatives, the Goto Medium-Size and the Goto Large projectors, S1 and S2 have optical lens projection systems throughout, producing sharp star and planet images on modest-sized domes.
- 2 All three essential motions of the projector are obtained with motor drives. Therefore, the daily motion (rising and setting) of the stars may be reproduced; the great change of precession of the equinoxes (a 25,800-year cycle); and change of latitude, permitting imaginary voyages **completely** around the world in either direction.
- 3 The celestial equator, ecliptic, and Milky Way are projected.
- 4 The supporting stand is heavy and rigid, and designed with a horizontal east-west axis, as it is in all large planetariums.
- 5 Setting rings for the diurnal and precessional motions.
- 6 Individual projectors for the planets, sun, and moon.



**GOTO
6-INCH (15-CM.)
EQUATORIAL**

A large refractor with the finest features, ready for immediate installation on its heavy metal pier. Equipped with setting circles and slow-motion controls—a versatile clock-driven equatorial.


OPTICS: The achromatic objective has a clear aperture of 150 millimeters, f/15, focal length 2,250 mm. It resolves double stars to about $\frac{1}{4}$ second of arc and detects stars fainter than magnitude $12\frac{1}{2}$. Seven eyepieces are provided, as well as a finder telescope, of 50-mm. aperture and 12x magnification.

ACCESSORIES: Star diagonal, Herschel prism, two sun filters, a moon filter, star spectroscopy, and accessories case.

FINISH: Instrument finished in white and light gray.

PATENTS PENDING

THE ALL-NEW ASTRO-DOMETTE




Astro-Domettes have been developed for high schools and junior colleges. These units are fabricated on an assembly line basis, resulting in high quality yet modest cost. Because sections are stamped or molded, dome components are uniform and interchangeable. Astro-Domettes are available in sizes of 8, 10 and 12 feet outside diameters. In the space age, progressive schools will want to include this new aid to education in their science departments. Domettes are available in pastel colors.

ASTRO-DOME
INCORPORATED

1801 BROWNLEE AVE., N. E. • CANTON 5, OHIO
GLendale 6-8361

Johnny isn't watching the late show tonight...



The shadows that shiver and shake on the TV screen are shivering and shaking in somebody else's living room tonight. Johnny has discovered something new.

He's traded the fleeting, flickering "thrills" of the 24 inch screen for the timeless excitement and majesty of the night sky.

He's traded the nervous rattle of the private eye's gun for a ringside seat at the stupendous nightly fireworks in the heavens.

He has, in short, discovered astronomy.

Nothing better could happen than what happened to Johnny. And it happened simply because someone took the trouble to awaken, nourish and satisfy a lifetime of curiosity in Johnny by making him the gift of a fine telescope.

Someone, not so long ago, gave Johnny a Unitron.

Johnny abandoned his 24" screen for this 2.4" Unitron refractor, complete with its handy, handsome, easily portable carrying case.



This is a close-up of the Unihex Johnny is using. It's 6 eyepieces in one, an exclusive with Unitron. One of a complete line of accessories to multiply your viewing pleasure.

UNITRON

INSTRUMENT COMPANY • TELESCOPE SALES DIV. — 66 NEEDHAM ST., NEWTON HIGHLANDS 61, MASS.

SEE PAGES 308 AND 309 FOR MORE ABOUT UNITRON.

